RESEARCH ARTICLE

REVISED B cell, CD8⁺ T cell and gamma delta T cell infiltration alters alveolar immune cell homeostasis in HIV-infected Malawian adults [version 2; peer review: 1 approved, 2 approved with reservations]

Previously titled: B cell, CD8⁺ T cell and gamma delta T cell lymphocytic alveolitis alters alveolar immune cell homeostasis in HIV-infected Malawian adults

Andrew Mwale¹, Annemarie Hummel ¹, Leonard Mvaya¹, Raphael Kamng'ona¹, Elizabeth Chimbayo¹, Joseph Phiri ¹, Rose Malamba¹, Anstead Kankwatira¹, Henry C Mwandumba ^{1,2}, Kondwani C Jambo ^{1,2}

¹Malawi-Liverpool-Wellcome Trust Clinical Research Programme, University of Malawi College of Medicine, Blantyre, Malawi

²Department of Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, UK

v2 First published: 27 Oct 2017, 2:105
<https://doi.org/10.12688/wellcomeopenres.12869.1>
 Second version: 21 Dec 2017, 2:105
<https://doi.org/10.12688/wellcomeopenres.12869.2>
 Latest published: 06 Apr 2018, 2:105
<https://doi.org/10.12688/wellcomeopenres.12869.3>

Abstract

Background: HIV infection is associated with increased risk to lower respiratory tract infections (LRTI). However, the impact of HIV infection on immune cell populations in the lung is not well defined. We sought to comprehensively characterise the impact of HIV infection on immune cell populations in the lung.

Methods: Twenty HIV-uninfected controls and 17 HIV-1 infected ART-naïve adults were recruited from Queen Elizabeth Central Hospital, Malawi. Immunophenotyping of lymphocyte and myeloid cell populations was done on bronchoalveolar lavage fluid and peripheral blood cells.

Results: We found that the numbers of CD8⁺ T cells, B cells and gamma delta T cells were higher in BAL fluid of HIV-infected adults compared to HIV-uninfected controls (all $p < 0.05$). In contrast, there was no difference in the numbers of alveolar CD4⁺ T cells in HIV-infected adults compared to HIV-uninfected controls ($p = 0.7065$). Intermediate monocytes were the predominant monocyte subset in BAL fluid (HIV-, 63%; HIV+ 81%), while the numbers of classical monocytes was lower in HIV-infected individuals compared to HIV-uninfected adults (1×10^5 vs. 2.8×10^5 cells/100ml of BAL fluid, $p = 0.0001$). The proportions of alveolar macrophages and myeloid

Open Peer Review

Approval Status

	1	2	3
version 3 (revision) 06 Apr 2018			 view
version 2 (revision) 21 Dec 2017	 view		 view
version 1 27 Oct 2017	 view	 view	

1. **David H. Dockrell**, University of Edinburgh, Edinburgh, UK

2. **Graeme Meintjes**, University of Cape Town, Cape Town, South Africa

Muki Shey , University of Cape Town, Cape Town, South Africa

dendritic cells was lower in HIV-infected adults compared to HIV-uninfected controls (all $p < 0.05$).

Conclusions: Chronic HIV infection is associated with broad alteration of immune cell populations in the lung, but does not lead to massive depletion of alveolar CD4⁺ T cells. Disruption of alveolar immune cell homeostasis likely explains in part the susceptibility for LRTIs in HIV-infected adults.

Keywords

BAL, HIV, B cell, CD8 T cell, gamma-delta T cells, CD4 T cell, alveolitis, adult

3. Dawn M.E. Bowdish , McMaster

University, Hamilton, Canada

Any reports and responses or comments on the article can be found at the end of the article.

This article is included in the [Malawi-Liverpool Wellcome Trust Clinical Research Programme gateway](#).

Corresponding author: Kondwani C Jambo (kjambo@mlw.mw)

Author roles: **Mwale A:** Conceptualization, Data Curation, Formal Analysis, Investigation, Writing – Original Draft Preparation, Writing – Review & Editing; **Hummel A:** Conceptualization, Data Curation, Formal Analysis, Investigation, Writing – Original Draft Preparation, Writing – Review & Editing; **Mvaya L:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Kamng'ona R:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Chimbayo E:** Investigation, Writing – Original Draft Preparation, Writing – Review & Editing; **Phiri J:** Data Curation, Investigation, Writing – Original Draft Preparation, Writing – Review & Editing; **Malamba R:** Conceptualization, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Kankwatira A:** Conceptualization, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Mwandumba HC:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Jambo KC:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Wellcome Trust [105831]; and the Bill and Melinda Gates Foundation [OPP1125279].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2017 Mwale A *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Mwale A, Hummel A, Mvaya L *et al.* **B cell, CD8⁺ T cell and gamma delta T cell infiltration alters alveolar immune cell homeostasis in HIV-infected Malawian adults [version 2; peer review: 1 approved, 2 approved with reservations]** Wellcome Open Research 2017, 2:105 <https://doi.org/10.12688/wellcomeopenres.12869.2>

First published: 27 Oct 2017, 2:105 <https://doi.org/10.12688/wellcomeopenres.12869.1>

REVISED Amendments from Version 1

The following changes have been made in this version 2:

1. The title has been revised based on the reviewers' comments
2. It contains more details regarding the study participants
3. It includes further details about the bronchoscopy procedure
4. We have corrected errors in [Figure 1](#) and [Figure 5](#)
5. The Discussion section has been expanded
6. We have also corrected an error in the [Supplementary Table 1](#)

See referee reports

Introduction

HIV-infected individuals have increased susceptibility to lower respiratory tract infections (LRTIs)^{1,2}, which account for 75–98% of lung complications in antiretroviral therapy (ART)-naïve HIV-infected adults worldwide^{3,4}. Predisposition to LRTIs is largely attributed to HIV-induced impairment of lung immunity, including reduced frequency of respiratory antigen-specific alveolar CD4⁺ T cells^{5–7} as well as impaired alveolar macrophage function^{5,8}. HIV infection is also associated with CD8⁺ T cell alveolitis, a condition characterized by the influx of HIV-specific CD8⁺ T cells into the lung^{9,10}. While these immune cell perturbations partly underlie propensity for LRTIs in HIV-infected individuals, the impact of HIV infection on the composition and functions of other immune cell populations in the lung is not well defined.

Several studies have reported alterations in the proportions and functions of different immune cell populations in peripheral blood in HIV-infected individuals^{11–14}. While peripheral blood CD4⁺ T cell depletion and an increase in CD8⁺ T cells are hallmarks of progressive untreated chronic HIV infection¹⁵, depletion of B cells¹¹ and aberrant NK cell function and redistribution from CD56^{dim} towards CD56^{neg} subsets has been observed during early and chronic HIV infection¹². Two major human $\gamma\delta$ T cells subsets (designated V δ 1 or V δ 2) are also altered in HIV-infected individuals, with an increase in the V δ 1 subset and a decrease in the V δ 2 subset¹³. Furthermore, increased proportions of non-classical and intermediate monocytes and depleted myeloid and plasmacytoid dendritic cell subsets have been reported in individuals with high plasma HIV viral load^{14,16,17}.

We, therefore, undertook a comprehensive characterisation of the impact of HIV infection on immune cell populations in the lung. We obtained paired bronchoalveolar lavage (BAL) fluid and peripheral blood from HIV-uninfected and asymptomatic HIV-infected, antiretroviral therapy (ART)-naïve Malawian adults. We analysed and compared the proportions and numbers of CD4⁺ and CD8⁺ T cells, B cells, NK cell subsets, $\gamma\delta$ T cells, monocytes, dendritic cell subsets, neutrophils and alveolar macrophages in samples from HIV-infected and uninfected individuals.

Methods

Study participants

The study was conducted at the Queen Elizabeth Central Hospital, a large teaching hospital in Blantyre, Malawi. Participants were recruited from the hospital's voluntary counselling and testing (VCT) and ART clinics. They were adults aged ≥ 18 years

comprising healthy HIV-1-uninfected and asymptomatic HIV-1-infected volunteers with no clinical evidence of active disease and willing to undergo bronchoscopy and BAL for research purposes¹⁸. Clinical diagnosis and radiographic examination were used to exclude individuals with active disease, including TB, from the study. No latent TB tests were carried out on the participants. The study participants were recruited from the same catchment area, with relatively similar environmental exposures. HIV testing was performed on whole blood using two commercial point-of-care rapid HIV test kits, Determine HIV 1/2 kit (Abbott Diagnostic Division) and Unigold HIV 1/2 kit (Trinity Biotech Inc.). A participant was considered HIV-uninfected if the test was negative by both kits or HIV-infected if the test was positive by both kits. If Determine and Unigold results were discordant, a third rapid test using Bioline HIV 1/2 kit (Standard Diagnostics Inc.) was performed to resolve the discordance. No baseline viral loads were done on the HIV-infected individuals. However, none of the participants were on ART at the time of recruitment to the study, but all initiated ART after sample collection according to the 'test and treat' Malawi national treatment guidelines. Exclusion criteria for the study were: current or history of smoking, use of immunosuppressive drugs, severe anaemia (Hb < 8g/dl) and known or suspected pregnancy. The research ethics committee of Malawi College of Medicine approved the study under approval number P.03/16/1907 and all participants provided written informed consent.

Sample collection and experimental procedures

Bronchoscopy and BAL were performed on all participants as previously described^{5,6,8}. Briefly, we used Instillagel to lubricate the nostrils and 2% lignocaine for vocal cords and the airways. A fiber-optic bronchoscope (Olympus, UK) was passed to the level of a sub-segmental bronchus of the right middle lobe and four 50 ml aliquots of sterile normal saline at 37°C instilled and removed using gentle hand suction. A typical BAL return from a 200ml instill is 100–140ml. The aspirated bronchoalveolar lavage fluid was placed into 50ml falcon tubes and transferred immediately to the laboratory for processing within 30 min. The fluid was filtered using sterile gauze and centrifuged at 500g for 10min. The supernatant was removed, the cell pellet was resuspended and washed with PBS by spinning in a centrifuge at 500g for 10min. The supernatant was removed and discarded while the cell pellet was resuspended in complete media. Upon microscopic examination, the cell pellet contained <5% bronchial epithelial cells or squamous cells. Due to time taken from BAL isolation and sample acquisition, there is a potential for changes in cell viability, especially neutrophils, however, the a standardised study protocol was applied to both study groups to minimise the time from isolation to acquisition. Peripheral blood was also obtained from study participants for full blood count (FBC) and peripheral blood mononuclear cell (PBMC) isolation using density gradient centrifugation. Cell counts in BAL cells and PBMCs isolated from each sample were performed using a haemocytometer.

Immunophenotyping

Whole BAL cells (1×10^6 cells) and PBMCs (1×10^6 cells) were stained with predetermined optimal concentration of fluorochrome-conjugated monoclonal antibodies against human cell surface proteins. Two separate antibody panels targeting lymphocytic and myeloid cells were used. The lymphocyte

panel consisted of anti-CD3 PE/Cy5, anti-CD4 Bv421, anti-CD8 APC-Cy7, anti-CD19 PE, anti-CD56 APC, anti-TCR $\gamma\delta$ FITC, and anti-CD45 PE-CF594. The myeloid panel consisted of anti-CD45 PE-CF594, anti CD14 Bv421, anti-CD16 PE/Cy7 PC7, anti-HLADR PE/Cy5, anti-CD66 FITC, anti-CD206 APC, anti-CD11c APC/Cy7 and anti-CD123 Bv510. Further details of the antibodies are in [Supplementary Table 1](#). All samples were analysed using a BD LSRFortessa flow cytometer (Becton Dickinson, USA).

Statistical analysis

Statistical analyses and graphical presentation were performed using GraphPad Prism 5 (GraphPad Software, USA). We used FlowJo v10 software (Treestar, USA) to analyse flow cytometry data. The numbers of cell subsets in BAL fluid were estimated by calculating the proportion of a particular subset relative to the total number (1×10^6 cells) of stained cells. The cell count was first calculated as number of cells relative to the BAL volume return of each individual, and this was then standardised to cells per 100ml BAL fluid as previously published^{5,6}. In PBMCs, the absolute numbers were obtained by calculating the proportion of a particular subset relative to the full blood count (FBC) data, specifically, lymphocyte and monocytes counts. Data were analysed using Mann Whitney U test. Results are given as median and interquartile range (IQR). Differences were considered statistically significant when $p < 0.05$.

Results

Study participants and samples

We recruited 20 HIV-uninfected healthy controls (median age [range] (32[18-52]; male:female, 12:8) and 17 asymptomatic

HIV-infected adults (median age [range] (33 [24-58]; male:female, 8:9). The CD4 count (median [range]) was lower in HIV-infected adults compared to the HIV-uninfected controls (365[218-541]) vs. 731[541-888] cells/ μ l, $p = 0.0024$). The main characteristics of the participants are summarised in [Table 1](#). Not all experimental assays were performed on all study participants.

CD8⁺ T cells, B cells and $\gamma\delta$ T cells contribute to HIV-associated lymphocyte infiltration in the alveolar space

We investigated the impact of HIV infection on the proportion and numbers of lymphocyte populations using flow cytometry. The gating strategy is illustrated in [Figure 1](#). We found that the proportions and numbers of lymphocytes in BAL fluid were higher in HIV-infected adults compared to HIV-uninfected (median 20.8% vs. 8.5%, $p = 0.0004$ and median 1×10^7 vs. 2.7×10^6 cells/100ml of BAL fluid, $p = 0.0005$, respectively) ([Figure 2A](#) and [2B](#)). We next determined the cell subsets that were responsible for the increased frequency of lymphocytes in

Table 1. Demographics of the study participants.

	HIV-uninfected controls (n=20)	HIV-infected ART-naive (n=17)
Age (years), median (range)	32(18-52)	33(24-58)
Sex (M:F)	12:8	8:9
CD4 count (cells/ μ l), median (IQR)	731(541-888)	365(218-541)

Figure 1. Representative flow cytometry plots for characterising lymphocytes in BAL fluid from HIV-uninfected adult. BAL cells were stained with fluorochrome-conjugated antibodies.

Figure 2. Proportions and numbers of CD4⁺ T cells, CD8⁺ T cells and CD19⁺ B cells in BAL fluid from ART-naïve HIV-infected compared to HIV-uninfected individuals. BAL cells were stained with fluorochrome-conjugated antibodies. **A)** Proportion of lymphocytes in BAL fluid. **B)** Numbers of lymphocytes in BAL fluid. **C)** Proportion of CD4⁺ and CD8⁺ T cells in BAL fluid. **D)** Numbers of CD4⁺ and CD8⁺ T cells in BAL fluid. **E)** Proportion of B cells in BAL fluid. **F)** Numbers of CD19⁺ B cells in BAL fluid. The horizontal bars represent median and 95% confidence intervals. Data were analyzed using Mann Whitney U test. (HIV-, n=20; HIV+ ART-, n=17).

the alveoli. We found that the proportions and numbers of CD8⁺ T cells (median, 68% vs. 32%, $p < 0.0001$ and median 7×10^6 vs. $7 \times 10^5/100\text{ml}$ of BAL fluid, $p < 0.0001$, respectively) and B cells (median 1.8% vs. 0.8%, $p = 0.0014$ and median 7×10^4 vs. $1 \times 10^4/100\text{ml}$ of BAL fluid, $p < 0.0001$, respectively) in BAL fluid were higher in HIV-infected adults compared to HIV-uninfected controls (Figure 2C–2F). The proportion and numbers of $\gamma\delta$ T cells were also higher in BAL fluid from HIV-infected adults compared HIV-uninfected controls (median 1.4% vs. 0.8%, $p = 0.036$ and median 1×10^5 vs. $2 \times 10^4/100\text{ml}$ of BAL fluid, $p = 0.0002$, respectively) (Figure 3A and 3B).

In contrast, the proportions of CD4⁺ T cells and NK cells in BAL fluid were lower in HIV-infected adults compared to HIV-uninfected controls (CD4⁺ T cell, median 2% vs. 4%, $p < 0.0001$; NK cells, median 1% vs. 2%, $p < 0.0001$) (Figure 2C and Figure 3C). However, the numbers showed no difference in CD4⁺ T cells (median 1.1×10^6 vs. $1.0 \times 10^6/100\text{ ml}$ of BAL fluid, $p = 0.7065$) and NK cells (median 5.4×10^4 vs. $4.9 \times 10^4/100\text{ml}$ of BAL fluid, $p = 0.8911$) between HIV-infected adults and

HIV-uninfected controls (Figure 2D and Figure 3D). Furthermore, there is no statistically significant difference between the alveolar CD4 T cell count in individuals with peripheral blood CD4 count of less than 350 compared to those with a peripheral blood CD4 count of greater than 350 (Median [Interquartile range] 1.9×10^6 [$0.9\text{--}2.2 \times 10^6$] vs. 0.9×10^6 [$0.4\text{--}2.2 \times 10^6$]/100ml of BAL fluid, $p = 0.1905$). These findings demonstrate that HIV infection has a differential impact on alveolar lymphocyte populations.

Differential impact of HIV infection on lymphocyte subsets in the alveolar and blood compartments

We then investigated the similarities and differences of HIV-associated changes in cell composition between BAL fluid and peripheral blood. In agreement with BAL fluid, the proportions of CD8⁺ T cells in peripheral blood were higher in HIV infected adults compared to HIV-uninfected controls (Median 47% vs. 24%, $p < 0.0001$) (Supplementary Figure 1). The proportions of CD4⁺ T cells in peripheral blood were lower in HIV-infected adults compared to HIV-uninfected controls (Median 20% vs. 46%, $p < 0.0001$) (Supplementary Figure 1). In contrast with BAL fluid,

Figure 3. Proportions and numbers of $\gamma\delta$ T cells and NK cells in BAL fluid from ART-naïve HIV-infected compared to HIV-uninfected individuals. BAL cells were stained with fluorochrome-conjugated antibodies. **A)** Proportion of $\gamma\delta$ T cell subsets in BAL fluid. **B)** Numbers of $\gamma\delta$ T cell subsets in BAL fluid. **C)** Proportion of NK cell subsets in BAL fluid. **D)** Numbers of NK cell subsets in BAL fluid. The horizontal bars represent median and 95% confidence intervals. Data were analyzed using Mann Whitney U test. (HIV-, $n=20$; HIV+ ART-, $n=17$).

the proportion of B cells in peripheral blood was lower in HIV-infected adults compared to HIV-uninfected controls (Median 5.8% vs. 9.4%, $p=0.0472$) (Supplementary Figure 1). The proportion of CD3⁺CD56⁺ NK T cells in peripheral blood was lower in HIV-infected adults compared to HIV-uninfected controls (Median 0.03% vs. 0.09%, $p=0.0386$) (Supplementary Figure 1). The proportion of CD8 $\gamma\delta$ T cells in peripheral blood was higher in HIV-infected adults compared to HIV-uninfected controls (Median 1.9% vs. 0.76%, $p=0.0229$) (Supplementary Figure 1). The findings show that HIV infection differentially impacts lymphocyte populations in the alveolar space and peripheral blood compartments.

Differential impact of HIV infection on monocyte subsets in the alveolar and blood compartments

Next, we investigated the impact of HIV infection on the monocyte population in BAL fluid compared to peripheral blood. The gating strategy is illustrated in Figure 4. First we determined the composition of the monocyte cell population in BAL fluid in comparison to peripheral blood. We found that irrespective of HIV status CD14⁺ CD16⁺ intermediate monocytes were the

predominant subset in BAL fluid, followed by CD14⁺⁺CD16^{lo} classical monocytes and then CD14^{lo}CD16⁺ non-classical monocytes (HIV-, Median 63% vs. 33% vs. 5%; HIV+, Median 81% vs. 13% vs. 9%) (Figure 5A and 5C). In blood, irrespective of HIV status, CD14⁺⁺ CD16^{lo} classical monocytes were the predominant monocyte subset, followed by CD14^{lo}CD16⁺ non-classical monocytes and then CD14⁺CD16⁺ intermediate monocytes (HIV-, median 74% vs. 18% vs. 9%; HIV+, median 73% vs. 23% vs. 8%) (Figure 5B and 5D).

Second, we compared the proportions and numbers of monocyte population in BAL fluid and peripheral blood between HIV-infected adults and HIV-uninfected controls. In BAL fluid, we found that the proportion and numbers of CD14⁺CD16^{lo} classical monocytes were lower in HIV-infected adults compared to HIV-uninfected controls (median 13% vs. 33%, $p=0.0002$ and median 1×10^5 vs. 2.8×10^5 cells/100ml of BAL fluid, $p=0.0001$, respectively) (Figure 5C and 5E). In contrast, the proportion of CD14⁺CD16⁺ intermediate monocytes was higher in HIV-infected adults compared the HIV-uninfected controls (median,

Figure 4. Representative flow cytometry plots for characterising myeloid cells in BAL fluid from an HIV-uninfected adult. BAL cells were stained with fluorochrome-conjugated antibodies.

Figure 5. Proportions and numbers of monocyte subsets in BAL fluid and peripheral blood from ART-naïve HIV-infected compared to HIV-uninfected individuals. BAL cells and PBMCs were stained with fluorochrome-conjugated antibodies. **A)** Flow cytometry representative plot of stained BAL sample from an HIV-uninfected control. **B)** Flow cytometry representative plot of stained peripheral blood sample from an HIV-uninfected control. **C)** Proportion of monocyte subsets in BAL fluid. **D)** Proportion of monocyte subsets in peripheral blood. **E)** Numbers of monocytes subsets in BAL fluid. **F)** Numbers of monocyte subsets in peripheral blood. The horizontal bars represent median and 95% confidence intervals. Data were analyzed using Mann Whitney U test. (BAL fluid, HIV-, n = 20; HIV+ ART-, n = 17; PBMC, HIV-, n = 16; HIV+ ART-, n = 14).

80% vs. 64%, $p=0.0011$) but the numbers were similar between the two groups (median 6.0×10^5 vs. 7.7×10^5 cells/100ml of BAL fluid, $p=0.8628$) (Figure 5C and 5E). In blood, we found that the numbers of CD14⁺ CD16^{lo} classical monocytes (median 110 vs. 60 cells/1000 mm³, $p=0.0237$), CD14⁺CD16⁺ intermediate monocytes (median 20 vs. 6 cells/1000 mm³,

$p=0.0362$) and CD14^{lo}CD16⁺ non classical monocytes (median 30 vs. 10 cells/1000 mm³, $p=0.0316$) were higher in HIV-infected adults compared to HIV-uninfected controls (Figure 5F). These findings underscore differences in the composition and the impact of HIV infection on immune cells in the lung and systemic compartments.

Altered proportions of alveolar macrophages and dendritic cell populations in HIV-infected adults

Lastly, we investigated the impact of HIV on alveolar macrophages (AM), neutrophils and dendritic cell populations in BAL fluid. We found that the proportions of alveolar macrophages and myeloid dendritic cells were lower in HIV infected adults compared to HIV-uninfected controls (AM, median 73% vs. 80%, $p=0.0109$; mDC, median 0.6% vs. 0.9%, $p=0.0036$) (Figure 6A and 6C). The proportion of neutrophils and plasmacytoid

dendritic cells was similar between HIV-infected adults and HIV-uninfected controls (neutrophils, median 0.34% vs. 0.14%, $p=0.0789$; pDC, median 0.04% vs. 0.05%, $p=0.1947$) (Figure 6A and 6C). The numbers of alveolar macrophages (median 2.0×10^6 vs. 2.2×10^6 cells/100ml of BAL fluid, $p=0.8628$), neutrophils (median 6.2×10^4 vs. 8.4×10^4 cells/100ml of BAL) and dendritic cells (mDC, median 2.1×10^5 vs. 2.7×10^5 cells/100ml of BAL fluid, $p=0.2676$; pDC, median 1.3×10^5 vs. 1.7×10^5 cells/100ml of BAL fluid, $p=0.5328$) were

Figure 6. Proportions and numbers of alveolar macrophages, neutrophils and dendritic cells in BAL fluid from ART-naïve HIV-infected compared to HIV-uninfected individuals. BAL cells were stained with fluorochrome-conjugated antibodies. **A)** Proportion of alveolar macrophages and neutrophils in BAL fluid. **B)** Numbers of alveolar macrophages and neutrophils in BAL fluid. **C)** Proportion of dendritic cell subsets in BAL fluid. **D)** Numbers of dendritic cell subsets in BAL fluid. The horizontal bars represent median and 95% confidence intervals. Data were analyzed using Mann Whitney U test. (HIV-, $n = 20$; HIV+ ART-, $n = 17$).

similar between the HIV-infected adults and HIV-uninfected controls (Figure 6B and 6D). Taken together, the findings show that chronic HIV-infection is associated with a disruption in the homeostatic proportions of alveolar macrophage and dendritic cell populations.

Discussion

We report the broad impact of HIV infection on immune cell populations in the alveolar space beyond the well-characterised CD8⁺T cell alveolitis observed in previous studies^{5,6,9,10}. We show that in addition to CD8⁺ T cells, B cells and $\gamma\delta$ T cells are increased, while classical monocytes are decreased in BAL fluid from ART-naïve HIV-infected adults compared to HIV-uninfected individuals. We further show generalised disruption in the proportions of immune cell subsets including alveolar macrophages, CD4⁺ T cells, myeloid dendritic cells, intermediate monocytes and NK cells in BAL fluid of asymptomatic chronic HIV-infected adults.

Although HIV-infection was associated with accumulation of B cells and $\gamma\delta$ T cells in BAL fluid, their contribution to pulmonary immunity during chronic HIV infection is incompletely understood. However, previous studies have reported HIV-associated impairment of function of these two cell subsets in peripheral blood^{19–21}. Consistent with what has been observed in the systemic circulation, hyperglobulinemia has been reported in BAL fluid of HIV-infected adults^{22,23}, but the antibodies have impaired opsonic function²⁴. It is plausible that the HIV-associated increase in B cells in the lung results in increased antibody production and BAL fluid hyperglobulinemia. Furthermore, the increase in $\gamma\delta$ T cells that we found in the present study supports the findings of Agostini *et al.*²⁵, who showed that HIV-infected individuals with CD8⁺ T cell alveolitis had increased $\gamma\delta$ T cells in BAL fluid, which were predominantly of the V δ 2 subset. However, HIV infection was also associated with anergic $\gamma\delta$ T cells that were characterised by their substantially deficient response to phosphoantigens^{26–28}. Taken together, the findings of previous studies lead us to postulate that despite the increase in numbers, lung B cells and $\gamma\delta$ T cells from HIV-infected individuals have impaired function as their blood counterparts.

HIV infection is associated with massive depletion of mucosal CD4⁺ T cells in the gut^{29,30} and gradual decline in peripheral blood CD4⁺ T cells¹⁵. We have shown preserved mucosal CD4⁺ T cells in BAL fluid from chronic HIV-infected adults, even in those with depleted peripheral blood CD4⁺ T cells. Our findings are consistent with previous work that showed lung CCR5⁺CD4⁺ T cells are not massively depleted during HIV infection³¹. However, it is not clear whether these preserved mucosal CD4⁺ T cells in BAL fluid would also be true in individuals with symptomatic chronic HIV infection. The mechanisms behind this preservation of alveolar CD4⁺ T cells is unclear and warrants further investigation. However, Mahlknecht *et al.* has shown that macrophages can prevent CD4⁺ T cell apoptosis *in vitro* via cell to cell contact using a mechanism that involves

stimulation of *nef*-expressing CD4⁺ T cells with macrophage membrane-bound TNF³². *Nef* in presence of TNF stimulation promotes activation of anti-apoptotic transcription factor NF- κ B, resulting in blockade of caspase-8 activation and subsequent apoptosis³². It is therefore plausible that alveolar macrophages could promote survival of CD4⁺ T cells in the lung through similar mechanisms, but this warrants further investigation. However, although alveolar CD4⁺ T cells are not massively depleted during chronic HIV infection, their functional capacity is perturbed^{5–7}.

Consistent with others^{33–35}, we have showed that CD16⁺CD14⁺ intermediate monocytes were the predominant subset in BAL fluid. The lower proportion of CD14⁺CD16⁺ classical monocytes in BAL compared to peripheral blood is consistent with work from Baharom *et al.*, which comprehensively characterized monocyte subsets and their function in the lung mucosa³⁶. CD16⁺ monocytes and AM have been shown to be permissive to HIV infection^{8,37}. The abundance of intermediate monocytes and AM in BAL fluid increases potential cellular targets for HIV. Our findings that AM are preserved during chronic HIV infection, may partly be attributed to the long life span of these cells^{38,39}, as well as their resistance to the cytopathic effects of HIV^{40,41}. In contrast, we observed a depletion in classical monocytes in BAL fluid from HIV-infected individuals. The mechanism for the selective depletion of classical monocytes is unclear, but might involve HIV-induced apoptosis⁴² or loss/downregulation of surface CD14⁴³. In steady state, alveolar macrophages originate from erythro-myeloid progenitors (EMPs), while monocytes originate from haematopoietic stem cells (HSCs)⁴⁴, hence the differential impact of HIV on these subsets might be due to the distinct nature of their source of origin. On the other hand, during an inflammatory state, classical monocytes are thought to differentiate into lung macrophages and contribute to clearance of invading pathogens⁴⁵. Presence of a wide array of HIV-permissive cells in the lung, including recruited and resident cells, could contribute to maintenance of local viral production and subsequent disruption of immune cell populations and homeostasis in this compartment.

A potential limitation of the study is that the numbers of BAL cell subsets are extremely difficult to measure with a very high degree of accuracy due to the variations in the dilution factor of epithelial lining fluid and differences in BAL fluid volume return. However, using a method utilised in previous studies^{5,6}, we calculated numbers of cell subsets using the BAL cell count obtained from a haemocytometer combined with proportions obtained by immunophenotyping. We have confidence in the reliability of this method to measure the numbers for the other cell subsets, as we have replicated the observation that the absolute number of CD8⁺ T cells is higher in HIV-infected adults compared with HIV-uninfected individuals^{5,6,9,10}. Furthermore, using data from our previous work that focused on measuring cytokines in BAL fluid, we found no statistically significant

difference in concentration of urea in BAL fluid between HIV-infected adults compared to HIV-uninfected individuals, suggesting that permeability of the alveolar space might not be different in the two groups (unpublished).

In conclusion, our findings show that HIV infection is associated with broad alteration of immune cell populations in the lung. Disruption in immune homeostasis has been shown to lead to increased susceptibility to both infectious and non-infectious diseases. The broad alteration of immune cell populations in the lung in part explain the propensity to LRTI in HIV-infected individuals. However, the degree to which successful anti-retroviral therapy restores the composition of immune cells in the lung warrants further investigation.

Data availability

The data underlying the results presented in this manuscript are available from OSF: osf.io/ykve4.

Competing interests

No competing interests were disclosed.

Grant information

This work was supported by the Wellcome Trust [105831]; and the Bill and Melinda Gates Foundation [OPP1125279].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

The authors thank all study participants, Mrs Kunkeyani (MLW, Blantyre, Malawi), Mrs Kanyandula (MLW, Blantyre, Malawi) and staff of MLW and QECH for their support and co-operation during the study. We thank Prof David Russell (Cornell University, Ithaca, USA) for proof reading the manuscript.

Supplementary material

Supplementary Figure 1. Proportions of CD4⁺ T cells, CD8⁺ T cells, CD19⁺ B cells, $\gamma\delta$ T cells and NK in peripheral blood from ART-naïve HIV-infected compared to HIV-uninfected individuals. PBMC were stained with fluorochrome-conjugated antibodies. A) Proportion of CD4⁺ and CD8⁺ T cells in peripheral blood. B) Proportion of B cells in peripheral blood. C) Proportion of $\gamma\delta$ T cell subsets in peripheral blood. D) Proportion of NK cell subsets in peripheral blood. The horizontal bars represent median and 95% confidence intervals. Data were analyzed using Mann Whitney U test. (HIV-, n=16; HIV+ ART-, n=14).

[Click here to access the data.](#)

Supplementary Table 1. Details of fluorochrome-conjugated antibodies used in the study.

[Click here to access the data.](#)

References

- Martinson NA, Hoffmann CJ, Chaisson RE: **Epidemiology of tuberculosis and HIV: recent advances in understanding and responses.** *Proc Am Thorac Soc.* 2011; **8**(3): 288–293.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Segal LN, Methé BA, Nolan A, *et al.*: **HIV-1 and bacterial pneumonia in the era of antiretroviral therapy.** *Proc Am Thorac Soc.* 2011; **8**(3): 282–287.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Murray JF: **Pulmonary complications of HIV-1 infection among adults living in Sub-Saharan Africa.** *Int J Tuberc Lung Dis.* 2005; **9**(8): 826–35.
[PubMed Abstract](#)
- Benito N, Moreno A, Miro JM, *et al.*: **Pulmonary infections in HIV-infected patients: an update in the 21st century.** *Eur Respir J.* 2012; **39**(3): 730–745.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jambo KC, Banda DH, Afran L, *et al.*: **Asymptomatic HIV-infected individuals on antiretroviral therapy exhibit impaired lung CD4⁺ T-cell responses to mycobacteria.** *Am J Respir Crit Care Med.* 2014; **190**(8): 938–947.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jambo KC, Sepako E, Fullerton DG, *et al.*: **Bronchoalveolar CD4⁺ T cell responses to respiratory antigens are impaired in HIV-infected adults.** *Thorax.* 2011; **66**(5): 375–382.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kalsdorf B, Scriba TJ, Wood K, *et al.*: **HIV-1 infection impairs the bronchoalveolar T-cell response to mycobacteria.** *Am J Respir Crit Care Med.* 2009; **180**(12): 1262–1270.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jambo KC, Banda DH, Kankwatira AM, *et al.*: **Small alveolar macrophages are infected preferentially by HIV and exhibit impaired phagocytic function.** *Mucosal Immunol.* 2014; **7**(5): 1116–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Neff CP, Chain JL, MaWhinney S, *et al.*: **Lymphocytic alveolitis is associated with the accumulation of functionally impaired HIV-specific T cells in the lung of antiretroviral therapy-naïve subjects.** *Am J Respir Crit Care Med.* 2015; **191**(4): 464–473.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

10. Twigg HL, Soliman DM, Day RB, *et al.*: **Lymphocytic alveolitis, bronchoalveolar lavage viral load, and outcome in human immunodeficiency virus infection.** *Am J Respir Crit Care Med.* 1999; **159**(5 Pt 1): 1439–1444.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Moir S, Buckner CM, Ho J, *et al.*: **B cells in early and chronic HIV infection: evidence for preservation of immune function associated with early initiation of antiretroviral therapy.** *Blood.* 2010; **116**(25): 5571–5579.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Mavilio D, Lombardo G, Benjamin J, *et al.*: **Characterization of CD56/CD16⁺ natural killer (NK) cells: a highly dysfunctional NK subset expanded in HIV-infected viremic individuals.** *Proc Natl Acad Sci U S A.* 2005; **102**(8): 2886–2891.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Poles MA, Barsoum S, Yu W, *et al.*: **Human immunodeficiency virus type 1 induces persistent changes in mucosal and blood gammadelta T cells despite suppressive therapy.** *J Virol.* 2003; **77**(19): 10456–10467.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Funderburg NT, Zidar DA, Shive C, *et al.*: **Shared monocyte subset phenotypes in HIV-1 infection and in uninfected subjects with acute coronary syndrome.** *Blood.* 2012; **120**(23): 4599–4608.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Cao W, Mehrav V, Kaufmann DE, *et al.*: **Elevation and persistence of CD8 T-cells in HIV infection: the Achilles heel in the ART era.** *J Int AIDS Soc.* 2016; **19**(1): 20697.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Pacanowski J, Kahi S, Baillet M, *et al.*: **Reduced blood CD123⁺ (lymphoid) and CD11c⁺ (myeloid) dendritic cell numbers in primary HIV-1 infection.** *Blood.* 2001; **98**(10): 3016–3021.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Donaghy H, Pozniak A, Gazzard B, *et al.*: **Loss of blood CD11c⁺ myeloid and CD11c⁺ plasmacytoid dendritic cells in patients with HIV-1 infection correlates with HIV-1 RNA virus load.** *Blood.* 2001; **98**(8): 2574–2576.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Mtunthama N, Malamba R, French N, *et al.*: **Malawians permit research bronchoscopy due to perceived need for healthcare.** *J Med Ethics.* 2008; **34**(4): 303–307.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Moir S, Malaspina A, Pickeral OK, *et al.*: **Decreased survival of B cells of HIV-viremic patients mediated by altered expression of receptors of the TNF superfamily.** *J Exp Med.* 2004; **200**(5): 587–599.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Moir S, Ho J, Malaspina A, *et al.*: **Evidence for HIV-associated B cell exhaustion in a dysfunctional memory B cell compartment in HIV-infected viremic individuals.** *J Exp Med.* 2008; **205**(8): 1797–1805.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Pauza CD, Poonia B, Li H, *et al.*: **γδ T Cells in HIV Disease: Past, Present, and Future.** *Front Immunol.* 2014; **5**: 687.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Fahy RJ, Diaz PT, Hart J, *et al.*: **BAL and serum IgG levels in healthy asymptomatic HIV-infected patients.** *Chest.* 2001; **119**(1): 196–203.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Collins AM, Batrawy SE, Gordon SB, *et al.*: **Increased IgG but normal IgA anti-pneumococcal protein antibodies in lung of HIV-infected adults.** *Vaccine.* 2013; **31**(35): 3469–3472.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Eagan R, Twigg HL 3rd, French N, *et al.*: **Lung fluid immunoglobulin from HIV-infected subjects has impaired opsonic function against pneumococci.** *Clin Infect Dis.* 2007; **44**(12): 1632–1638.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Agostini C, Zambello R, Trentin L, *et al.*: **Gamma delta T cell receptor subsets in the lung of patients with HIV-1 infection.** *Cell Immunol.* 1994; **153**(1): 194–205.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. De Paoli P, Gennari D, Martelli P, *et al.*: **A subset of gamma delta lymphocytes is increased during HIV-1 infection.** *Clin Exp Immunol.* 1991; **83**(2): 187–191.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Poccia F, Boullier S, Lecoœur H, *et al.*: **Peripheral V gamma 9/V delta 2 T cell deletion and anergy to nonpeptidic mycobacterial antigens in asymptomatic HIV-1-infected persons.** *J Immunol.* 1996; **157**(1): 449–461.
[PubMed Abstract](#)
28. Wallace M, Scharko AM, Pauza CD, *et al.*: **Functional gamma delta T-lymphocyte defect associated with human immunodeficiency virus infections.** *Mol Med.* 1997; **3**(1): 60–71.
[PubMed Abstract](#) | [Free Full Text](#)
29. Guadalupe M, Reay E, Sankaran S, *et al.*: **Severe CD4⁺ T-cell depletion in gut lymphoid tissue during primary human immunodeficiency virus type 1 infection and substantial delay in restoration following highly active antiretroviral therapy.** *J Virol.* 2003; **77**(21): 11708–11717.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Brechley JM, Paiardini M, Knox KS, *et al.*: **Differential Th17 CD4 T-cell depletion in pathogenic and nonpathogenic lentiviral infections.** *Blood.* 2008; **112**(7): 2826–2835.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Brechley JM, Knox KS, Asher AI, *et al.*: **High frequencies of polyfunctional HIV-specific T cells are associated with preservation of mucosal CD4 T cells in bronchoalveolar lavage.** *Mucosal Immunol.* 2008; **1**(1): 49–58.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Mahlknecht U, Deng C, Lu MC, *et al.*: **Resistance to apoptosis in HIV-infected CD4⁺ T lymphocytes is mediated by macrophages: role for Nef and immune activation in viral persistence.** *J Immunol.* 2000; **165**(11): 6437–6446.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Brittan M, Barr L, Conway Morris A, *et al.*: **A novel subpopulation of monocyte-like cells in the human lung after lipopolysaccharide inhalation.** *Eur Respir J.* 2012; **40**(1): 206–214.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Brittan M, Barr LC, Anderson N, *et al.*: **Functional characterisation of human pulmonary monocyte-like cells in lipopolysaccharide-mediated acute lung inflammation.** *J Inflamm (Lond).* 2014; **11**: 9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Yu YR, Hotten DF, Malakhau Y, *et al.*: **Flow Cytometric Analysis of Myeloid Cells in Human Blood, Bronchoalveolar Lavage, and Lung Tissues.** *Am J Respir Cell Mol Biol.* 2016; **54**(1): 13–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Baharom F, Thomas S, Rankin G, *et al.*: **Dendritic Cells and Monocytes with Distinct Inflammatory Responses Reside in Lung Mucosa of Healthy Humans.** *J Immunol.* 2016; **196**(11): 4498–509.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Ellery PJ, Tippett E, Chiu YL, *et al.*: **The CD16⁺ monocyte subset is more permissive to infection and preferentially harbors HIV-1 *in vivo*.** *J Immunol.* 2007; **178**(10): 6581–6589.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Murphy J, Summer R, Wilson AA, *et al.*: **The prolonged life-span of alveolar macrophages.** *Am J Respir Cell Mol Biol.* 2008; **38**(4): 380–385.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Williams M, De Kleer I, Henri S, *et al.*: **Alveolar macrophages develop from fetal monocytes that differentiate into long-lived cells in the first week of life via GM-CSF.** *J Exp Med.* 2013; **210**(10): 1977–1992.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Cai Y, Sugimoto C, Araing M, *et al.*: **Preferential Destruction of Interstitial Macrophages over Alveolar Macrophages as a Cause of Pulmonary Disease in Simian Immunodeficiency Virus-Infected Rhesus Macaques.** *J Immunol.* 2015; **195**(10): 4884–4891.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Swingle S, Mann AM, Zhou J, *et al.*: **Apoptotic killing of HIV-1-infected macrophages is subverted by the viral envelope glycoprotein.** *PLoS Pathogens.* 2007; **3**(9): 1281–1290.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Laforge M, Campillo-Gimenez L, Monceaux V, *et al.*: **HIV/SIV infection primes monocytes and dendritic cells for apoptosis.** *PLoS Pathogens.* 2011; **7**(6): e1002087.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Ruppert J, Schutt C, Ostermeier D, *et al.*: **Down-regulation and release of CD14 on human monocytes by IL-4 depends on the presence of serum or GM-CSF.** *Adv Exp Med Biol.* 1993; **329**: 281–286.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Gomez Perdiguero E, Klapproth K, Schulz C, *et al.*: **Tissue-resident macrophages originate from yolk-sac-derived erythro-myeloid progenitors.** *Nature.* 2015; **518**(7540): 547–551.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Italiani P, Boraschi D: **From Monocytes to M1/M2 Macrophages: Phenotypical vs. Functional Differentiation.** *Front Immunol.* 2014; **5**: 514.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 27 March 2018

<https://doi.org/10.21956/wellcomeopenres.14695.r31691>

© 2018 Bowdish D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dawn M.E. Bowdish

Department of Pathology and Molecular Medicine, McMaster University, Hamilton, ON, Canada

The applicants have clearly demonstrated differences in the immune composition of the lungs in HIV+ and negative individuals. Overall, I am very supportive of this work but I have a few minor concerns.

Methodological concerns:

My only concern is in the immunophenotyping of myeloid cells in blood and BAL. It would be helpful to label the gating strategy in Fig 4 with what cell types each group represents (e.g. mDC, monocyte). Is the gating strategy for the lung and blood the same? The proportions of the three monocyte subsets in blood looks slightly different than reported elsewhere and I wonder if this is due to differences in gating? Specifically intermediate monocytes are HLADR+ and it looks like in Fig 4, this is not a component of the gating strategy? CD11c is also found on circulating monocytes (esp non-classical) – is it possible that these are included in the DC gates rather than the monocyte gates? I find the gating strategy in Patel et al. Journal of Experimental Medicine Jun 2017, jem.20170355; DOI: 10.1084/jem.20170355 is consistent for monocytes in blood and tissues. It also appears that the monocytic proportion of the BAL are gated first on CD206 – although CD206 is found on alveolar and interstitial alveolar macrophages, I'm unaware of it being expressed on lung monocytes and so this might explain the discrepancy.

Grammar: In the heading "Sample collection and experimental procedures, the speeds at which the samples are centrifuged at should be "x g" not "g" (otherwise you are measuring in grams).

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 29 Mar 2018

Kondwani Jambo, Malawi Liverpool Wellcome Trust Programme of Clinical Tropical Research, Malawi

Reviewer comment: My only concern is in the immunophenotyping of myeloid cells in blood and BAL. It would be helpful to label the gating strategy in Fig 4 with what cell types each group represents (e.g. mDC, monocyte).

Response: We have labelled the gating strategy in Figure 4

Reviewer comment: Is the gating strategy for the lung and blood the same?

Response: No. We utilised two different gating strategies for the blood and lung, as the composition of the cells in the two compartments is different.

Reviewer comment: The proportions of the three monocyte subsets in blood looks slightly different than reported elsewhere and I wonder if this is due to differences in gating?

Response: The population of monocytes in blood is identified using standard gating (see Supplementary Figure 2). We have data from a similar cohort (unpublished), in which we made a direct comparison of monocyte subsets in an African cohort compared to European cohort, which shows that the proportions of the monocyte subsets in blood are different in the two groups. Specifically, the proportion of classical monocytes is higher in European individuals compared to African individuals, and the proportion of non-classical monocytes is higher in the African individuals compared to European individuals.

Reviewer comment: Specifically intermediate monocytes are HLADR+ and it looks like in Fig 4, this is not a component of the gating strategy?

Response: We can confirm that the monocyte population was HLADR positive. See plots in Supplementary Figure 3

Reviewer comment: CD11c is also found on circulating monocytes (esp non-classical) – is it

possible that these are included in the DC gates rather than the monocyte gates?

Response: We have noted an error in the presentation of the gating strategy. The DC gate actually is derived from the monocyte plot (CD14^{lo/-} CD16⁻ population), and not from the CD206⁻ CD66b⁻ population. The error on gating strategy has now been corrected (See Figure 4).

Furthermore, we acknowledge that the Myeloid DCs population might not be pure. However, upon further examination of the cells denoted as dendritic cells, we found that the cells are smaller than alveolar macrophages, less granular than neutrophils and exhibit low expression for CD206 (in comparison to alveolar macrophages) (see Supplementary Figure 4).

Reviewer comment: I find the gating strategy in Patel et al. Journal of Experimental Medicine Jun 2017, jem.20170355; DOI: 10.1084/jem.20170355 is consistent for monocytes in blood and tissues. It also appears that the monocytic proportion of the BAL are gated first on CD206 – although CD206 is found on alveolar and interstitial alveolar macrophages, I'm unaware of it being expressed on lung monocytes and so this might explain the discrepancy.

Response: Thank you for the manuscript. We agree peripheral blood monocytes do not express CD206. However, we know that when monocytes are differentiating into macrophages they gain expression of CD206. We suspect the expression of CD206 on the monocytes in BAL might be related to their differentiation process into monocyte-derived macrophages. According to Supplementary Figure 3, both subsets express CD206 at lower levels than alveolar macrophages, but CD14⁺CD16⁺ monocytes express higher levels of CD206 compared to CD14⁺CD16⁻.

Reviewer comment: Grammar: In the heading "Sample collection and experimental procedures, the speeds at which the samples are centrifuged at should be "x g" not "g" (otherwise you are measuring in grams).

Response: We agree with the reviewer. We have corrected the typo

Competing Interests: None

Reviewer Report 22 December 2017

<https://doi.org/10.21956/wellcomeopenres.14695.r29252>

© 2017 Dockrell D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

David H. Dockrell

MRC Centre for Inflammation Research, University of Edinburgh, Edinburgh, UK

The revised manuscript has addressed the reviewers comments for the most part. It is still not clear from the rather brief responses what if anything has been done to resolve the queries about HLA DR staining and use of CR2 staining etc. The answer may well be nothing further was possible to address this but some clarification would have been helpful unless I have missed this response. Otherwise it seems the authors have carefully addressed the comments.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 27 November 2017

<https://doi.org/10.21956/wellcomeopenres.13949.r27371>

© 2017 Meintjes G et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Graeme Meintjes

Groote Schuur Hospital, University of Cape Town, Cape Town, South Africa

Muki Shey

Department of Medicine, University of Cape Town, Cape Town, South Africa

Review by Graeme Meintjes

Mwale and colleagues report a cross-sectional study conducted in Blantyre, Malawi, in which 17 HIV-1 infected, ART naïve, asymptomatic adults and 20 HIV-1 uninfected adults were compared with respect to proportions and total number of leukocyte subsets in BAL obtained at bronchoscopy and in peripheral blood. The study is descriptive, providing a detailed profile of the leukocyte alterations associated with HIV-1 infection in these compartments.

My comments are:

1. More details regarding whether there was appropriate matching of cases and controls should be provided. It is unclear whether co-morbidities and environmental exposures that may impact lung immune responses were well matched across arms. For example, were respiratory diseases or symptoms present in any participants? Was a history of environmental exposures taken (eg. indoor smoke, occupational)? Were participants asked about previous TB, current TB symptoms and was any TB diagnostic work-up undertaken in those with symptoms? Was there a chest X-ray done in participants? Was any testing for TB infection ("latent TB") undertaken?
2. The term alveolitis is used in the title. "Alveolitis" suggests a clinical syndrome characterized by lung inflammation focused on the alveoli (as is the case with CD8+ alveolitis in HIV-infected patients – this manifests with a clinical syndrome). The authors have described increased numbers of certain immune cell types in the alveoli. I do not think this represents alveolitis and suggest changing this term in the title.
3. In the abstract it is stated that HIV "does not lead to massive depletion of alveolar CD4+ T cells". This is true for the participants studied but most had CD4 count in blood > 200 thus this statement should be qualified by recognising this may not apply to patients with severe CD4 depletion in blood.
4. In the Methods, the delay between taking the BAL sample and processing should be described. This may affect cell viability, especially neutrophils.
5. On page 3 for this statement: "In PBMCs, the absolute numbers were obtained by calculating the proportion of a particular subset relative to the full blood count (FBC) data." Suggest providing details about which cell populations from the FBC data were used for denominators in calculating cell numbers from the FACS data.
6. Page 7: "In blood, we found that the numbers of CD14+ CD16lo classical mono- cytes (median 110 vs. 60 cells/1000 mm³, p=0.0237), CD14+CD16+ intermediate monocytes (median 20 vs. 6 cells/1000 mm³, p=0.0362) and CD14loCD16+ non classical monocytes (median 10 vs. 30 cells/1000 mm³, p=0.0316) were higher in HIV-infected adults compared to HIV-uninfected controls." ... It appears the non classical monocytes were lower (10 vs 30) in HIV-infected adults not higher. Please check this.
7. Page 10: "We have shown preserved mucosal CD4+ T cells in BAL fluid from chronic HIV-infected adults, even in those with depleted peripheral blood CD4+ T cells." The statement I underlined should be justified by data presented. Was a stratified analysis of those with CD4<200 or <350 conducted?

8. The authors acknowledge that the accuracy of measurement of cell numbers in BAL may be inaccurate because of variations in the dilution factor of epithelial lining fluid and differences in BAL fluid volume return. They do not provide details on any methods they used to standardise this measurement across participants. For example in the BAL sampling technique or in standardising to a constituent of ELF. Suggest discuss this in more detail.

Additional comments from my colleague Dr Muki Shey:

General comments

I was not certain whether clinically the presence of immune cells in the lungs of (HIV-infected) individuals constitute alveolitis, otherwise definition and contextualization was missing.

There is no mention of TB infection in the participants or location where participants were recruited. Was there any screening for TB?

Some more discussion is necessary about the differences between cell numbers and proportions in BAL and Blood.

Specific comments

- Figure 1 needs revision: Different antibody fluorochromes are given in method section while different ones appear in the figure. For example, CD3 PE-Cy5 vs PerCP Cy5.5; and CD45 PE-Texas Red s CF594 (even though the two fluorochromes may be detected in the same channel, there needs to be consistency)
- Figure1: The CD4/CD8 axes labelling seems to be reversed. I am quite certain that the frequencies of CD4⁺ T cells cannot be more than frequencies of CD8⁺ T cells in HIV infected individuals
- The TCR Va chain is mentioned in the methods section while the TCR gd appears in the plots.
- The gating of CD45⁺ cells looks more stringent in Figure 1 (for T cells) than in Figure 4 (for myeloid cells). Was there any reason for that?
- The definition of different cell subsets by expression of markers may be necessary, for example the definition of NK T cells (CD3⁺CD56⁺?)
- In the results, giving the various median values in the text which are in the figures seems a duplication.
- HLA-DR antibody is mentioned in methods section but not shown either as a marker of activation or identification of myeloid cells. HLA-DR is an important marker of myeloid cell identification, especially in whole blood.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

No

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: HIV-associated tuberculosis

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however we have significant reservations, as outlined above.

Author Response 15 Dec 2017

Kondwani Jambo, Malawi Liverpool Wellcome Trust Programme of Clinical Tropical Research, Malawi

Thank you for the helpful comments. We have revised the manuscript to incorporate your suggestions.

Specifically, we have revised the title of manuscript, provided more details in the methodology section and corrected errors in the results section.

Competing Interests: None

Reviewer Report 30 October 2017

<https://doi.org/10.21956/wellcomeopenres.13949.r27370>

© 2017 Dockrell D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

David H. Dockrell

MRC Centre for Inflammation Research, University of Edinburgh, Edinburgh, UK

Mwale and co-workers examine the cellular composition of BAL asymptomatic HIV-positive Malawians. They show increased numbers of CD8 T-cells, B-cells and gd T-cells, while also identifying a reduction in classical monocytes in HIV which resulted in an increased proportion but not absolute number of intermediate monocytes. The data is important and adds a significant contribution to the literature, in particular through providing clinically relevant samples in HIV in patients naïve to ART. While the data makes a valuable contribution to the literature some areas of further detail would be informative.

1. Background demographic data shows the control group are well matched in terms of age and gender. More details on the HIV-positive group are required. What steps were taken to exclude TB. Is there any data on baseline CXR or TB screening or is the assumption that patients were negative purely based on lack of symptoms? Is there any data on baseline HIV viral load?
2. Some further methodological details would aid interpretation. While the differences in proportions of cell subsets are clear interpretation of absolute numbers requires evidence that volumes of instillation and BAL recovery are similar. Can the authors report their standard methodology involved instillation of 200 mls in four aliquots, into the right middle lobe, or whatever? Have they specific data on the volume recovered and its variability. Can they confirm there were low (<5%) numbers of bronchial epithelial cells or squamous cells?

Although the focus is analysis of cellular components have they any information on the permeability of the alveolar space in the two groups through measurement of albumin or a related marker? The authors appropriately remark in the discussion that these considerations limit interpretation of absolute numbers so this comment is meant only to provide detail not as a significant criticism.

3. The absolute number of monocytes in BAL should also be reported. In the abstract, the primary finding is presented as an increase in the proportion of intermediate monocytes in HIV BAL but in reality, the main finding appears to be a reduction in absolute number of classical monocytes which results in a relative rather than absolute increase in intermediate monocytes in HIV. Subsets of intermediate monocytes may also be defined by HLA-DR and the authors appear to have also used antibodies against HLA-DR. Did they find any differences in intermediate subsets by HLA-DR in their HIV positive population?
4. While reductions in non-classical monocytes are described by many groups the magnitude of the reduction in numbers of non-classical monocytes in the BAL is a little surprising since these cells are thought to be a source of alveolar macrophages. The authors suggest they can detect very few classical cells in BAL in contrast to the blood. Some more discussion of this point seems needed. Have the authors any data with alternative markers e.g. CCR2, CX3CR1 to confirm such low numbers? Or may they be missing some non-classical monocytes? This finding should be developed further and discussed a little further and related to other BAL-specific lung data.

5. In the discussion the authors highlight the different origins of monocytes and lung

macrophages but some qualification of the differences in origin of lung macrophages in inflammatory settings and the potential for classical monocytes to contribute to lung macrophage numbers in inflammation may be pertinent.

6. In Figure 5D the HIV + dot has been labelled HIV – and needs to be altered.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 15 Dec 2017

Kondwani Jambo, Malawi Liverpool Wellcome Trust Programme of Clinical Tropical Research, Malawi

Thank you for the helpful comments. We have revised the manuscript to incorporate your suggestions.

Specifically, provided more details in the methodology section and added more information in the discussion section.

Competing Interests: None