RESEARCH ARTICLE

REVISED Eco-geographic patterns of child malnutrition in India and its association with cereal cultivation: An analysis using demographic health survey and agriculture datasets [version 3; peer review: 2 approved with reservations, 1 not approved]

Rama Krishna Sanjeev¹, Prashanth Nuggehalli Srinivas², Bindu Krishnan³, Yogish Channa Basappa², Akshay S. Dinesh⁴, Sabu K. Ulahannan²

¹Pediatrics, Rural Medical College, Pravara Institute of Medical Sciences, Loni (BK), Ahmednagar district, Maharashtra, 413736, India

²Health equity cluster, Institute of Public Health Bengaluru, Bengaluru, Karnataka, 560070, India

³Physiology, Rural Medical College, Pravara Institute of Medical Sciences, Loni (BK), Ahmednagar district, Maharashtra, 413736, India

⁴Metastring Foundation, Bengaluru, Karnataka, 560020, India

v3 First published: 02 Jun 2020, 5:118
<https://doi.org/10.12688/wellcomeopenres.15934.1>
 Second version: 09 Nov 2020, 5:118
<https://doi.org/10.12688/wellcomeopenres.15934.2>
 Third version: 30 Sep 2021, 5:118
<https://doi.org/10.12688/wellcomeopenres.15934.3>
 Latest published: 22 Feb 2022, 5:118
<https://doi.org/10.12688/wellcomeopenres.15934.4>

Abstract

Background: High prevalence of maternal malnutrition, low birth-weight and child malnutrition in India contribute substantially to the global malnutrition burden. Rural India has disproportionately higher levels of child malnutrition. Stunting and wasting are the primary determinants of child malnutrition and their district-level distribution shows clustering in different geographies and regions. Cereals, particularly millets, constitute the bulk of protein intake among the poor, especially in rural areas in India where high prevalence of wasting persists.

Methods: The last round of National Family Health Survey (NFHS4) has disaggregated data by district, enabling a more fine-scale characterisation of the prevalence of markers of malnutrition. We used data from NFHS4 and agricultural statistics datasets to analyse relationship of prevalence of malnutrition at the district level and area under cereal cultivation. We analysed malnutrition through data on under-5 stunting and wasting by district.

Results: Stunting and wasting patterns across districts show a distinct geographical and age distribution; districts with higher wasting showed relatively higher prevalence before six months of age. Wasting prevalence at district level was associated with higher

Open Peer Review

Approval Status ✓ ✗ ? ?

	1	2	3	4
version 4				
(revision)	✓			?
22 Feb 2022	view			view
version 3				
(revision)	?		?	
30 Sep 2021	view		view	
version 2				
(revision)	?	✗		
09 Nov 2020	view	view		
version 1				
02 Jun 2020	✗	✗		
	view	view		

1. Edward J.M. Joy¹, London School of Hygiene & Tropical Medicine, London, UK
2. Suparna Ghosh-Jerath, Public Health Foundation of India, Gurgaon, India
3. Nicholas J. Kassebaum¹, University of

cultivation of millets, with a stronger association seen for jowar and other millets (Kodo millet, little millet, proso millet, barnyard millet and foxtail millet). District level stunting was associated with higher district level cultivation of all crops (except other millets). The analysis was limited by lack of fine-scale data on prevalence of low birth-weight and type of cereal consumed.

Conclusions: Better cereal cultivation and consumption data will be needed to confirm causal pathways contributing to potential ecogeographic patterns. The cultivation of other millets has a strong association with prevalence of wasting. State-of-the-art studies that improve our understanding of bio-availability of amino acids and other nutrients from the prevalent dietary matrices of rural poor communities will be needed to confirm causal pathways contributing to potential eco-geographic patterns.

Keywords

Millets, malnutrition, wasting, stunting, MTORC1 (mechanistic target of Rapamycin complex1), GCN2 (general control non derepressible 2), DSCQ (District subsistence cultivation quantum)

Washington, Seattle, USA

4. **Ilana Cliffer** , Harvard T.H. Chan School of Public Health, Boston, USA

Any reports and responses or comments on the article can be found at the end of the article.

This article is included in the [Wellcome Trust/DBT India Alliance](#) gateway.

Corresponding author: Prashanth Nuggehalli Srinivas (prashanthns@iphindia.org)

Author roles: **Sanjeev RK:** Conceptualization, Data Curation, Formal Analysis, Methodology, Project Administration, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Nuggehalli Srinivas P:** Conceptualization, Formal Analysis, Methodology, Project Administration, Software, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Krishnan B:** Conceptualization, Data Curation, Formal Analysis, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Basappa YC:** Data Curation, Formal Analysis, Methodology, Software, Validation, Visualization, Writing – Review & Editing; **Dinesh AS:** Formal Analysis, Software, Visualization, Writing – Review & Editing; **Ulahannan SK:** Conceptualization, Data Curation, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Wellcome Trust through a DBT/Wellcome Trust India Alliance Fellowship to PNS [IA/CPHI/16/1/502648]. This award also supports YCB and SKU.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Sanjeev RK *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Sanjeev RK, Nuggehalli Srinivas P, Krishnan B *et al.* **Eco-geographic patterns of child malnutrition in India and its association with cereal cultivation: An analysis using demographic health survey and agriculture datasets [version 3; peer review: 2 approved with reservations, 1 not approved]** Wellcome Open Research 2021, 5:118 <https://doi.org/10.12688/wellcomeopenres.15934.3>

First published: 02 Jun 2020, 5:118 <https://doi.org/10.12688/wellcomeopenres.15934.1>

REVISED Amendments from Version 2

We have revised V2 of the manuscript extensively in response to comments of reviewers 1 & 2 as well as inputs we sought from two other researchers working on topics related to the manuscript (now added in the revised acknowledgements). The major changes are:

The title has been changed to further narrow scope in line with the revised results.

The abstract has been rewritten to bring it in line with the revised results. The exploratory hypothesis-building focus on the manuscript has been clarified. Assertions related to causal claims in linking staple millet use to particular forms of childhood malnutrition have been examined and revised.

The introduction has been reworked for greater emphasis on the impact of cereals on malnutrition.

The low maternal BMI & maternal short stature component has been removed as it was not adding value to the current analysis.

Correction of minor errors in results. The maps have undergone minor changes due to a 1 % change in stunting and wasting figures which are in synchrony with cited literature. Table 2 having districts with higher prevalence of stunting and wasting has other additions which were overlooked in the earlier versions. The number of districts with higher prevalence of stunting has been updated in the results.

In the discussion we clarified the effects of cooking on the availability of nutrients in greater detail with recent literature. The rationale for use of amino acid availability in cereals has also been strengthened.

New figure with district-level proportions of cereal cultivation to visually depict relative proportions of cultivation of different cereal types in districts with high wasting and stunting has been added.

Limitations have been expanded considerably in response to reviewer comments.

Any further responses from the reviewers can be found at the end of the article

Introduction

Undernutrition in childhood includes intrauterine growth restriction (defined as below the 10th centile of intergrowth standards for the gestational age and sex), apart from stunting (z score for length/height for age less than 5 years below -2), wasting (z score for weight for length/height for age less than 5 years below -2) and underweight (z score for weight for age below -2)¹. Low pre-pregnancy BMI, low maternal BMI (< 18.5 kg/m²), maternal short stature and maternal micronutrient deficiency or anemia all contribute to small for gestational age, low birth weight and prematurity². Out of the estimated 20.5 million babies born low birth weight, 48% are born in South Asia. India alone is estimated to have 100 million adult women with low BMI^{1,2}. Globally, the World Health Organization (WHO) estimates that among children under five, about 151 million suffer from stunting and 51 million from wasting with consequent risks of mortality, morbidity and delayed development³. The latest stunting trends indicate increases in Africa along with substantial reductions across Asia. However, with regards to wasting,

with a regional prevalence of 12%, South Asia accounts for half of all wasted children globally^{1,4,5}. India reports 15.4% wasting of children under 5 years numbering about 27 million⁴. South Asia is also estimated to have ~45% of the global burden of stunting. The socio-economic gains and poverty reduction of the past decades have not translated into commensurate reduction of stunting and wasting in children, often characterised as, *the Asian enigma*^{6–8}.

Subsistence farming and millet dependence

Indian states consist of 640 districts (at the time of NFHS4) with wide differences in geography, climate and the main agricultural crops. India has a large and poor rural population (68.9% rural with 25.5 % rural poverty prevalence), and over half (54%) of the working rural population (481.9 million) are cultivators and agricultural labourers^{9,10}. Small land-holding farmers (owning less than two hectares of land) and their families constitute more than half the country's population. Only half (96.46 million hectares) of the total area under cultivation (198.36 million hectares) is irrigated¹¹. Although, rice and wheat together constitute 75% of total area under food grain cultivation, Jowar (Sorghum) and Bajra (pearl millet) make up a significant 13.8%. However, the distribution of food grain cultivation in irrigated land varies, with rice (60%) and wheat (94.2%), expectedly being grown largely on irrigated land. In contrast, Sorghum (Jowar) and Pearl millet (Bajra) are grown largely in non-irrigated lands, most likely by small land-holding farmers in monsoon-dependent arid or semi-arid regions of the country, which are also among the poorest^{12,13}. Cereal cultivation and consequently household food grain consumption and diets in such regions are likely driven by these strong linkages between agro-climatic, edaphic, and eco-geographic factors, more so among poorer households with socio-economic barriers to achieve dietary diversity.

A study based on National Family Health Survey-3, which reported results at the state level for India in 2005–6, demonstrated considerable geographic variation among the states of India with regards to child malnutrition, with higher levels of stunting seen in Uttar Pradesh, Uttaranchal & Gujarat¹⁴. In contrast, higher wasting levels were seen in Madhya Pradesh, a state in central India. A nutritional survey among preschool children in three tribal regions belonging to different ecological zones in the state of Madhya Pradesh, India, namely Jhabua, Bastar and Sarguja, showed greater extent and severity of malnutrition among children in Jhabua. The staple cereals reported in the study for Jhabua was maize and Sorghum, while for Bastar and Sarguja, it was rice¹⁵. Sorghum, as staple, has also been linked to endemic pellagra among farm workers in Hyderabad by Gopalan¹⁶.

The Lancet 2008 series too has brought out this aforementioned pattern of child malnutrition, with areas having similar prevalence of stunting demonstrating substantial differences in wasting¹⁷. Likewise, low women's BMI (15–49 years age) too, has numerous geographical subnational hotspots in South Asia¹.

Geo-spatial heterogeneity in prevalence of child malnutrition across Indian districts has been reported¹⁸. The National Family Health Survey 4 (NFHS 4) was published in 2015 with district-level data for the first time¹⁹. Based on unpublished field observations of wasting prevalence among populations depending on millet as staple in rural Maharashtra (spanning western and central India), we critically examined the spatial patterns of prevalence of stunting and wasting at the district level across India with the objective of exploring the role of dietary staple cereal consumption pattern using cultivation pattern as a Proxy. Ragi (finger millet; *Eleusine coracana*) was excluded because it belongs to a distinct sub-family in the grass family Poaceae and has a relatively better nutritional profile^{20–22}. We also propose a hypothetical pathway that integrates evidence emerging from agro-climatic and geographic patterns with physiological mechanisms of malnutrition.

Methods

We analysed district-level secondary data on under-5 stunting and wasting as reported in NFHS4 with district-wise crop cultivation data to assess geo-spatial overlaps and risk relationships between pre-school child malnutrition and cultivation of staple cereal crops. NFHS is a standardised and periodic nationally representative survey. NFHS4 covered 601,509 households, 699,686 women aged 15–49 years and 103,525 men aged 15–54 years that provides comprehensive data on various aspects of maternal and child health^{18,23}. NFHS-4 provides unit level data (for each of the 640 districts of India at the time of survey) for download upon request via the demographic health survey data repository^{23,24}. We extracted data on population of each district from the 2011 Census¹⁰. We included other socio-demographic variables with known associations with malnutrition from NFHS4 to assess their relative contribution to childhood wasting and stunting, at district level using multivariable linear regression.

Definitions and data sources

We considered the following cereal crops widely grown and reported in Indian agriculture databases and the Directorate of Millets Development (under Department of Agriculture, Co-operation and Farmers Welfare) in our analysis: rice (*Oryza sativa*), wheat (*Triticum aestivum*), maize (*Zea mays*), jowar (sorghum; *Sorghum bicolor*), bajra (pearl millet; *Pennisetum glaucum*) and other millets (kodo millet: *Paspalum scrobiculatum*, little millet: *Panicum sumatrense*, proso millet: *P. miliaceum*, barnyard millet: *Echinochloa esculenta* and foxtail millet: *Setaria italica*). Henceforth, millets in the text means Jowar, Bajra and other millets combined.

We adopted the definitions of districts with high prevalence of wasting and stunting from district-level malnutrition analysis by Junaid and Mohanty¹⁸, which has considered >46% district-level stunting prevalence (Z score ≤ 2), and >28% district-level wasting prevalence (Z score ≤ 2) as representing high prevalence districts for stunting and wasting respectively.

We extracted variables of interest from NFHS4 (see variables listed below). For data on cultivation of cereal crops, we used DACNET, a web-based land use statistics information

system maintained by the Agriculture Informatics Division of the National Informatics Centre of the Government of India²⁵.

The following data were extracted to prepare a district-level dataset for analysis²⁶:

1. From the 2011 census data, district-wise total and rural population
2. From the NFHS4 data,
 - a. using appropriate weights BMI less than 18.5 and short stature less than 145cm of women aged 15–49 years, utilization of Anganwadi, dietary diversity (age 6–23months), women with 10years or more of education, household wealth quintiles (lowest and second), open defecation and rural population.
 - b. district-level percentage of wasting and stunting was calculated from the children dataset
 - c. percentage of people in household wealth quintiles, open defecation for a given district was calculated from household dataset
3. Various crop data is available in state-wise reports compiled by the Ministry of Agriculture and Farmers Welfare. We extracted district-level area under cultivation of cereals: rice, wheat, maize, ragi, bajra, jowar, and millets (by type as defined above) into a spreadsheet. Data was from the latest state-wise reports available at the time of analysis at DACNET²⁵ (data for most states ranged for years between 2014–17 except Maharashtra 2002–03, Manipur 2004–05 and Gujarat 2007–08; all data in hectares converted to acres).

Using district names as the common variable in all three datasets, the variables from these three datasets were merged into a single dataset²⁶. Any errors due to district spellings and duplicate district names across differing states were handled with caution to ensure proper merging. For each district we estimated the population of poor by multiplying the census figures for population of the district by the proportion of the population in the fourth and fifth wealth quintiles (from NFHS4). This was based on the assumption that subsistence cereal consumption is largely restricted to poor small land-holding farmers^{27,28}. Since, Sorghum and other millets are largely cultivated by poor farmers with small land holdings for subsistence purposes with the exception of economically better-off and well-irrigated regions, particularly in northern India^{28–30} District Subsistence Cultivation Quantum (DSCQ) for each district was obtained by multiplying the per capita area (cereal cultivation area in acres/total population) by the proportion of the poor (in the lowest two wealth quintiles as per NFHS) followed by normalising data using logarithmic transformation.

Independent variables. We used the DSCQ for each cereals and other sociodemographic variables listed above while describing the district level dataset. Other socio-demographic

variables included in our analysis have been listed above while describing the district level dataset.

Outcome variables: percentage of children with wasting (Z score of weight for height <2), percentage of children with stunting (Z score of height for age <2).

Analysis

Spatial malnutrition patterns. We assessed overlaps between high prevalence of stunting and/or wasting with cereal cultivation data by generating maps derived from The [Database of Global Administrative Areas \(GADM\)](#)³¹. We merged tabular data (from a spreadsheet file) with geographic data (from a geojson file), chose variables of interest, created map legends dynamically and rendered multiple maps using a custom-built wrapper software written in javascript which internally uses [Mapbox GL JS library](#) (version 1.10.0) for rendering maps³². Further information on what this software wrapper does and how it works is present in the README file of the [source code](#)³³. As a base layer, DSCQ was shaded using a linear interpolator with manually chosen colour levels for the legend. A transparent layer of outcome variables (stunting and wasting) marked with distinct stripe patterns was overlaid on the base layer for visualizing overlap.

Examining relationship between subsistence millet cultivation, childhood malnutrition and its early onset. For each cereal, we examined its association with district-level prevalence of stunting and wasting and DSCQ (normalised using logarithmic transformation) by linear regression. We also examined the relationship of age with wasting and stunting at the district level.

Results

In all, 107 districts had a high prevalence of stunting (ranging from 46–65% district prevalence) with risk concentrated in poorer states: Uttar Pradesh (31; 29%) Bihar (28; 26%) and Madhya Pradesh (13; 12%) (numbers in brackets are number of districts followed by percentage). Among the 112 districts, those with higher rates of wasting (ranging from 28–47% district prevalence) were in districts with pre-dominantly tribal population in Jharkhand (14; 12.5%), Madhya Pradesh (20; 17.8%), Maharashtra (12; 11%), Rajasthan (11; 9.8%) and Gujarat (10; 9%) (numbers in brackets are number of districts followed by percentage). High stunting areas were concentrated in north and eastern India, whereas areas of high wasting were primarily in central India, which had high prevalence of both childhood stunting and wasting ([Figure 1](#)). There were 21 districts with high levels of both stunting and wasting, of which 16 had millets of any type or maize as either the most dominant (n=6) or second most dominant crop (n=10) ([Table 1](#)). Of these 21 districts, there were three from Rajasthan, which had more maize cultivation than any other cereal crop: Udaipur (62 %), Banswara (51%), and Dungarpur (47%).

On examining the district-level patterns of subsistence cultivation of jowar by district overlaid over districts having higher prevalence of stunting and wasting, we find that there is an overlap of districts with wasting alone and those with stunting and wasting with higher DSCQ for jowar ([Figure 2](#)). However, large areas with higher DSCQ particularly in North, West and some parts of central India do not show either stunting or wasting. Similar maps, separately showing overlap of high stunting and high wasting with per-capita cultivation of jowar, wheat, rice, bajra, maize, and other millets are also available³³. There is an overlap of districts with high wheat and rice cultivation in the well irrigated Gangetic plain (North and Eastern parts) with stunting ([Figure 5](#) & [Figure 6](#)). Cultivation of other millets is scattered throughout the country with an overlap with high prevalence of wasting. The large, irrigated areas in the Northwest & Central India with high DSCQ of Bajra & Jowar also have higher DSCQ of rice and wheat as seen in [Figure 2](#), [Figure 3](#), [Figure 5](#) & [Figure 6](#). Maize cultivation is all over the country with no clear overlap with either stunting or wasting.

Overall, increase in cultivation of jowar, bajra and other millets is independently associated with increase in prevalence of both stunting and wasting (see [Figure 3–Figure 5](#)). When the association was examined for individual millets, whereas jowar cultivation did show an association with increase in both stunting and wasting, increase in bajra cultivation was associated only with increase in stunting. Increase in cultivation of other millets was associated with increase in wasting only (a reverse trend was seen with stunting). As expected, there was either no change or decrease seen when we examined association between increase in rice or wheat cultivation with wasting (with an increase in stunting associated with increase in rice or wheat cultivation).

On examining the age of children in districts with higher prevalence of stunting and wasting the following observations are evident, as seen in [Figure 7](#) & [Figure 8](#). In 112 districts with high wasting, wasting showed an early onset with highest wasting (40%) less than 6 months of age ([Figure 7](#)). The age-distribution of stunting was similar for both groups of districts with highest age-specific stunting prevalence at 12 months and a plateau thereafter till five years of age ([Figure 7](#) & [Figure 8](#)). The earlier onset of wasting indicates the possibility of maternal nutritional factors affecting intra-uterine growth during pregnancy and possibly continuing in early infancy while being breast fed.

After accounting for the effects of known co-variables of child malnutrition, cultivation of jowar ($\beta = 0.32$, 95% CI: 0.163 - 0.488), other millets ($\beta = 3.372$, 95% CI: 1.404 - 5.341), women BMI less than 18.5 ($\beta = 0.211$, 95% CI: 0.135 - 0.287) are significant predictors of under-five wasting at district level ([Table 2](#)). On the other hand, cultivation of wheat ($\beta = 0.315$, 95% CI: 0.191 - 0.439), and women short stature

Figure 1. Map of India showing areas with higher prevalence of stunting ($>46\%$) in horizontal bars and those with higher prevalence of wasting ($>28\%$) in vertical bars. Districts with higher prevalence of both stunting and wasting are numbered cross referenced to [Table 1](#) and marked with oblique bars.

Table 1. Districts with high prevalence of stunting(>46%) and high prevalence of wasting(>28%) as per the integrated dataset from NFHS4 and agriculture statistics²⁶.

S.No	District	State	Major crop1	Percentage	Major crop 2	Percentage	Major crop 3	Percentage
1	Arwal	Bihar	Rice	73.64	Wheat	25.21	Maize	0.9
2	Sheikhpura	Bihar	Wheat	49.37	Rice	49.12	Maize	1.5
3	Narayanpur	Chhatisgarh	Rice	87.17	Other millets	7.66	Maize	3.8
4	Narmada	Gujarat	Rice	59.38	Jowar	27.07	Wheat	7.86
5	The Dangs	Gujarat	Rice	64.86	Other millets	16.98	Jowar	16.60
6	Chatra	Jharkhand	Rice	78.44	Maize	11.7	Wheat	8.99
7	Pashchimi Singhbhum	Jharkhand	Rice	98.89	wheat	0.61	Maize	0.47
8	Gumla	Jharkhand	Rice	88.18	Ragi	6.5	Other cereals	2.4
9	Gulbarga	Karnataka	Jowar	86.11	Bajra	5.10	Wheat	3.89
10	Yadgir	Karnataka	Rice	62.26	Jowar	24.99	Bajra	12.35
11	Alirajpur	Madhya Pradesh	Maize	42.4	Wheat	21.99	Bajra	14.42
12	Barwani	Madhya Pradesh	Maize	35.02	Wheat	33.77	Jowar	26.04
13	Bhind	Madhya Pradesh	Wheat	67.57	Bajra	22.00	Jowar	4.06
14	Morena	Madhya Pradesh	Wheat	50.46	Bajra	48.11	Rice	0.44
15	Shajapur	Madhya Pradesh	Wheat	99.44	Jowar	0.53	Rice	0.02
16	Sheopur	Madhya Pradesh	Wheat	66.24	Rice	21.57	Bajra	10.58
17	Banswara	Rajasthan	Maize	51.63	Wheat	34.42	Rice	12.17
18	Dungarpur	Rajasthan	Maize	47.50	Wheat	35.06	Rice	13.19
19	Udaipur	Rajasthan	Maize	62.65	Wheat	29.25	Jowar	2.62
20	Chitrakoot	Uttar Pradesh	Wheat	62.43	Bajra	13.36	Rice	11.23
21	Kaushambi	Uttar Pradesh	Wheat	53.69	Rice	35.95	Bajra	6.33

($\beta = 0.474$, 95% CI: 0.376 - 0.571) predict district level stunting (Table 3)

Discussion

Poverty and its antecedents affecting dietary intake, healthcare and poor environmental conditions can broadly explain malnutrition prevalence, particularly stunting worldwide². Nearly three decades ago, Victora identified the several-fold variation in wasting prevalence in areas with similar stunting prevalence, also highlighted more recently by Black *et al.* (2008)^{17,34}. In a particularly illuminating paper, Martorell *et al.* (2012) highlighted higher levels of early wasting with similar levels of stunting in India, in comparison to Guatemala³⁵. They have linked the higher prevalence of low BMI and maternal anaemia in India, while suggesting that improvements in maternal nutrition through prenatal interventions and during breastfeeding could address this problem of early wasting.

Our analysis attempts to show that ultimately, disaggregation of the data on type of malnutrition prevalence within India at a finer-scale along with proximate dietary factors in terms of staple cereal consumption holds the clues to such differences.

Global patterns of wasting linked to millet subsistence

The explanations for ecogeographic patterns of wasting in India appear to at least partially rest in subsistence cultivation patterns and staple consumption of millets as seen in our results. This pattern can be reproduced globally. Subsistence farming of millets overlaps with other global regions with some of the highest prevalence of wasting worldwide in Yemen and Sub-Saharan Africa^{3,30}. Similarly, in the low-lying areas of Jizan province of Saudi Arabia (which is adjoining Yemen), pearl millet is cultivated and widely used as a staple; expectedly, wasting prevalence in Jizan is the highest

Figure 2. Plots examining relationship between jowar cultivated with stunting and wasting at district level along with map showing the overlap of jowar cultivated with stunting and wasting. A) Scatterplot of stunting v/s district subsistence cultivation quantum (DSCQ) of jowar by poor **B)** Scatterplot of wasting v/s DSCQ of jowar by poor **C)** Geographic distribution of DSCQ of jowar by poor, stunting > 46 & wasting > 28.

among all provinces in Saudi Arabia^{36–38}. Malnutrition in early infancy (< six months of age) has been shown to be highly prevalent in India, Niger, Nigeria, Burkina Faso and Mali³⁹. These are in fact the top countries that produce millets for human consumption through subsistence farming⁴⁰. A paper by Grellety and Golden⁴¹ brought out differences in patterns of wasting in countries with some having more proportion of wasting assessed by MUAC, while others had greater proportion of wasting represented by low weight for height or having both low weight for height and low MUAC. On examining FAOSTAT data⁴⁰ to explain Grellety and Golden's assessment of differences in types of acute malnutrition across various countries, we found higher prevalence of wasting by MUAC to be a feature of maize-cultivating countries, while the countries reporting low weight for height are typically cultivating millets^{40–42}. These apparent differences in type of malnutrition linked to cereals has also been reproduced in observational studies. In a comparison of Bwamanda district (Democratic Republic of Congo with maize and Cassava staple) and Niakhar region (in Senegal with staple millet consumption), the former had higher proportion of wasting measured

by lower MUAC, while the latter had earlier onset of wasting and higher prevalence of low weight for heights⁴³. At a global level, the top 50 countries ranked among the hidden hunger scores⁴⁴ had either maize or millets/sorghum as staple among the top two cereals (as seen from FAOSTAT data of the top two cereals produced), an observation that may be linked to protein quality of these cereal crops^{40,42}.

Cereal protein quality

Sorghum, millets and maize share a common evolutionary ancestor in the grass family (Family *Poaceae*, sub-family *Panicoideae*), and can grow in arid/semi-arid agro-climatic regions where other crops often do not produce optimal yields through their dependence on the C4 carbon fixation pathway of photosynthesis^{45–48}. Millets are also usually not traded in markets but consumed directly by poor subsistence farmers^{28,29}. In rural India, the chief source of proteins are cereals⁴⁹. Further, there is a socio-economic gradient to protein quality; tribal populations and the poor consume lesser lysine-containing proteins primarily through cereals^{49,50}. The diet of rural pregnant and lactating women is particularly inadequate with

Figure 3. Plots examining relationship between bajra cultivated with stunting and wasting at district level along with map showing the overlap of bajra cultivated with stunting and wasting. A) Scatterplot of stunting v/s DSCQ of bajra by poor **B)** Scatterplot of wasting v/s DSCQ of bajra by poor **C)** Geographic distribution of DSCQ of bajra by poor, stunting > 46 & wasting > 28.

respect to quality of protein, thereby contributing to early malnutrition. Lysine content of millet (22 mg/g of protein) and sorghum (24 mg/g of protein) is the least among cereals in comparison to rice (35 mg/g of protein) and wheat (27 mg/g of protein)⁴⁹. The proportionate amino-acid requirement at infancy is the highest and shows an age-related decline^{49–53}. Extensive dependence on millets and sorghum as a staple among millions of poor rural communities where subsistence farming is the mainstay, in semi-arid and rainfed agricultural landscapes, motivated the FAO to commission a detailed assessment of their dietary protein quality³⁰. The report unequivocally highlights the inadequacy of millet and sorghum proteins for infants and young children based on amino acid scores³⁰. The other inexpensive and subsistence crop in India is maize with only 20% consumption, which (apart from its association with Kwashiorkor in the initial description by Cecily Williams) too has been extensively investigated for its causation of Pellagra^{54–57}. Similarly, Sorghum has been shown to be associated with Pellagra in Indian studies^{16,58}. Protein quality is assessed currently by the Digestible Indispensable amino acid scores (DIAAS) as per guidelines of the FAO⁵⁹. The DIAAS of different cereals is shown in Table 4,

drawn from a compendium curated by Hans-Henrik Stein of the University of Illinois, Urbana-Champaign (pers comm).

Sorghum protein is stored in Kafirins and is deficient in amino acids lysine with an excess of leucine^{16,21,30,58}. Millets in general have higher tannins^{21,30}; pearl millet has antinutrients like phytic acid, goitrogenic polyphenols, and tannins⁶⁰. Despite its better amino-acid profile (among the millets) the digestibility of proteins in Pearl millet (bajra) is probably less than other major grains²¹ due to antinutrients⁶⁰.

Effect of cooking and the dietary matrix: In India, the commonest mode of consumption of millets is by milling followed by removal of bran and making unleavened bread using (typically) dry heat⁶¹. Porridge-like preparations and cooked grains are also common. Sorghum protein becomes much less soluble after cooking⁶². These modes of processing are inferior to processes such as fermentation, germination or in combination which increase the availability of micronutrients, such as iron and zinc^{63,64}. Such cultural and social norms are important determinants of bioavailability of nutrients. For instance, maize is consumed in Latin America after nixtamalization,

Figure 4. Plots examining relationship between Other millets (kodo millet, little millet, proso millet, barnyard millet and foxtail millet) cultivated with stunting and wasting at district level along with map showing the overlap of Other millets cultivated with stunting and wasting. **A)** Scatterplot of stunting v/s DSCQ of other millets by poor **B)** Scatterplot of wasting v/s DSCQ of Other millets by poor **C)** Geographic distribution of DSCQ of Other millets by poor, stunting > 46 & wasting > 28 .

unlike in India where it is consumed directly, possibly explaining the stunting and wasting in the few districts in India where maize-growing for staple consumption is high⁶⁵. In separate studies for pearl millet and sorghum, conducted at the Hospital for sick children in Toronto (Canada), lysine availability was checked by assessing Indicator amino acid oxidation method (IAAO) after overnight soaking and boiling of the respective cereal in a rice cooker at 90°C. The pearl millet used had a lysine composition of 30.6 g/kg protein while sorghum had 27.31 g/kg protein. The studies yielded availability of cooked Lysine of 97 % for pearl millet and 94% for Sorghum⁶⁶⁻⁶⁸. However, the above method of cooking is significantly different from that used commonly in India for both sorghum and pearl millet. The flour is cooked by dry heat (at temperatures in the range 200°C-300°C) to make unleavened bread (variously called *bhakri* or *roti*)⁶⁹⁻⁷². Heat is known to adversely affect cereal protein availability by means of *Maillard reaction* and Lysinoalanine-like product formation, particularly in sorghum^{70,73}. So, the already lower lysine (as per composition per kg of protein) can get compromised further⁷³. In southern India, porridge like preparations are more common⁶⁹.

A study using stable isotope method with Indian rice-, wheat- and *ragi*-based meals for iron studies showed the least absorption with millet-based meals among the three. The study also indicates the importance of dietary matrix for iron absorption, even in cereals like *ragi*, which is replete in micro-nutrients like iron and calcium⁷⁴. Overall, there is an urgent need for state-of-the-art stable isotope studies in India to determine the metabolic availability of key nutrients from the common diets prevalent among rural poor.

Ready to use therapeutic food and aminoacids: The focus on protein quality has important biomedical and policy implications. Recent success with the use of peanut paste with milk based ready-to-use therapeutic food (RUTF) is being supplemented with well-intentioned attempts to use locally available ingredients in community-based malnutrition management approaches^{75,76}. These soya-maize-sorghum (SMS) formulations have been reported to be inferior in trials, particularly in children less than two years of age *vis-a-vis* peanut-based RUTF^{76,77}. However, when supplemented with free amino acids (free aminoacid soya-maize-sorghum RUTF), it has been shown to be as efficacious as standard peanut milk

Figure 5. Plots examining relationship between rice cultivated with stunting and wasting at district level along with map showing the overlap of rice cultivated with stunting and wasting. A) Scatterplot of stunting v/s DSCQ of rice by poor **B)** Scatterplot of wasting v/s DSCQ of rice by poor **C)** Geographic distribution of DSCQ of rice by poor, stunting > 46 & wasting > 28 .

based RUTF⁷⁸ thereby highlighting amino acids to be the missing link in the millet-based RUTF in the first 1000 days of life.

Micronutrient availability from cereals: However, in a poor family with low dietary diversity on a cereal based diet, in contrast to amino acid availability, the intake of micronutrients, particularly Zinc, could be lesser in the pre-dominantly rice and wheat cultivating areas (Table 5). Zinc levels, however, remains unaffected by maternal status and intake in the breast milk⁷⁹. However, calcium and possibly, Vitamin D are affected by maternal intake in the breast milk. Ragi is replete with both Calcium and Vitamin D in comparison to other cereals (Table 5). Figure 9 has pie charts showing cultivation of cereal crops all over India (A) as well as in districts with high wasting (9B) and high stunting (9C). The figures show that districts with high wasting have higher proportion of cultivation of coarse cereals (jowar, bajra, other millets and maize) with a commensurate reduction in proportion of rice and wheat. The converse is true for the proportion of crops in the 107 districts with high stunting. Hence, the higher prevalence of stunting and short stature in rice- and wheat-growing areas (Figure 2–Figure 8) could be due to micronutrient deficiencies, particularly of zinc^{80–82}.

Stunting, wasting and amino acid level

A metabolomic study involving hospitalized children with acute malnutrition (kwashiorkor & marasmus) in Malawi showed lower essential and conditionally essential amino acid levels in comparison to both stunted and non-stunted community controls⁸³. A similar community-based study showed lower amino acid profiles (particularly essential amino acids) among stunted children in comparison to non-stunted controls⁸⁴. These graded differences have raised questions about the possible causal role of amino acids in stunting⁸⁵. Similarly, it could be reasoned that further lower intake of amino acids could be linked to wasting in children with Kwashiorkor and marasmus as seen in the Malawi hospital study. Hence, in a comparable caloric intake among populations consuming different cereal based diets there could be different levels of amino acids and other key nutrients like glucose (because of varying glycemic indices of cereals⁸⁶) which are likely to have a bearing on growth as explained below.

Cellular pathways to malnutrition

Over the last three decades, pioneering research on cell growth has helped elucidate the critical role of complex intracellular nutrient-sensing mechanisms and their linkages with upstream and downstream pathways incorporating endocrine

Figure 6. Plots examining relationship between wheat cultivated with stunting and wasting at district level along with map showing the overlap of wheat cultivated with stunting and wasting. **A)** Scatterplot of stunting v/s DSCQ of wheat by poor **B)** Scatterplot of wasting v/s DSCQ of wheat by poor **C)** Geographic distribution of DSCQ of wheat by poor, stunting > 46 & wasting > 28 .

Figure 7. Age profile of stunted and wasted children in 108 high wasting (28-47%) districts.

Figure 8. Age profile of stunted and wasted children in 112 high stunting (46-67%) districts.

Table 2. Multivariable Regression models exploring the association between poor, women ≥ 10 years of education, Proportion of rural, Open defecation, Minimum dietary diversity, Utilization of anganwadi, Women's short stature (<145 cms) in 15-49 years of age, Women's BMI less than 18.5 in the 15-49 years of age, cultivation of Jowar, Bajra, other millets, rice and ragi and the outcome of interest is under 5 wasting.

Variables	Model-1	Model-2	Model-3	Model-4	Model-5	Model-6	Model-7
	Unadjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)
Poor	0.073*** (0.050 - 0.095)	-0.029 (-0.072 - 0.014)				0.005 (-0.036 - 0.047)	-0.025 (-0.072 - 0.022)
women ≥ 10 years of education	-0.074*** (-0.105 - -0.042)	-0.007 (-0.049 - 0.035)				0.016 (-0.026 - 0.058)	0.016 (-0.027 - 0.058)
Proportion of rural	0.030* (0.002 - 0.058)	-0.042* (-0.075 - -0.008)				-0.075*** (-0.107 - -0.042)	-0.058*** (-0.092 - -0.025)
Open defecation	0.125*** (0.106 - 0.145)	0.157*** (0.129 - 0.185)				0.120*** (0.092 - 0.148)	0.070*** (0.039 - 0.101)
Minimum dietary diversity	-0.168*** (-0.206 - -0.129)		-0.173*** (-0.210 - -0.136)			-0.119*** (-0.157 - -0.080)	-0.056** (-0.099 - -0.014)
Utilization of anganwadi	0.093*** (0.064 - 0.122)		0.098*** (0.070 - 0.125)			0.088*** (0.061 - 0.115)	0.046** (0.016 - 0.075)
Women short stature	0.026 (-0.073 - 0.125)						

Variables	Model-1	Model-2	Model-3	Model-4	Model-5	Model-6	Model-7
	Unadjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)
Women BMI less than 18.5	0.372*** (0.321 - 0.422)			0.372*** (0.321 - 0.422)			0.211*** (0.135 - 0.287)
Jowar	0.620*** (0.482 - 0.757)				0.670*** (0.500 - 0.840)		0.326*** (0.163 - 0.488)
Bajra	0.228** (0.085 - 0.371)				-0.306*** (-0.481 - -0.131)		-0.194* (-0.366 - -0.022)
Wheat	0.363*** (0.243 - 0.483)				0.365*** (0.243 - 0.488)		0.135 (-0.003 - 0.272)
Other millets	6.689*** (4.472 - 8.905)				4.299*** (2.185 - 6.412)		3.372*** (1.404 - 5.341)
Rice	0.052 (-0.128 - 0.233)						
Ragi	0.490*** (0.296 - 0.684)				0.488*** (0.303 - 0.674)		0.237* (0.057 - 0.418)
Observations	640	640	640	640	640	640	640
R-squared		0.218	0.167	0.247	0.221	0.299	0.382

*** p<0.001, ** p<0.01, * p<0.05

Table 3. Multivariable Regression models exploring the association between poor, women =>10 years of education, Proportion of rural, Open defecation, Minimum dietary diversity, Utilization of anganwadi, Women short stature (<145 cms) in 15-49 years age, Women's BMI less than 18.5, cultivation of Jowar, Bajra, other millets, rice and ragi and the outcome of interest is under 5 stunting.

Variables	Model-1	Model-2	Model-3	Model-4	Model-5	Model-6	Model-7
	Unadjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)
Poor	0.248*** (0.225 - 0.271)	0.052* (0.010 - 0.093)				0.062** (0.023 - 0.102)	0.031 (-0.014 - 0.076)
women =>10 years of education	-0.335*** (-0.367 - -0.304)	-0.197*** (-0.237 - -0.156)				-0.137*** (-0.177 - -0.097)	-0.104*** (-0.143 - -0.065)
Proportion of rural	0.164*** (0.130 - 0.198)	-0.044** (-0.077 - -0.012)				-0.030 (-0.061 - 0.001)	-0.012 (-0.043 - 0.019)
Open defecation	0.235*** (0.214 - 0.257)	0.146*** (0.119 - 0.173)				0.132*** (0.106 - 0.159)	0.086*** (0.057 - 0.114)

Variables	Model-1	Model-2	Model-3	Model-4	Model-5	Model-6	Model-7
	Unadjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)	Adjusted coefficients (95% CI)
Minimum dietary diversity	-0.339***		-0.336***			-0.149***	-0.082***
	(-0.384 - -0.294)		(-0.381 - -0.292)			(-0.186 - -0.113)	(-0.121 - -0.043)
Utilization of anganwadi	-0.058**		-0.048**			-0.068***	-0.038**
	(-0.096 - -0.019)		(-0.081 - -0.015)			(-0.093 - -0.042)	(-0.065 - -0.011)
Women short stature	0.902***			0.612***			0.474***
	(0.796 - 1.007)			(0.512 - 0.712)			(0.376 - 0.571)
Women BMI less than 18.5	0.571***			0.427***			0.065
	(0.511 - 0.631)			(0.368 - 0.487)			(-0.005 - 0.134)
Jowar	0.511***				0.265*		0.079
	(0.328 - 0.694)				(0.053 - 0.476)		(-0.072 - 0.230)
Bajra	0.467***				-0.052		0.146
	(0.285 - 0.648)				(-0.274 - 0.170)		(-0.011 - 0.304)
Wheat	0.964***				0.901***		0.315***
	(0.825 - 1.103)				(0.749 - 1.052)		(0.191 - 0.439)
Other millets	2.039						
	(-0.875 - 4.954)						
Rice	0.420***				0.361***		-0.102
	(0.191 - 0.650)				(0.153 - 0.570)		(-0.258 - 0.055)
Ragi	-0.222						
	(-0.475 - 0.030)						
Observations	640	640	640	640	640	640	640
R-squared		0.557	0.266	0.473	0.246	0.617	0.684

*** p<0.001, ** p<0.01, * p<0.05

inputs for growth, primarily centred on the role of protein kinases, MTORC 1 and MTORC2⁸⁷⁻⁹². Another protein kinase, GCN2 acts in concert with MTORC1 in sensing amino acid deficiencies^{90,91}.

From the results above, clearly, there are distinct geographic patterns of both stunting and wasting. There is also similarity in patterns of association between low maternal BMI with child wasting and maternal short stature with child stunting. Large-scale subsistence cultivation of millets in dry/semi-arid

areas of central India are more likely to be associated with higher wasting. Other factors like quantity of food, intake of proteins through milk and other animal sources during and before the first 1000 days by the mother, the dietary matrix and diversity and presence of infections are also important. However, the ecogeographic patterns of malnutrition can probably be explained to some extent by staple cereal cultivation. Hence, the proportionate shares of cultivation of cereals crops in the country as whole and in the district with high wasting and stunting is shown in [Figure 9](#)

Table 4. Digestible indispensable amino acid scores (DIAAS) determined for human foods using the pig or rat model (Data from published studies compiled by Hans-Henrik Stein, University of Illinois, Urbana-Champaign).

Cereal grains	Animal model/human	Reference Protein pattern			
		Infants(0–6 months)	Young children (6 months–3 years)	Older children, adolescents and adults	Reference
Rice, cooked	Rat	--	60 (lysine)	--	Rutherford <i>et al.</i> , 2015
Rice, polished, cooked	Rat	--	37 (lysine)	--	Han <i>et al.</i> , 2019
Rice protein	Pig	--	48 (lysine)	57 (lysine)	Exp. 620 [#]
Rice, white, polished, raw	Pig	--	--	64 (lysine)	Cervantes-Pahm <i>et al.</i> , 2014
Sorghum, raw	Pig	--	--	29 (lysine)	Cervantes-Pahm <i>et al.</i> , 2014
Millet, foxtail, cooked	Rat	--	10 (lysine)	--	Han <i>et al.</i> , 2019
Millet, proso, cooked	Rat	--	7 (lysine)	--	Han <i>et al.</i> , 2019
Corn, yellow dent, raw	Pig	--	--	48 (lysine)	Cervantes-Pahm <i>et al.</i> , 2014
Wheat, whole, cooked	Rat	--	20 (lysine)	--	Han <i>et al.</i> , 2019

[#] Exp 620 stands for Experiment 620 in the Laboratory of Prof Hans-Henrik Stein, University of Illinois, Urbana-Champaign

An attempt has been made here to build a hypothetical framework explaining the plausible pathways through which staple based cereal based diet could produce stunting and wasting (Figure 10)

A pre-dominantly millet diet has a low glycaemic index, putative deficiencies in amino acids and micronutrients with variable digestibility after cooking (Figure 10 pathway 2) due to the presence of a tough cell wall and antinutrients^{21,30,55,60,93}. This could lead to a greater lowering of amino acids and glucose, which in turn could lead to a substantial lowering of MTORC1 and rise in GCN2 contributing to wasting (Figure 10 pathway 5)^{87–91}. The lowering of MTORC1 & GCN2 at the cellular level causes low BMI in women (particularly growing adolescents) with consequent decreased acquisition of nutrients by placental syncytiotrophoblast^{87,88,94,95}. This in turn leads to intra-uterine growth restriction and low birth weight. These nutritional effects on the malnourished mother persist during breast feeding⁷⁹ (Figure 10 pathway 6) leading to early wasting and lower weight for height seen in regions with higher wasting (Figure 10 pathway 9)^{35,41,43,96,97}. Maize consumption without nixtamalization could lead to lesser unbound niacin (Figure 10 pathway 7)^{42,65,92,98}, and both stunting and wasting with lower MUAC (Figure 10 pathway 7)^{41–43}. Poverty in highly populous areas growing rice & wheat could be contributory to stunting. Possible greater stress, lower dietary quantity and poor sleep could be contributory (Figure 10 pathway 8)^{12,19,38,99,100}. Lower lysine, high phytates with lowered micronutrients like zinc and iron in cereal based diet in wheat and rice growing areas with poor dietary diversity could lead

to stunting (Figure 10 pathway 8)^{8,18,19,98–100}. Infections in the growing child can decrease stimulation of both MTORC1 and 2 through T cell mediated mechanisms (Figure 10 pathway 3)¹⁰¹. Millets are indeed gluten-free, have high fibre and antioxidant content and have recently seen a spike in use^{102–104}. However, their use by pregnant as well as nursing mothers and children in poor communities, with limited dietary diversity, during the first 1000 days could be associated with malnutrition.

Study limitations

An important limitation of our analysis is the limited fine scale data on low birth weight and food grain consumption (as opposed to cultivation) which would have allowed for confirmation of our hypothesis at household level. One of the reasons for this is that the NFHS and other country/regional demographic health surveys record cereal consumption without paying attention to type of cereal. Moreover, consumption is likely to be guided by choice and availability through food subsidy or open-market access to other cereals and food staples, apart from those cultivated for subsistence. Our analysis indicates the need for NFHS and demographic health surveys worldwide to include type of cereal consumption to gain better understanding of pathways to malnutrition. The use of cereal cultivation as a proxy for consumption too is a source of noise in our data, as some of the cultivation is likely to be for non-human use (primarily fodder for animal use). Factors leading to lack of dietary diversity like poverty, prevalence of infections like worm infestations or tuberculosis and other possible unaccounted confounding factors could also be contributing to these patterns. The data

Table 5. Micronutrient table for cereals compiled from Indian Food composition Tables 2017, National Institute of Nutrition, Hyderabad, India⁸².

S.no	Cereal Name	Fe (mg)	Zn (mg)	Ca (mg)	Se (mg)	In 100 gm of edible portion		Expressed per 100 gm edible portion		
						Total Carotenoids µg	Vit D2 µg	B1 (mg)	B3 (mg)	Total Folate B 9 mg
1	Bajra (Pennisetum typhoideum)	6.42 +/- 1.04	2.76 +/- 0.36	27.35 +/- 2.16	30.4 +/- 5.22	293 +/- 55.7	5.65 +/- 0.27	0.25 +/- 0.04	0.86 +/- 0.10	36.11 +/- 5.05
2	Jowar (Sorghum vulgare)	3.95 +/- 0.94	1.96 +/- 0.31	27.60 +/- 3.71	26.29 +/- 11.08	9.08 +/- 1.77	3.96 +/- 0.30	0.35 +/- 0.039	2.10 +/- 0.09	39.42 +/- 3.13
3	Maize, dry (Zea mays)	2.49 +/- 0.32	2.27 +/- 0.23	8.91 +/- 0.61	8.69 +/- 1.81	893 +/- 154	33.6 +/- 2.82	0.33 +/- 0.032	2.69 +/- 0.06	25.81 +/- 1.44
4	Ragi (Eleusine coracana)	4.62 +/- 0.36	2.53 +/- 0.51	364 +/- 58	15.30 +/- 6.23	154 +/- 25.6	41.46 +/- 3.12	0.37 +/- 0.041	1.34 +/- 0.02	34.66 +/- 4.97
5	Little millet or Samai (Panicum miliare)	1.26 +/- 0.44	1.82 +/- 0.14	16.06 +/- 1.54	40.41 +/- 24.09	120 +/- 9	0.28 +/- 0.80	0.26 +/- 0.042	1.29 +/- 0.02	36.20 +/- 7.04
6	Foxtail millet (Setaria italica)	2.34 +/- 0.46	1.65 +/- 0.18	15.27 +/- 1.28	14.12 +/- 2.26	272 +/- 25.1	-----	0.29 +/- 0.054	1.49 +/- 0.08	39.49 +/- 4.52
7	Rice, raw milled (Oryza sativa)	0.65 +/- 0.11	1.21 +/- 0.17	7.49 +/- 1.26	1.01 +/- 0.13	16.87 +/- 5.61	-----	0.05 +/- 0.019	1.69 +/- 0.13	9.32 +/- 1.93
8	Wheat flour,atta (Triticum aestivum)	1.77 # +/- 0.38	0.88 # +/- 0.07	20.4 # +/- 2.46	-----	284 +/- 31.9	13.43 +/- 1.77	0.42 +/- 0.044	2.37 +/- 0.10	29.22 +/- 1.92
9	Barley (Hordeum vulgare)	1.56 +/- 0.15	1.5 +/- 0.27	28.64 +/- 3.49	18.61 +/- 1.32	69.87 +/- 28.88	-----	0.36 +/- 0.059	0.86 +/- 0.10	31.58 +/- 3.79

For wheat flour refined

Figure 9. Pie charts showing Proportionate shares of cereal crops in districts with higher stunting (>46%), higher wasting (>28%) and all over India. (A) pie chart showing proportionate share of cereal crops in India as a whole (B) Pie chart showing proportionate share of cereal crops in 112 districts with higher prevalence of wasting (C) Pie chart showing proportionate share of cereal crops in 107 districts with higher prevalence of stunting.

Figure 10. A schematic figure showing the possible pathways linking staple cereal consumption with child malnutrition with key steps highlighted (with references to evidence from literature).

on availability of nutrients from cereal consumption from nutritional assays (stable isotope-based) is also meagre to the best of our efforts in reviewing peer-reviewed evidence-base. Such data from cereal consumption could help in linking the dietary matrix to the effects described above.

Conclusion

Higher wasting and stunting prevalence among children in India has an ecogeographic pattern with plausible links of pre-dominant millet consumption to higher prevalence of wasting. MUAC and type of cereal consumed should be

incorporated in NFHS4 and all global demographic surveys and health surveys to enable better assessment of patterns of malnutrition. State of the art research in nutrient sensing should be integrated with agriculture, food science, delivery systems and dietary matrix for translational benefits to accrue to the wider population.

Data availability

Underlying data

Figshare: Dataset used to assess relationship between millet cultivation and malnutrition patterns in India at district level. <https://doi.org/10.6084/m9.figshare.12236789.v2>²⁶

This project contains the following underlying data:

- malnutrition_dataset_for_publication.xlsx (Dataset used for analysis described in the paper)
- Malnutrition and millets – India – DACNET NFHS 4.docx (Word document explaining how the dataset was prepared)

Extended data

Figshare: Plots examining relationship between type of millet cultivated with stunting and wasting at district level along with map showing the overlaps for each type of millet with stunting and wasting. <https://doi.org/10.6084/m9.figshare.12206135.v4>¹⁰⁵

This project contains the following extended data:

- Malnutrition_millets and malnutrition.pdf (PDF file with panel of seven plots and maps, each showing relationship between type of millet cultivated with stunting and wasting at district level and the corresponding map showing the overlaps of each type of millet with stunting and wasting)

Figshare: Plots examining relationship between low BMI and short stature in women 15–49 with stunting and wasting at district level along with map showing the overlaps for

each type of millet with low BMI and short stature in women (15–49). <https://doi.org/10.6084/m9.figshare.12206264.v4>

This project contains the following extended data:

- malnutrition_bmi_short_status.pdf (PDF file with panel of seven plots and maps, each showing relationship between low BMI and short stature in women 15–49 with stunting and wasting at district level along with maps showing overlaps for each type of millet with low BMI and short stature in women (15–49))

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Software availability

Source code available from: <https://gitlab.com/asdofindia/malnutrition-crops-maps>

Archived source code at time of publication: <http://doi.org/10.5281/zenodo.3828725>³³

License: [MIT license](#)

Acknowledgements

RKS and BK would like to thank the Pravara Institute of Medical Sciences (PIMS) for encouraging research on child malnutrition, in particular, the Dr Rajendra Vikhe Patil (CEO & Pro-Chancellor), late Professor (Col) DY Shrikhande (Head of the department of Pediatrics), Professor Rahul Kunkulol (Director of Research) and Professor KV Somasundaram, Director of the Centre for Social Medicine at PIMS. All authors thank Awdhesh Yadav for help with organisation of the final dataset and statistical analysis, Shivanand Savatagi for help with figures, Megha for help with literature and Mahantesh Kamble for help with references. Authors would like to particularly thank Professor Keith West from Johns Hopkins Bloomberg School of Public Health for extensive inputs in the writing of the third version of the paper.

References

1. Victora CG, Christian P, Vaidaletti LP, *et al.*: **Revisiting maternal and child undernutrition in low-income and middle-income countries: variable progress towards an unfinished agenda.** *Lancet*. 2021; **397**(10282): 1388–1399. [PubMed Abstract](#) | [Publisher Full Text](#)
2. Christian P, Smith ER, Zaidi A: **Addressing inequities in the global burden of maternal undernutrition: the role of targeting.** *BMJ Glob Health*. 2020; **5**(3): e002186. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. UNICEF, WHO, International Bank of Reconstruction and Development, World Bank: **Levels and Trends in Child Malnutrition: Key Findings of the 2019 Edition of the Joint Child Malnutrition Estimates.** Geneva; 2019. [Reference Source](#)
4. Harding KL, Aguayo VM, Webb P: **Factors associated with wasting among children under five years old in South Asia: Implications for action.** *PLoS One*. 2018; **13**(7): e0198749. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Richard SA, Black RE, Gilman RH, *et al.*: **Wasting Is Associated with Stunting in Early Childhood.** *J Nutr*. 2012; **142**(7): 1291–1296. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Government of India: **Press Note on Poverty Estimates.** New delhi; 2013. [Reference Source](#)
7. World Bank: **Data bank poverty and equity data base.** World bank. 2020. [Reference Source](#)
8. Ramalingaswami V, Jonson U, Rohde J: **The Asian Enigma, the progress of nations.** 1997.
9. Motkuri V, Naik SV: **Growth and Structure of Workforce in India.** *Indian Econ J*. 2016; **64**(1–4): 57–74. [Publisher Full Text](#)

10. Office of the Registrar General & Census Commissioner: **Provisional Population Totals Paper 1 of 2011 India Series 1**. 2011; 188.
[Reference Source](#)
11. Singh R, Kumar PWT: **Smallholder Contributions to Agriculture: Smallholder Farmers in India: Food Security and Agricultural Policy**. Bangkok, Thailand; 2002.
[Reference Source](#)
12. Department of Agriculture and Farmers Welfare: **Pocket book of Agricultural statistics**. 2017.
[Reference Source](#)
13. Food and Agriculture Organization: **State of the world's forests**. 2005.
[Reference Source](#)
14. Yadav A, Ladusingh L, Gayawan E, *et al.*: **Does a geographical context explain regional variation in child malnutrition in India?** *J Public Health (Germany)*. 2015; **23**: 277–87.
[Publisher Full Text](#)
15. Rao DH, Rao KM, Radhaiah G, *et al.*: **Nutritional status of tribal preschool children in three ecological zones of Madhya Pradesh**. *Indian Pediatrics*. 1994; **31**(6): 635–640.
[PubMed Abstract](#)
16. Gopalan N, Srikantia SG: **Leucine and pellagra**. *Lancet*. 2018; **1**(7131): 954–957.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Black RE, Allen LH, Bhutta ZA, *et al.*: **Maternal and child undernutrition: global and regional exposures and health consequences**. *Lancet*. 2008; **371**(9608): 243–260.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Khan J, Mohanty SK: **Spatial heterogeneity and correlates of child malnutrition in districts of India**. *BMC Public Health*. 2018; **18**(1): 1027.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. International Institute for Population Sciences: **National Family Health Survey (NFHS-4) 2015-16 India**. 2017.
[Reference Source](#)
20. Shobana S, Krishnaswamy K, Sudha V, *et al.*: **Finger Millet (*Ragi*, *Eleusine Coracana* L.): A Review of Its Nutritional Properties, Processing, and Plausible Health Benefits**. *Adv Food Nutr Res*. 2013; **69**: 1–39.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. National Research Council: **Lost Crops of Africa**. Washington DC National Academies Press; 1996.
[Reference Source](#)
22. Cannarozzi G, Plaza-Wüthrich S, Esfeld K, *et al.*: **Genome and transcriptome sequencing identifies breeding targets in the orphan crop tef (*Eragrostis tef*)**. *BMC Genomics*. 2014; **15**(1): 581.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. NFHS: **National Family Health Survey-4 (NFHS-4)**. Ministry Heal Fam Serv India. 2016.
[Reference Source](#)
24. ICF International: **The DHS Program: demographic and health surveys**. DHS Model Questionnaires.
[Reference Source](#)
25. **Special Data Dissemination Standard Division, Directorate Of Economics & Statistics, Ministry Of Agriculture and Farmers Welfare, Govt. Of India. Web-based land use statistics information system of India: District-wise land use statistics**. Agriculture Informatics Division, National Informatics Centre, Ministry Of Communication & IT, Govt. Of India, New Delhi.
[Reference Source](#)
26. Basappa YC, Ulahannan SK, Dinesh AS, *et al.*: **Dataset used to assess relationship between millet cultivation and malnutrition patterns in India at district level**. *Figshare*.
<http://www.doi.org/10.6084/m9.figshare.12236789.v2>
27. Rao PP, Birlhal PS, Reddy BVS, *et al.*: **Diagnostics of sorghum and pearl millet grains-based nutrition in India**. *SAT eJournal*. 2006; **2**(1).
[Reference Source](#)
28. Rao PP, Basavaraj G: **Status and prospects of millet utilization in India and global scenario**. 2015; 197–209.
[Reference Source](#)
29. Basavaraj G, Rao PP, Bhagavatula S, *et al.*: **Availability and utilization of pearl millet in India**. *SAT eJournal*. 2010; **8**: 1–6.
[Reference Source](#)
30. Food and Agriculture Organization of the United Nations: **Sorghum and millets in human nutrition**. David Lubin Meml Libr Cat Publ Data FAO, Rome. 1995.
[Reference Source](#)
31. Global Administrative Areas: **GADM database of Global Administrative Areas, version 2.8**.
[Reference Source](#)
32. Mapbox: **Mapbox GL JS v1.10.0 Javascript library**.
[Reference Source](#)
33. Dinesh AS, Basappa YC, Ulahannan SK, *et al.*: **Cereal Cultivation and Wasting Patterns Maps: Fully contained release**. 2020.
<http://www.doi.org/10.5281/ZENODO.3828725>
34. Victora CG: **The Association between Wasting and Stunting: An International Perspective**. *J Nutr*. 1992; **122**(5): 1105–1110.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Martorell R, Young MF: **Patterns of Stunting and Wasting: Potential Explanatory Factors**. *Adv Nutr*. 2012; **3**(2): 227–233.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Al-Mendalawi MD: **Regional Disparity in Prevalence of Malnutrition in Saudi Children**. *Saudi Med J*. 2010; **31**(10): 1184.
[PubMed Abstract](#)
37. Khalid ME, Al-Hashem FH: **The relationship of body weight to altitude in preschool children of southwestern Saudi Arabia**. *J Family Community Med*. 2007; **14**(2): 71–76.
[PubMed Abstract](#) | [Free Full Text](#)
38. Gassem MA, Osman MA: **Chemical composition and nutritional potential of pearl millet grown in Jazan region of Saudi Arabia**. *J Saudi Soc Agric Sci*. 2008; **7**: 1–22.
39. Kerac M, Frison S, Connell N, *et al.*: **Informing the management of acute malnutrition in infants aged under 6 months (MAMI): risk factor analysis using nationally-representative demographic & health survey secondary data**. *PeerJ*. 2019; **6**: e5848.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Food and agricultural organization, Food, agricultural organization: **FAOSTAT main data base**. United nations. 2020; Accessed April 26, 2020.
[Reference Source](#)
41. Grellety E, Golden MH: **Weight-for-height and mid-upper-arm circumference should be used independently to diagnose acute malnutrition: policy implications**. *BMC Nutr*. 2016; **2**(1): 10.
[Publisher Full Text](#)
42. Ranum P, Peña-Rosas JP, Garcia-Casal MN: **Global maize production, utilization, and consumption**. *Ann N Y Acad Sci*. 2014; **1312**(1): 105–112.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Garenne M, Willie D, Maire B, *et al.*: **Incidence and duration of severe wasting in two African populations**. *Public Health Nutr*. 2009; **12**(11): 1974–1982.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Muthayya S, Rah JH, Sugimoto JD, *et al.*: **The Global Hidden Hunger Indices and Maps: An Advocacy Tool for Action**. Noor AM ed. *PLoS One*. 2013; **8**(6): e67860.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. Zhou H, Helliker BR, Huber M, *et al.*: **C₄ photosynthesis and climate through the lens of optimality**. *Proc Natl Acad Sci U S A*. 2018; **115**(47): 12057–12062.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Sage RF: **The evolution of C₄ photosynthesis**. *New Phytol*. 2004; **161**(2): 341–370.
[Publisher Full Text](#)
47. Osborne CP, Freckleton RP: **Ecological selection pressures for C₄ photosynthesis in the grasses**. *Proc Soc Sci*. 2009; **276**(1663): 1753–1760.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Badi SM, Hoseney RC, Casady AJ: **Characterization By Sem Amino Acid Analysis, Lipid Composition, And Prolamine Solubility**. 1976.
[Reference Source](#)
49. Swaminathan S, Vaz M, Kurpad AV: **Protein intakes in India**. *Br J Nutr*. 2012; **108**(Suppl 2): S50–S58. Swaminathan S, Hemalatha R, Pandey A, *et al.*: **The burden of child and maternal malnutrition and trends in its indicators in the states of India: the Global Burden of Disease Study 1990–2017**. *Lancet Child Adolesc Heal*. 2019; **3**(12): 855–870.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Minocha S, Thomas T, Kurpad AV: **Dietary Protein and the Health-Nutrition-Agriculture Connection in India**. *J Nutr*. 2017; **147**(7): 1243–1250.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. Uauy R, Kurpad A, Tano-Debrah K, *et al.*: **Role of Protein and Amino Acids in Infant and Young Child Nutrition: Protein and Amino Acid Needs and Relationship with Child Growth**. *J Nutr Sci Vitaminol (Tokyo)*. 2015; **61** Suppl: S192–S194.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Young VR, Pellett PL: **Plant proteins in relation to human protein and amino acid nutrition**. *Am J Clin Nutr*. 1994; **59**(5 Suppl): 1203S–1212S.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. Young VR, Pellett PL: **Wheat proteins in relation to protein requirements and availability of amino acids**. *Am J Clin Nutr*. 1985; **41**(5 Suppl): 1077–1090.
[PubMed Abstract](#) | [Publisher Full Text](#)
54. Khatkar BS, Chaudhary N, Dangi P: **Production and Consumption of Grains in India**. In: *Reference Module in Food Science*. Elsevier; 2016.
[Publisher Full Text](#)
55. Proietti I, Frazzoli C, Mantovani A: **Exploiting Nutritional Value of Staple Foods in the World's Semi-Arid Areas: Risks, Benefits, Challenges and Opportunities of Sorghum**. *Healthcare (Basel)*. 2015; **3**(2): 172–193.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Williams CD: **A nutritional disease of childhood associated with a maize diet**. *Arch Dis Child*. 1933; **8**(48): 423–433.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. Morabia A: **Joseph Goldberger's research on the prevention of pellagra**. *J R Soc Med*. 2008; **101**(11): 566–568.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

58. Gopalan C: **Pellagra in sorghum eaters.** 1972.
[Reference Source](#)
59. **Dietary protein quality evaluation in human nutrition. Report of an FAQ Expert Consultation.** *FAO Food Nutr Pap.* 2013; **92**: 1–66.
[PubMed Abstract](#)
60. Boncompagni E, Orozco-Arroyo G, Cominelli E, *et al.*: **Antinutritional factors in pearl millet grains: Phytate and goitrogens content variability and molecular characterization of genes involved in their pathways.** *Kashkush K, ed. PLoS One.* 2018; **13**(6): e0198394.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. Subramanian V, Jambunathan R: **Traditional Methods of Processing Sorghum Land Millets in India.** Patancheru, India; 1980.
[Reference Source](#)
62. Hamaker BR, Kirleis AW, Mertz ET, *et al.*: **Effect of cooking on the protein profiles and in vitro digestibility of sorghum and maize.** *J Agric Food Chem.* 1986; **34**(4): 647–649.
[Publisher Full Text](#)
63. Svanberg U, Lorri W, Sandbeag AS: **Lactic Fermentation of Non-Tannin and High-Tannin Cereals: Effects on In Vitro Estimation of Iron Availability and Phytate Hydrolysis.** *J Food Sci.* 1993; **58**(2): 408–412.
[Publisher Full Text](#)
64. Gibson RS, Ferguson EL: **Nutrition intervention strategies to combat zinc deficiency in developing countries.** *Nutr Res Rev.* 1998; **11**(1): 115–131.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. Nuss ET, Tanumihardjo SA: **Maize: A Paramount Staple Crop in the Context of Global Nutrition.** *Compr Rev Food Sci Food Saf.* 2010; **9**(4): 417–436.
[Publisher Full Text](#)
66. Alyssa P: **Metabolic Availability of Lysine in Cooked Sorghum in Healthy Adult Males Using the Combined Indicator Amino Acid Oxidation and Slope Ratio Method.** [dissertation]: University of Toronto; 2019.
67. Fakiha A, Tul-Noor Z, Paoletti A: **Bioavailable Lysine, Assessed in Healthy Young Men Using Indicator Amino Acid Oxidation, is Greater when Cooked Millet and Stewed Canadian Lentils are Combined.** *J Nutr.* 2020; **150**(10): 2729–2737.
[PubMed Abstract](#) | [Publisher Full Text](#)
68. Elango R, O Ball R, Pencharz PB: **Indicator amino acid oxidation: concept and application.** *J Nutr.* 2008; **138**(2): 243–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
69. Rakshit S, Wang YH: **The Sorghum Genome.** (ISBN 978-3-319-47787-9). Published by Springer Verlag. 2016; 318.
[Publisher Full Text](#)
70. Taylor JRN, Duodu KG: **Traditional Sorghum and Millet Food and Beverage Products and Their Technologies.** *Sorghum and Millets.* 2019; 259–292.
[Publisher Full Text](#)
71. Subramanian V, Jambunathan R, Ramaiah CE: **Physical and Chemical Characteristics of Pearl Millet Grains in Relationship to Roti Quality.** *J Food Sci Tech Mys.* 1986; 259–292.
[Reference Source](#)
72. Murty DS, Subramanian V: **Sorghum Roti: I. Traditional methods of consumption and standard procedures for evaluation.** ICRISAT Proceedings of the International Symposium on Sorghum Grain Quality. October 28–31, Hyderabad, India. 73–78.
[Reference Source](#)
73. Sarwar Gilani G, Wu Xiao C, Cockell KA: **Impact of antinutritional factors in food proteins on the digestibility of protein and the bioavailability of amino acids and on protein quality.** *Br J Nutr.* 2012; **108** Suppl 2: S315–32.
[PubMed Abstract](#) | [Publisher Full Text](#)
74. Kalasuramath S, Kurpad AV, Thankachan P: **Effect of iron status on iron absorption in different habitual meals in young south Indian women.** *Indian J Med Res.* 2013; **137**(2): 324–30.
[PubMed Abstract](#) | [Free Full Text](#)
75. Golden MHN: **The Development of Concepts of Malnutrition.** *J Nutr.* 2002; **132**(7): 2117S–2122S.
[PubMed Abstract](#) | [Publisher Full Text](#)
76. Bahwere P, Balaluka B, Wells JCK, *et al.*: **Cereals and pulse-based ready-to-use therapeutic food as an alternative to the standard milk- and peanut paste-based formulation for treating severe acute malnutrition: a noninferiority, individually randomized controlled efficacy clinical trial.** *Am J Clin Nutr.* 2016; **103**(4): 1145–1161.
[PubMed Abstract](#) | [Publisher Full Text](#)
77. Kohlmann K, Callaghan-Gillespie M, Gauglitz JM, *et al.*: **Alternative Ready-To-Use Therapeutic Food Yields Less Recovery Than the Standard for Treating Acute Malnutrition in Children From Ghana.** *Glob Heal Sci Pract.* 2019; **7**(2): 203–214.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
78. Bahwere P, Akomo P, Mwale M, *et al.*: **Soya, maize, and sorghum-based ready-to-use therapeutic food with amino acid is as efficacious as the standard milk and peanut paste-based formulation for the treatment of severe acute malnutrition in children: a noninferiority individually randomized controlled efficacy clinical trial in malawi.** *Am J Clin Nutr.* 2017; **106**(4): 1100–1112.
[PubMed Abstract](#) | [Publisher Full Text](#)
79. Dror DK, Allen LH: **Overview of Nutrients in Human Milk.** *Adv Nutr.* 2018; **9**(suppl_1): 278S–294S.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
80. DeFries R, Chhatre A, Davis KF, *et al.*: **Impact of Historical Changes in Coarse Cereals Consumption in India on Micronutrient Intake and Anemia Prevalence.** *Food Nutr Bull.* 2018; **39**(3): 377–392.
[PubMed Abstract](#) | [Publisher Full Text](#)
81. Kapil U, Jain K: **Magnitude of zinc deficiency amongst under five children in India.** *Indian J Pediatr.* 2011; **78**(9): 1069–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
82. Longvah T, Angantan I, Bhaskarachary K, *et al.*: **Indian food composition tables.** Hyderabad: National Institute of Nutrition, Indian Council of Medical Research; 2017.
[Reference Source](#)
83. Di Giovanni V, Bourdon C, Wang DX, *et al.*: **Metabolomic Changes in Serum of Children with Different Clinical Diagnoses of Malnutrition.** *J Nutr.* 2016; **146**(12): 2436–2444.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
84. Semba RD, Shardell M, Sakr Ashour FA, *et al.*: **Child Stunting is Associated with Low Circulating Essential Amino Acids.** *EBioMedicine.* 2016; **6**: 246–252.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
85. Uauy R, Suri DJ, Ghosh S, *et al.*: **Low Circulating Amino Acids and Protein Quality: An Interesting Piece in the Puzzle of Early Childhood Stunting.** *EBioMedicine.* 2016; **8**: 28–29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
86. Bora P, Ragaei S, Marcone M: **Characterisation of several types of millets as functional food ingredients.** *Int J Food Sci Nutr.* 2019; **70**(6): 714–724.
[PubMed Abstract](#) | [Publisher Full Text](#)
87. Semba RD, Trehan I, Gonzalez-Freire M, *et al.*: **The Potential Role of Essential Amino Acids and the Mechanistic Target of Rapamycin Complex 1 (mTORC1) Pathway in the Pathogenesis of Child Stunting.** *Adv Nutr.* 2016; **7**(5): 853–865.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
88. Jansson T, Aye ILMH, Goberdhan DCI: **The emerging role of mTORC1 signaling in placental nutrient-sensing.** *Placenta.* 2012; **33** Suppl 2(Suppl 2): e23–e29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
89. Saxton RA, Sabatini DM: **mTOR Signaling in Growth, Metabolism, and Disease.** *Cell.* 2017; **168**(6): 960–976.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
90. Shimobayashi M, Hall MN: **Making new contacts: the mTOR network in metabolism and signalling crosstalk.** *Nat Rev Mol Cell Biol.* 2014; **15**(3): 155–162.
[PubMed Abstract](#) | [Publisher Full Text](#)
91. Efeyan A, Comb WC, Sabatini DM: **Nutrient-sensing mechanisms and pathways.** *Nature.* 2015; **517**(7534): 302–310.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
92. Lin F, Xu W, Guan C, *et al.*: **Niacin protects against UVB radiation-induced apoptosis in cultured human skin keratinocytes.** *Int J Mol Med.* 2012; **29**(4): 593–600.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
93. Vinoth A, Ravindhran R: **Biofortification in Millets: A Sustainable Approach for Nutritional Security.** *Front Plant Sci.* 2017; **8**: 29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
94. Dimasuay KG, Boeuf P, Powell TL, *et al.*: **Placental Responses to Changes in the Maternal Environment Determine Fetal Growth.** *Front Physiol.* 2016; **7**: 12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
95. Dimasuay KG, Aitken EH, Rosario F, *et al.*: **Inhibition of placental mTOR signaling provides a link between placental malaria and reduced birthweight.** *BMC Med.* 2017; **15**(1): 1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
96. Victora CG, de Onis M, Hallal PC, *et al.*: **Worldwide Timing of Growth Faltering: Revisiting Implications for Interventions.** *Pediatrics.* 2010; **125**(3): e473–e480.
[PubMed Abstract](#) | [Publisher Full Text](#)
97. Young MF, Martorell R: **The Public Health Challenge of Early Growth Failure in India.** *Eur J Clin Nutr.* 2013; **67**(5): 596–500.
[PubMed Abstract](#) | [Publisher Full Text](#)
98. Food and Agriculture Organization of the United Nations: **Maize in Human Nutrition.** Rome; 1992.
[Reference Source](#)
99. Millward DJ: **Nutrition, infection and stunting: the roles of deficiencies of individual nutrients and foods, and of inflammation, as determinants of reduced linear growth of children.** *Nutr Res Rev.* 2017; **30**(1): 50–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
100. Roselló-Díez A, Joyner AL: **Regulation of Long Bone Growth in Vertebrates; It Is Time to Catch Up.** *Endocr Rev.* 2015; **36**(6): 646–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
101. Zeng H, Chi H: **mTOR signaling in the differentiation and function of regulatory and effector T cells.** *Curr Opin Immunol.* 2017; **46**: 103–111.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

102. Hunt JR: **Moving toward a Plant-based Diet: Are Iron and Zinc at Risk?** *Nutr Rev.* 2002; **60**(5 Pt 1): 127–134.
[PubMed Abstract](#) | [Publisher Full Text](#)
103. Sharma N, Niranjana K: **Foxtail millet: Properties, processing, health benefits, and uses.** *Food Rev Int.* 2018; **34**(4): 329–363.
[Publisher Full Text](#)
104. Bhat S, Nandini C, Tipreswamy VP: **Significance of small millets in nutrition and health-A review.** *Asian J Dairy Food Res.* 2018; **37**(1): 35–40.
[Publisher Full Text](#)
105. Basappa YC, Dinesh AS, Sanjeev RK, *et al.*: **Plots examining relationship between type of millet cultivated with stunting and wasting at district level along with map showing the overlaps for each type of millet with stunting and wasting.** *Figshare.*
<http://www.doi.org/10.6084/m9.figshare.12206135.v4>

Open Peer Review

Current Peer Review Status: ? X ?

Version 3

Reviewer Report 10 November 2021

<https://doi.org/10.21956/wellcomeopenres.18904.r46237>

© 2021 Joy E. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Edward J.M. Joy

Faculty of Epidemiology and Population Health, London School of Hygiene & Tropical Medicine, London, UK

- In the abstract, the authors suggest that 'better data' are required to confirm a causal pathway. What is actually required is a well-designed intervention study. Please revise.
- In the methods, the authors write "Other socio-demographic variables included in our analysis have been listed above...". It would be clearer to specify them here, and the authors should also specify whether these variables were controlled for in the multivariate regression analysis.
- In the methods where the outcome variables are specified, the z-scores should be 'negative 2'
- In the study limitations, the authors write "The use of cereal cultivation as a proxy for consumption too is a source of noise in our data...". I don't think 'noise' is the right term. It is likely to introduce substantial error and is a major limitation of the study.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Dietary micronutrient assessment, population micronutrient assessment

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 22 Feb 2022

Prashanth N Srinivas, Institute of Public Health Bengaluru, Bengaluru, India

Thank you for your review.

Regarding suggestion on revision of abstract, the revised abstract addresses this comment.

Regarding the clarification on variables, listing them and on the details of controlling for them in the multivariable regression, the variables have been listed in Table 1 and mention has been made of them being controlled in multivariable regression.

The error regarding z-scores is regretted and fixed.

Amendments pointed out In the study limitations has been addressed.

In addition to this, extensive revisions have now been made in the Discussion section to address other reviewer concerns as well and we hope these have comprehensively addressed the observations in the reviews.

Competing Interests: None declared

Reviewer Report 26 October 2021

<https://doi.org/10.21956/wellcomeopenres.18904.r46243>

© 2021 Kassebaum N. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Nicholas J. Kassebaum

Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA, USA

This study reports on associations between grain cultivation patterns and patterns of HIGH wasting and HIGH stunting prevalence. This is a useful addition to the field of nutrition and child anthropometrics, which ironically has not focused as much on food supply as on other factors. However, I have a few concerns:

Small points:

1. Opening sentence of intro is very difficult to read. After several tries, I'm still not sure what it is trying to say.
2. Last sentence of intro says NFHS4 was "published" in 2015. Wasn't it conducted in 2015-2016 and published in 2018?

More substantial comments:

1. Figure 1- the legend is so similar for the three categories that it is very difficult to tell them apart. Suggest to revise to increase legibility.

2. Birth weight data for children <5 were routinely collected in NFHS-4. The authors state that this is "unavailable." Please explain.
3. Choice of "high" vs. "low" is a bit arbitrary. There is nothing in the paper by Khan and Mohanty (other than their making severity classifications in a table), or indeed in the NFHS-4 anthropometrics data itself, that suggests a particular inflection point where the authors of this study have defined "high" wasting / stunting. Stunting/ wasting/ underweight are already over-simplified to be dichotomous. Further dichotomizing to high vs. low prevalence locations may be a simplification that makes the analysis easier, but it substantially weakens the broad applicability of the findings of the paper. Can the authors comment on why they were unable to perform this as an analysis where the dependent variable was not also/ instead chosen to be continuous z score?
4. Discussion doesn't talk at all about the results of the paper and its implications, instead focusing on nutrition science. Suggest to revise and focus on this paper. Summarize the biology/ clinical information much more briefly and refer interested readers to where they can find this additional information. The tables and figures in the discussion do not add to the narrative. Figure 9 in particular – these don't look substantially different; if there is an important difference to show, this needs to be a different visual (and moved to the results).
5. The most important finding of this analysis has largely been glossed over in both the results and the discussion. This is presented in Table 2 and Table 3, where the authors' multivariate analysis suggests that the previously described association between sociodemographic factors (poverty, education, rural status, open defecation) are all actually [potential] proxies for dietary diversity as evidenced by dramatically reduced – or reversed – beta coefficients in multivariate regressions. What is the authors interpretation of this finding? This contradicts, or at least refines, the findings of Khan and Mohanty (cited by authors, ref 18) and many others. If true, the implications of this are potentially profound. What are the potential policy implications? What does the literature say about the mechanistic association between sociodemographics and food supply – is it more effective to target education/ poverty/ living conditions? Or more effective to target food supply and dietary diversity? What would be the next steps in gathering data to answer this question if unknown?

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.**Reviewer Expertise:** child health, epidemiology, health metrics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 22 Feb 2022

Prashanth N Srinivas, Institute of Public Health Bengaluru, Bengaluru, India

Thank you for your detailed review.

On the minor points you have highlighted:

1. We have made suitable edits to improve readability. Regarding teh
2. Error related to year of NFHS4 regretted and corrected.

Response to major/substantive comments below; extensive changes made in v4 to address these:

1. Figure 1 has been re-done the figures to improve visual contrast of the pattern.
2. References to birth weight data from NFHS have now been removed as we do not use it in our analysis.
3. On the assertion regarding arbitrariness of "high" vs. "low" stunting/wasting districts: In the paper by Junaid and Mohanty, the districts were classified as low stunting (less than 26%), medium stunting (prevalence 26.1-46%) and high stunting (prevalence higher than 46%) based on mean and standard deviation, as per their results. Highest wasting prevalence districts were categorised as more than 28%. We utilised this classification for illustrating the districts with high prevalence of stunting, wasting as well as both high stunting and wasting in the maps. However, Z scores of less than -2 SD have been used for stunting & wasting cut off to assess prevalence in NFHS4. Their prevalence in districts (without distinction into high and low) have been used for the scatter plots and multivariable regression. A table (table 1) indicating different variables used in the maps and other results is shown for clarity in this version. We accept that our choosing of these cut-offs will affect the wider applicability of these findings. However, in the lack of a globally comparable standard, we have chosen this. Furthermore, this categorisation is only applicable to our map-based visualisations. The multivariable analysis indeed uses district-level percentages of stunting and wasting.
4. On the points raised in Discussion that are not in the results: We have re-written the Discussion section to focus better on the results reported. WE have now removed any tables and figures from our discussion and have modified earlier figure 9, a pie chart into a bar and moved it up to the results. The figure showing pathways to nutrition and information about mechanisms through MTORC1 has been only referred to and all narrative/figures on the pathway have now been shifted out of the paper.
5. On the reviewers points about findings presented in Table 2 and Table 3: Thank you

for the opportunity to clarify this further and thank you for your comments on the potential applications of our findings. In our revision, we have attempted to address these important comments to the extent possible. However, some of these comments being policy and practice implications of the underlying hypotheses we are examining, we have been unable to comprehensively address them even in our revised discussion. Moreover, many of these important observations by the reviewer are in age-groups outside of our analysis (nonetheless having important implications). We hope that our paper shall contribute to expanding the literature on the points raised by the reviewer. However, given the importance of these points, we attempt to provide some response hereunder to those aspects that are yet not addressed in our revised discussion. We hope to develop these further once this analysis achieves peer acceptability.

6. The literature and evidence on food security potentially connects socio-demographic factors mechanistically to food supply. One definition of food security is "Access by all people at all times to sufficient food and nutrition for a healthy and productive life"¹. The opposite term, food insecurity has been defined by one source as "A person is food insecure when they lack regular access to enough safe and nutritious food for normal growth and development and an active and healthy life"². Moderate food insecurity, like seen among the poor in high-income countries, is also associated with obesity³. Severe food insecurity is associated with hunger. A study based on data suitable for measuring food security from Maharashtra, India indicated that food insecurity is no longer associated with statistically significant stunting, wasting or underweight, when controlled for dietary diversity and all other child, maternal or household characteristics implying the importance of dietary diversity as a determinant of malnutrition⁴.
7. The Lancet 2013 series clearly brought that the Nutrition specific interventions, when brought up to scale of over 90% could help tackle about 20% of global burden of malnutrition⁸. Noticeably, the emphasis in the Lancet series of both 2008 & 2013 was on the first 1000 days as far as Nutrition specific interventions are concerned^{8,9}. However, longitudinal studies point to the fact that children both become stunted and recover in growth, from infancy through adolescence, thereby extending the period of intervention for healthy growth from the first 1000 days till adolescence^{10,11}. Greater persistent stunting after 5 years till 15 years, in one study in India and Ethiopia vis-à-vis Peru and Vietnam, points to an environmental context in the former countries¹¹. India also has the largest number of low birth weight among countries in the world, a reflection on the health and nutrition of women through their lives¹². A study analysing secular trends in improvement of child malnutrition indices in Vietnam and Bangladesh, attributed rapid improvements in HAZ scores to changes in underlying determinants like socioeconomic status, maternal education, hygiene and food security¹³. Changes in maternal nutritional knowledge and child dietary diversity were associated with only 20% of the total change¹². A cross country study about drivers of nutritional change suggested four factors which predict reduction in undernutrition as, women's secondary education, reduction in fertility, increased access to health services and increase in household assets¹⁴. Clearly, a combination of nutrition sensitive and specific measures as per context would be the way forward¹⁵.
8. The Integrated Child Development Services (ICDS) of India covers a wide range of

services involving teenage nutrition, pre-school feeding, health and education as well as nutrition of pregnant & breast feeding mothers, conducted through 1.3 million Anganwadi centres all over India^{16,17}. In 2006, the ICDS was scaled up substantially following a Supreme Court directive¹⁸. There was a significant improvement in ICDS coverage as seen through comparison of NFHS 3 & 4 after this scaling up. A threefold increase in percentage of women and children receiving food supplementation during pregnancy, lactation and early childhood was noted during the period^{17,18}. There were also significant improvements recorded in breast feeding practices, antenatal coverage, immunization, age of marriage, women's education and several other health related practices¹⁸. Concomitantly, a 10 percentage reduction in stunting in children aged 0-5 years was recorded in 2016 vis-à-vis 2006, as per NFHS data^{17,18}. The improvement is heterogenous across states of India, with intra-state disparities seen particularly among the poor and lower castes. However, in IYCF practices, complementary feeding and minimal meal frequency showed decline in 2016 over 2006. Clearly, the ICDS is a work in progress and there is much room for improvement with respect to coverage and efficacy^{17,18}.

9. A meta-analysis on school feedings showed significant improvements in height and weight over 12 months, compared to controls¹⁹. A recent study has clearly indicated substantial benefits of India's mid-day meal (MDM) scheme, in estimated improvement in stunting up to 0.4 SD among children born to mothers with greater exposure to MDM²⁰. This points to the intergenerational benefits of school feeding apart from its multisectoral economic benefits which extend from health, social protection, education, fertility reduction to local agriculture^{20,21}. A closer look at countries which have shown impressive declines in stunting and other mother and child nutrition related parameters, such as China and Peru, show presence of robust school feeding programs including provision of milk, over the last few decades^{22,23}. Improving the quality of school feeding (coupled with deworming, nutrition education etc) and extending it to include adolescents could arguably bring about undernutrition reduction substantially by synergising the nutrition specific and sensitive nature of interventions involved.
10. A practice like eating down in pregnancy, wherein mothers eat less for fear of labour related complications due to a large baby is not unheard of in many parts of rural India²⁴. Health care systems should be made more accessible and effective, particularly in terms of priority areas like vaccination services, as well as obstetric and child care. This will also help deal with hidden issues like maternal depression as well as common infectious causes of child wasting such as tuberculosis, Malaria, diarrhoeas and respiratory infections^{25,26}. Replacement of iron folate to multiple micronutrient supplements, targeted balanced energy micronutrient supplements to pregnant mothers with low BMI have been recommended⁸. This could be extended to breast feeding mothers given the high rates of malnutrition in early infancy in India. The MDM should be expanded to include older children with provision of a breakfast with milk to start the day. Greater emphasis on play at ICDS and school should address both child stunting and rising prevalence of obesity (twin burdens of malnutrition) well. Conditional cash transfers for poorer households like the Oportunidades in Mexico²⁷, could be used to improve dietary diversity among women and children of the poorer states through encouraging use of Nutrition, education and health services. Optimum resource allocation for multisectoral

improvements in both nutrition specific and sensitive areas with close monitoring of benefits to the targeted population nation-wide, could bring about transformational benefits.

11. Dietary matrix data like studies by Kurpad et al on commonly consumed food preparations by the poor with a focus on means to enhance nutrient availability through simple measures can be utilised for better quality nutritional advice or education²⁸. The effect of cereal type on the variability in nutrient(s) availability will have to be factored while designing/ recommending appropriate diets. Well-designed biofortification on a large scale could be explored to improve micronutrient levels in a population. Quality of breast milk, nutrient availability in the breast milk and maternal depression among mothers consuming different dietary matrices(like, say, by using different cereal types) would be immensely beneficial to focus on type of diets needed by lactating mothers in different ecogeographic areas. Given that most of the countries with high rates of malnutrition in early infancy have high cultivation of coarse cereals (our observation), it would be relevant and important to explore the relationship between type of cereal consumed and foetal growth restriction in communities with low dietary diversity and a cereal based diet. Considering that one third of Indian women in NFHS report never consuming eggs and 11% never consuming dairy products, research in optimal diet to support this group for the first 1000 days is urgently needed²⁹. Use of mobile services for personalised dietary advice to the general population is another unexplored avenue. Research into quality of India's preschool services should be extended to states with poorer indicators of performance with an aim to translate best practices consistently across the states^{30,31}.

References:

1. Gibson M. Food Security-A Commentary: What Is It and Why Is It So Complicated? Foods. 2012 Dec 3;1(1):18-27. doi: 10.3390/foods1010018. PMID: 28239088; PMCID: PMC5302220.
2. FAO, IFAD, UNICEF, WFP and WHO. 2021. *The State of Food Security and Nutrition in the World 2021. Transforming food systems for food security, improved nutrition and affordable healthy diets for all*. Rome, FAO.
3. Dhurandhar EJ. The food-insecurity obesity paradox: A resource scarcity hypothesis. Physiol Behav. 2016 Aug 1;162:88-92. doi: 10.1016/j.physbeh.2016.04.025. Epub 2016 Apr 26. PMID: 27126969; PMCID: PMC5394740.
4. Chandrasekhar S, Aguayo VM, Krishna V, Nair R. Household food insecurity and children's dietary diversity and nutrition in India. Evidence from the comprehensive nutrition survey in Maharashtra. Matern Child Nutr. 2017 Oct;13 Suppl 2(Suppl 2):e12447. doi: 10.1111/mcn.12447. PMID: 29032621; PMCID: PMC6866156.
5. Arimond M, Ruel MT. Dietary diversity is associated with child nutritional status: evidence from 11 demographic and health surveys. J Nutr. 2004 Oct;134(10):2579-85. doi: 10.1093/jn/134.10.2579. PMID: 15465751.
6. Millward DJ. Nutrition, infection and stunting: the roles of deficiencies of individual

nutrients and foods, and of inflammation, as determinants of reduced linear growth of children. *Nutr Res Rev.* 2017 Jun;30(1):50-72. doi: 10.1017/S0954422416000238. Epub 2017 Jan 23. PMID: 28112064.

7. Roselló-Díez A, Joyner AL: Regulation of Long Bone Growth in Vertebrates; It Is Time to Catch Up. *Endocr Rev.* 2015; 36(6): 646–80. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

8. Bhutta ZA, Das JK, Rizvi A, Gaffey MF, Walker N, Horton S, Webb P, Lartey A, Black RE; Lancet Nutrition Interventions Review Group, the Maternal and Child Nutrition Study Group. Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? *Lancet.* 2013 Aug 3;382(9890):452-477. doi: 10.1016/S0140-6736(13)60996-4. Epub 2013 Jun 6. Erratum in: *Lancet.* 2013 Aug 3;382(9890):396. PMID: 23746776.

9. Black RE, Allen LH, Bhutta ZA, *et al.*: Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet.* 2008; 371(9608): 243–260. [PubMed Abstract](#) | [Publisher Full Text](#)

10. Martorell R, Zongrone A. Intergenerational influences on child growth and undernutrition. *Paediatr Perinat Epidemiol.* 2012 Jul;26 Suppl 1:302-14. doi: 10.1111/j.1365-3016.2012.01298.x. PMID: 22742617.

11. Gausman J, Kim R, Subramanian SV. Stunting trajectories from post-infancy to adolescence in Ethiopia, India, Peru, and Vietnam. *Matern Child Nutr.* 2019 Oct;15(4):e12835. doi: 10.1111/mcn.12835. Epub 2019 May 23. PMID: 31042809; PMCID: PMC7938413.

12. Ramalingaswami V, Jonson U, Rohde J: The Asian Engima, the progress of nations. 1997.

13. Nguyen PH, Headey D, Frongillo EA, Tran LM, Rawat R, Ruel MT, Menon P. Changes in Underlying Determinants Explain Rapid Increases in Child Linear Growth in Alive & Thrive Study Areas between 2010 and 2014 in Bangladesh and Vietnam. *J Nutr.* 2017 Mar;147(3):462-469. doi: 10.3945/jn.116.243949. Epub 2017 Jan 25. PMID: 28122930; PMCID: PMC5320405.

14. Headey D. Developmental drivers of nutritional change: a cross-country analysis. *World Dev* 2013; 42: 76–88.

15. Ruel MT, Alderman H; Maternal and Child Nutrition Study Group. Nutrition-sensitive interventions and programmes: how can they help to accelerate progress in improving maternal and child nutrition? *Lancet.* 2013 Aug 10;382(9891):536-51. doi: 10.1016/S0140-6736(13)60843-0. Epub 2013 Jun 6. Erratum in: *Lancet.* 2013 Aug 10;382(9891):506. PMID: 23746780.

16. Rao N, Kaul V. India's integrated child development services scheme: challenges for scaling up. *Child Care Health Dev.* 2018 01;44(1):31–40. doi:

<http://dx.doi.org/10.1111/cch.12531> PMID: 29235171

17. Chakrabarti S, Raghunathan K, Alderman H, Menon P, Nguyen P. India's Integrated Child Development Services programme; equity and extent of coverage in 2006 and 2016. *Bull World Health Organ*. 2019 Apr 1;97(4):270-282. doi: 10.2471/BLT.18.221135. Epub 2019 Feb 25. PMID: 30940984; PMCID: PMC6438246.

18. Nguyen PH, Avula R, Headey D, Tran LM, Ruel MT, Menon P. Progress and inequalities in infant and young child feeding practices in India between 2006 and 2016. *Matern Child Nutr*. 2018 Nov;14 Suppl 4(Suppl 4):e12663. doi: 10.1111/mcn.12663. PMID: 30499255; PMCID: PMC6518921.

19. Wang D, Shinde S, Young T, Fawzi WW. Impacts of school feeding on educational and health outcomes of school-age children and adolescents in low- and middle-income countries: A systematic review and meta-analysis. *J Glob Health*. 2021 Sep 4;11:04051. doi: 10.7189/jogh.11.04051. PMID: 34552720; PMCID: PMC8442580.

20. Chakrabarti S, Scott SP, Alderman H, Menon P, Gilligan DO. Intergenerational nutrition benefits of India's national school feeding program. *Nat Commun*. 2021 Jul 12;12(1):4248. doi: 10.1038/s41467-021-24433-w. PMID: 34253719; PMCID: PMC8275765.

21. Verguet S, Limasalle P, Chakrabarti A, Husain A, Burbano C, Drake L, Bundy DAP. The Broader Economic Value of School Feeding Programs in Low- and Middle-Income Countries: Estimating the Multi-Sectoral Returns to Public Health, Human Capital, Social Protection, and the Local Economy. *Front Public Health*. 2020 Dec 3;8:587046. doi: 10.3389/fpubh.2020.587046. PMID: 33344398; PMCID: PMC7744483.

22. Jacoby ER, Cueto S, Pollitt E. When science and politics listen to each other: good prospects from a new school breakfast program in Peru. *Am J Clin Nutr*. 1998 Apr;67(4):795S-797S. doi: 10.1093/ajcn/67.4.795S. PMID: 9537631.

23. Huang X, Yang B, Liu Q, Zhang R, Tang S, Story M. Improving maternal and child nutrition in China: an analysis of nutrition policies and programs initiated during the 2000-2015 Millennium Development Goals era and implications for achieving the Sustainable Development Goals. *J Health Popul Nutr*. 2020 Dec 2;39(1):12. doi: 10.1186/s41043-020-00221-y. PMID: 33267909; PMCID: PMC7709233

24. Nichter M. The ethnophysiology and folk dietetics of pregnancy: a case study from South India. *Human Organization* 1983; 42:235-246.

25. Goutines J, Miller LC, Sorge F. Infections and nutritional status of internationally adopted children in France. *Acta Paediatr*. 2021 Apr;110(4):1359-1365. doi: 10.1111/apa.15612. Epub 2020 Nov 9. PMID: 33040353.

26. Shidhaye P, Giri P. Maternal depression: a hidden burden in developing countries. *Ann Med Health Sci Res*. 2014 Jul;4(4):463-5. doi: 10.4103/2141-9248.139268. PMID: 25221688; PMCID: PMC4160664.

27. . Morley, Samuel & David Coady, 2003. "From Social Assistance to Social Development: Targeted Education Subsidies in Developing Countries," Peterson Institute Press: All Books, Peterson Institute for International Economics, number cgd376, July.

28. Kalasuramath S, Kurpad AV, Thankachan P. Effect of iron status on iron absorption in different habitual meals in young south Indian women. Indian J Med Res. 2013 Feb;137(2):324-30. PMID: 23563376; PMCID: PMC3657856.

29. Nguyen PH, Avula R, Tran LM, Sethi V, Kumar A, Baswal D, Hajeebhoy N, Ranjan A, Menon P. Missed opportunities for delivering nutrition interventions in first 1000 days of life in India: insights from the National Family Health Survey, 2006 and 2016. BMJ Glob Health. 2021 Feb;6(2):e003717. doi: 10.1136/bmjgh-2020-003717. PMID: 33627359; PMCID: PMC7908280.

30. Isley, B.J. (2000). *Tamil Nadu Early Childhood Environment Rating Scale (TECERS)*. Chennai, India: M.S. Swaminathan Research Foundation.

31. Rao N. Preschool Quality and the Development of Children from Economically Disadvantaged Families in India. *Early Educ. Dev.* 2010;21:167–185. doi: 10.1080/10409281003635770.[CrossRef] [Google Scholar]

Competing Interests: None disclosed.

Version 2

Reviewer Report 09 June 2021

<https://doi.org/10.21956/wellcomeopenres.18006.r41339>

© 2021 Ghosh-Jerath S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Suparna Ghosh-Jerath

Indian Institute of Public Health-Delhi, Public Health Foundation of India, Gurgaon, Haryana, India

I would request you to kindly transfer this paper for a second round of review by another reviewer for the following reasons:

- I have gone through the responses provided by the authors. Though the authors have tried to respond to some of the concerns raised by both the reviewers, the core concern related to the basic hypothesis remains. The authors seem to be making the classic mistake of assuming causality from correlations with erroneous and troublesome conclusions. Further, these conclusions are based on several assumptions especially translating production into

consumption, limited exploration of intra-household food distribution, access to other food sources (dietary diversity), nutrition sensitive factors and a simple fact that a nutritious diet cannot be constituted or assessed based on "only cereals", one needs to look at the diet in totality and production and access to a diverse food basket. If this message is disseminated without further scrutiny, it may give strong negative messages countering the benefits of cultivating nutrient rich, climate friendly millets, that are sometimes grown as mixed cropping with other pulses/legumes and cereals, and are incidentally produced by several of the nutritionally vulnerable, marginalized, economically backward communities of India.

- Unfortunately, the authors fail to understand that production of cereal cannot be a proxy to actual consumption, which along with several other intervening factors (both nutritional as well as non nutritional) are resulting in poor nutritional status of vulnerable communities in question.
- Most of the responses are deviating from the queries raised. Removing a nutrient rich millet from the analysis and assuming that the underlying cause of wasting could be coarse cereals seems somewhat indefensible.
- In case they wish to continue with the coarse cereal /millets and wasting argument, they need to highlight in discussion that pre-processing can address most concerns about the anti-nutritional effects e.g. as given in many papers, instead of trying to dissuade millet cultivation and use. (<https://fppn.biomedcentral.com/articles/10.1186/s43014-020-0020-5>)
- In case the authors extensively revise the paper, they need to focus on all components of infant and young child feeding practices in totality along with other nutrition sensitive factors rather than focusing on "cereals only" and exploring association of wasting with cultivation of specific coarse cereals in isolation
- At its current status, the paper cannot be accepted for indexing.

References

1. Samtiya M, Aluko R, Dhewa T: Plant food anti-nutritional factors and their reduction strategies: an overview. *Food Production, Processing and Nutrition*. 2020; **2** (1). [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response 25 Jun 2021

Prashanth N Srinivas, Institute of Public Health Bengaluru, Bengaluru, India

In the paper we analyse data in order to explore the possible link between type of cereal consumed and the occurrence of child malnutrition, given that cereals constitute upwards of 70% of diet in the rural areas where malnutrition is the highest. We maintain that this is a novel exploratory study, as we have not come across a study that has explored this relationship.

In the interest of advancing critical inquiry at the heart of the scientific process, we believe that overwhelming existing evidence should NOT be a reason for non-acceptance of new kinds of evidence demonstrating plausibility if not causation. A reasonable approach in our opinion for anyone with differences from the line of thought that we pursue here would be to demonstrate mistakes in our analysis and/or over-reach of conclusions (which in fact was pointed out by RI and has been significantly revised in our version 2).

The reviewer repeatedly brings out the argument that millets are nutritious and in the same breath says that malnutrition is more common in communities consuming millets (see for eg. R2 assertion in response to v1 ""It's not surprising that households with high coarse cereals consumption have high wasting and malnutrition") This is in fact the relationship that we attempt to explain (we do not attempt to causally link the two). Such an explanation requires reaching across disciplines ranging from cellular pathways, agriculture and nutrition which we attempt in this paper.

It is the dietary matrix and how it impacts on the absorption of various nutrients that is important as opposed to making millet a singular causal candidate. Our approach is that since there is substantial variability in the dietary availability of various key nutrients in different cereals there are likely to be differences in their effects in different populations. The low glycaemic index of some millets which may be advantageous to the obese diabetic may be a disadvantage to the unborn of a pregnant adolescent with poor dietary diversity. We have indicated in our version2 that most millets indeed have greater micronutrient levels in comparison to rice & wheat which could be contributory to stunting in areas where they are predominantly consumed. Our contention is that the lower amino acid availability could be similarly causing wasting in areas where millets are consumed. To emphasize this

aspect, we have brought out the Digestible ingested amino acid scores(DIAAS), as acceptable by WHO, of different cereals from published literature. The correct approach mitigate this lack of nutritional availability by pre-processing like nixtamalization or fermentation or to supplement above by appropriate changes in the matrix or with added nutritional supplements in the way Iron and Folate is supplemented in pregnancies in most developing countries.

So what we are asking is for a closer scientific scrutiny into the nutritional availability of key components in commonly consumed diets by rural Indians. To reject the question itself, as the reviewer has done in conclusion, demonstrates the lack of engagement with our analysis and results presented possibly due to an overwhelming reliance on the existing assertions about nutritious value of millets rather than an openness to engage with evidence critical of a given hypothesis.

We believe that our assertions are reasonably demonstrated by the evidence we present and were hoping to be proved wrong by demonstrating errors in our analysis or conclusions. In the lack of that, we shall continue working harder on communicating our findings better to the scientific community.

Competing Interests: None declared.

Reviewer Report 12 January 2021

<https://doi.org/10.21956/wellcomeopenres.18006.r41340>

© 2021 Joy E. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Edward J.M. Joy

Faculty of Epidemiology and Population Health, London School of Hygiene & Tropical Medicine, London, UK

The authors have made substantial revisions which have largely addressed my earlier comments. I would prefer the authors to expand on the study limitations, which are inherent in a cross-sectional, ecological study design.

The multi-variate analysis is a better approach. It appears that the association between wheat consumption and wasting is similarly strong as for jowar consumption, which I suppose illustrates the potential pitfalls with over-interpreting the findings and implying causality via a specific metabolic pathway driven by (anti-)nutritional factors in millets. It would be consistent with an underlying driver of poverty/wealth which will determine not only access to diverse diets but also other risk factors of undernutrition.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Dietary micronutrient assessment, population micronutrient assessment

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Version 1

Reviewer Report 27 July 2020

<https://doi.org/10.21956/wellcomeopenres.17478.r39290>

© 2020 Ghosh-Jerath S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Suparna Ghosh-Jerath

Indian Institute of Public Health-Delhi, Public Health Foundation of India, Gurgaon, Haryana, India

The manuscript provides insight on impact of production of coarse cereals (selective millets) on the nutritional status of women and children in India.

I have the following concerns with the methodology and conclusions drawn in this manuscript

1. Title: The title has issues in terms of including food (cereals) and nutrients (proteins and micronutrients) in the same basket, which is not clear.

2. Hypothesis: According to the manuscript, families depending on subsistence farming and

growing millets may have higher levels of wasting in under 5 children and malnourishment in women. Assessing causality by relating production of millets to nutritional status without looking at actual consumption and other factors [underlying and basic factors (ref to UNICEF conceptual framework of Causes of maternal and child malnutrition)] may be flawed.

3. The issue of assuming causality from correlations with perhaps erroneous conclusions is a major concern in this manuscript. It is crucial to look at dietary consumption data, from NSSO and NNMB and it is also important to adjust for all the other factors like SES, endemicity to diseases, access to potable water before superimposing millet cultivation and prevalence of malnutrition in women and wasting in children. These communities are poverty stricken and have limited access to several resources mentioned above. So the yield of crop is questionable as well. What % of their farming are millets? What % of their food plate are millets? All these need to be explored before drawing conclusions. It's not surprising that households with high coarse cereals consumption have high wasting and malnutrition. To assume that the underlying cause of wasting is coarse cereals (and that wasting would be less without coarse cereals) seems erroneous and perhaps indefensible. Comparing similar households with and without coarse cereal consumption could be a convincing way to draw conclusion in the present manuscript.
4. Why have the authors excluded Ragi (finger millets) from the analysis, which is one of the most nutritious millets?
5. With Public distribution system (PDS) , a food security program in India distributing majorly rice and wheat throughout the country at subsidized rates, it might be difficult to conclude that the millet growing communities, who have access to PDS are only consuming millets and hence have poor nutritional status?
6. The authors mention that at 6 months, the wasting in children is the highest, does the data say that disaggregated data on millet cultivation and wasting at 6 months has the most significant association?
7. It is somewhat worrisome to see the concluding line stating that "Policies and programs targeting malnutrition need to address type of cereal consumed in order to impact childhood malnutrition in parts of India where subsistence cultivation of millets for staple consumption is prevalent." The millet consumption can actually add diversity to the Indian diets and the presence of anti-nutrients can be assessed in the light of anti-nutrient and micronutrient molar ratio (e.g. phytate: iron molar ration before concluding that the millets can lead to malnutrition.

Recommendation: In its current state, this manuscript may be rejected.

Is the work clearly and accurately presented and does it cite the current literature?

No

Is the study design appropriate and is the work technically sound?

No

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

No

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Food systems research, assessment of nutritional status, nutrient analysis, nutritional status of indigenous communities of India

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response 09 Oct 2020

Prashanth N Srinivas, Institute of Public Health Bengaluru, Bengaluru, India

Thank you for your careful review of our submission and for raising important concerns. We have now comprehensively revised the paper and hope that the revised version has adequately responded to your concerns raised. We first provide a comprehensive overview of the main arguments underlying our revision and subsequently provide point-by-point response to the specific concerns raised. We kindly point you to review substantive responses we have made in response to reviewer 1 under the following headings which are also relevant to concerns raised by you: (a) *On protein and amino acid content of cereals*, (b) *On the matter of why some cereals such as millet and sorghum have lower digestibility*, (c) *Protein quality in the first 1000 days of conception*, (d) *On the micronutrient content in cereals*

On the choice of title: There have been questions linked to our choice of the title by both reviewers. Overall, there are 108 districts each with higher prevalence of stunting and wasting, with an overlap of both in 18 districts (Figure 1 showing overlaps between two sets of districts in response to reviewer 1). Undeniably, cereals constitute upwards of 70% protein consumption in rural India and malnutrition has a strong rural preponderance worldwide^{1,2,3}. Looking from the prism of predominantly cereal based diets, the wide range of protein (or amino acid) and micronutrient availabilities among cereals (as explained in our overarching response above) could well determine both wasting and stunting. Furthermore the results provided in the revised version strengthen the evidence-base for an association of wasting and low BMI with millet cultivation (as a proxy for consumption).

On the hypothesis: Our hypothesis is based on the distinct patterns of higher prevalence of stunting and wasting as well as the wide diversity in the cultivation and nutritional values

of both micronutrients and proteins among cereals. We agree that other factors linked to poverty like dietary diversity, infections, low birth weight could well be contributing to malnutrition. This emphasizes a need for well-designed studies to look for the contribution of cereals consumed to the patterns of malnutrition. But, given that some states with higher poverty (e.g., UP and Bihar) with relatively lower levels of low BMI and wasting than states in peninsular (or central) India (also see Table 1 in response to Reviewer 1 and the section titled *On the early onset of malnutrition in India and ecogeographic patterns* & section titled *On the micronutrient content of cereals*) points to other factors.

On the exclusion of *ragi*(finger millet): As mentioned in the text we excluded *ragi* from the millets because of its purported nutritional richness and its distinct subfamily in the grass family *Poaceae*^{4,5,6}. This is also corroborated by patterns in the African continent where the countries which had the highest production of *ragi* (Uganda and Tanzania) are in the highlands of East Africa with lower prevalence of wasting when compared to other millet-growing regions in the continent^{4,7}. The figure showing the distribution of *Ragi* in eastern highlands is available from p.42 (Link); the geo-spatial analysis of prevalence of malnutrition in low- and middle-income countries by Kinyoki et. al figure 2 showing low prevalence of wasting in areas having pre-dominant *ragi* cultivation is available (Link).

About Poverty, PDS and food security in India and its effects on our findings (see table uploaded on Figshare as tables are not allowed in response to reviewers): That the Public Distribution system (PDS) is invaluable in managing food security in the country is undeniable. The share of rice or wheat consumption from PDS in different states indicates that the percentage is quite less (ranging from 7.6% in Gujarat to 34.3% in Chhattisgarh). The percentage of improvement in PDS use from 2004-5 to 2011-12 has not led to a commensurate decline in undernutrition in the country^{8,9}. Moreover, the coverage of PDS is much higher in hill-states of northern India and in South Indian states as compared to other regions where malnutrition prevalence is higher⁹. Hence, the likelihood of effects of PDS especially in the regions that we analyse in this paper by the marginal farmers and rural poor is likely to be low with subsistence cultivation continuing to play an important part, especially among the poorest communities.

About validity of data for study purposes: The major nutrition surveys completed in this decade are summarised by Rathi et. al. (see table 1; Link to journal)⁷. The Comprehensive National Nutrition Survey 2016-18 has data by state and their respective urban/rural areas and not by districts¹⁰. The district level household survey 4 (DLHS4 in 2012-14) covered only 336 districts among 640 present at that time. The minimum disaggregation available on such a scale is at the district level only. The malnutrition data in our study was extracted from NFHS4 which is a nationwide survey encompassing all districts^{11,12}. The Hemalatha et al¹³ study uses multiple data inputs with the latest being, NFHS4 and India Urban Nutrition Survey Data 2015-2016. As seen from table 4 above taken from Rathi et al¹⁴ NFHS4 is the only nationwide survey encompassing all districts completed in the present decade. The food consumption data too has dietary groups with cereals aggregated in a group, as grains, roots and tubers. For the purposes of our study we needed separate consumption of cereal (by type) in each district which is not available in CNNS or NFHS4. The older NSSO (NSS 55) has data on type of cereals consumed but that too is not available by district but by 77 ecogeographic zones¹⁵.

Point-by-point response to review observations

- Title: The title has issues in terms of including food (cereals) and nutrients (proteins and micronutrients) in the same basket, which is not clear.

Thank you for bringing the title to attention. We have explained above under the heading “*on the choice of the title*”.

- Hypothesis: According to the manuscript, families depending on subsistence farming and growing millets may have higher levels of wasting in under 5 children and malnourishment in women. Assessing causality by relating production of millets to nutritional status without looking at actual consumption and other factors [underlying and basic factors (ref to UNICEF conceptual framework of Causes of maternal and child malnutrition)] may be flawed.

We accept that obtaining actual consumption would have been ideal for establishing or negating such a hypothesis. In the version 2 of our paper, we have attempted to include other factors pertinent to malnutrition (available through nationwide surveys) in our analysis. We have explained this in detail under the sections *On the hypothesis* and *On the validity of data for study purposes*. Please also see the more extended multivariate analysis in version 2.

- The issue of assuming causality from correlations with perhaps erroneous conclusions is a major concern in this manuscript. It is crucial to look at dietary consumption data, from NSSO and NNMB and it is also important to adjust for all the other factors like SES, endemicity to diseases, access to potable water before superimposing millet cultivation and prevalence of malnutrition in women and wasting in children. These communities are poverty stricken and have limited access to several resources mentioned above. So the yield of crop is questionable as well. What % of their farming are millets? What % of their food plate are millets? All these need to be explored before drawing conclusions. It's not surprising that households with high coarse cereals consumption have high wasting and malnutrition. To assume that the underlying cause of wasting is coarse cereals (and that wasting would be less without coarse cereals) seems erroneous and perhaps indefensible. Comparing similar households with and without coarse cereal consumption could be a convincing way to draw conclusion in the present manuscript.

We have strengthened the analysis in the revised version as well as edited the language implying causality. We, in fact do not want to attribute causality but wish to describe a consistent pattern and provide a plausible hypothesis in the epidemiology of malnutrition in India. Hence, we have placed the pathway incorporating the mechanisms of causation in discussion rather than results in version 2. We have incorporated few of the variables pertaining to causation of malnutrition in a multivariate analysis as well. Dietary consumption data from nationwide surveys have limitations as mentioned under the section *About validity of data for study purposes*. We concede that well designed field studies are required to come to a clear position on this issue.

- Why have the authors excluded Ragi (finger millets) from the analysis, which is one of

the most nutritious millets?

We have dwelt on this under the above section titled *On the exclusion of ragi(finger millet)*.

- With Public distribution system (PDS) , a food security program in India distributing majorly rice and wheat throughout the country at subsidized rates, it might be difficult to conclude that the millet growing communities, who have access to PDS are only consuming millets and hence have poor nutritional status?

This has been explained in the section above titled *About Poverty, PDS and food security in India and its effects on our findings*.

- The authors mention that at 6 months, the wasting in children is the highest, does the data say that disaggregated data on millet cultivation and wasting at 6 months has the most significant association?

In Fig 5 of version 1 and Figs 9 & 10 of version 2 the age profile of children in the 108 high stunting and 108 high wasting districts is plotted. The districts with higher wasting (which have higher cultivation of millets as brought out in the results) have higher wasting at 6 months. However, data are unavailable to establish this at the household level due to lack of data on type of cereal consumed (which we have elaborated under the discussion).

- It is somewhat worrisome to see the concluding line stating that "Policies and programs targeting malnutrition need to address type of cereal consumed in order to impact childhood malnutrition in parts of India where subsistence cultivation of millets for staple consumption is prevalent." The millet consumption can actually add diversity to the Indian diets and the presence of anti-nutrients can be assessed in the light of anti-nutrient and micronutrient molar ratio (e.g. phytate: iron molar ration before concluding that the millets can lead to malnutrition.

We have indeed taken note of this suggestion and changed the manuscript in version 2 accordingly. We have taken into consideration the diversity of nutritional profiles of various cereals and brought out tables (Table 3 & 4) to highlight this. Further detailed explanation has been given under the headings *On protein and amino acid content of cereals*, *Protein quality in the first 1000 days of conception* and *On the micronutrient content in cereals* in the response to reviewer 1. The concluding line as stated in the comment has been removed in version 2.

References:

1. Swaminathan S, Vaz M, Kurpad AV: Protein intakes in India. *Br J Nutr*. 2012; **108**(Suppl 2): S50-S58. [PubMed Abstract](#) | [Publisher Full Text](#)
2. Manary MJ, Heikens GT, Golden M. Kwashiorkor: more hypothesis testing is needed to understand the aetiology of oedema. *Malawi Med J*. 2009 Sep;21(3):106-7. doi: 10.4314/mmj.v21i3.45630. PMID: 20345018; PMCID: PMC3717490.
3. Paciorek CJ, Stevens GA, Finucane MM, Ezzati M; Nutrition Impact Model Study Group (Child Growth). Children's height and weight in rural and urban populations in low-income and middle-income countries: a systematic analysis of population-representative data. *Lancet Glob Health*. 2013 Nov;1(5):e300-9. doi: 10.1016/S2214-

- 109X(13)70109-8. Epub 2013 Oct 24. PMID: 25104494; PMCID: PMC4547325.
4. National Research Council: Lost Crops of Africa. Washington DC National Academies Press; 1996. [Reference Source](#)
 5. Shobana S, Krishnaswamy K, Sudha V, et al.: Finger Millet (*Ragi*, *Eleusine Coracana* L.): A Review of Its Nutritional Properties, Processing, and Plausible Health Benefits. *Adv Food Nutr Res*. 2013;69:1–39. 23522794 10.1016/B978-0-12-410540-9.00001
 6. Cannarozzi G, Plaza-Wüthrich S, Esfeld K, et al.: Genome and transcriptome sequencing identifies breeding targets in the orphan crop tef (*Eragrostis tef*). *BMC Genomics*. 2014;15(1):581. 25007843 10.1186/1471-2164-15-581 4119204
 7. Kinyoki, D.K., Osgood-Zimmerman, A.E., Pickering, B.V. *et al.* Mapping child growth failure across low- and middle-income countries. *Nature* **577**, 231–234 (2020). <https://doi.org/10.1038/s41586-019-1878-8>
 8. Kumar, A., Ayyappan, S. Food Security and Public Distribution System in India. *Agric Res* **3**, 271–277 (2014). <https://doi.org/10.1007/s40003-014-0115-9>
 9. NITI Ayog. Evaluation Study on Role of Public Distribution System in Shaping Household and Nutritional Security India. Available online: http://niti.gov.in/writereaddata/files/document_publication/Final%20PDS%20Report-new.pdf. Accessed on 16 August 2020.
 10. Ministry of Health and Family Welfare (MoHFW), Government of India, UNICEF and Population Council. 2019. Comprehensive National Nutrition Survey (CNNS) National Report. New Delhi.
 11. Khan J, Mohanty SK: Spatial heterogeneity and correlates of child malnutrition in districts of India. *BMC Public Health*. 2018; **18**(1): 1027. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 12. NFHS: National Family Health Survey-4 (NFHS-4). *Ministry Heal Fam Serv India*. 2016. [Reference Source](#)
 13. India State-Level Disease Burden Initiative CGF Collaborators. Mapping of variations in child stunting, wasting and underweight within the states of India: the Global Burden of Disease Study 2000-2017. *EClinicalMedicine*. 2020;22:100317. Published 2020 May 13. doi:10.1016/j.eclinm.2020.100317
 14. Rathi K, Kamboj P, Bansal PG, Toteja GS. A review of selected nutrition & health surveys in India. *Indian J Med Res*. 2018;148(5):596-611. doi:10.4103/ijmr.IJMR_1808_18
 15. Rao PP, Birthal PS, Reddy BVS, Rai KN, Ramesh S. Diagnostics of sorghum and pearl millet grains-based nutrition in India. *SAT eJournal* 2006;2(1).

Competing Interests: None declared

Reviewer Report 21 July 2020

<https://doi.org/10.21956/wellcomeopenres.17478.r39353>

© 2020 Joy E. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Edward J.M. Joy

Faculty of Epidemiology and Population Health, London School of Hygiene & Tropical Medicine, London, UK

The authors assess whether staple cereal consumption patterns in India underlie malnutrition patterns. They test this through an association between District-level production of staple cereals, and prevalence of stunting/wasting in children or low BMI/short stature in adult women.

The article is quite clearly written and I appreciate that the authors have made the datasets and analysis code available for reviewers/readers. This is commendable.

However, there are a number of areas that concern me about the study:

The study title calls this an exploratory analysis, and if this is to be indexed I think the interpretation of the results and framing of the whole paper needs to be consistent with this. For example, the authors currently write "Policies and programs targeting malnutrition need to address type of cereal consumed in order to impact childhood malnutrition in parts of India where subsistence cultivation of millets for staple consumption is prevalent". I don't think such a statement is justified based on the findings of this paper, given the exploratory nature of the analysis, quite apart from the methodological shortcomings.

The title also specifies wasting, where 4 different anthropometric outcomes were assessed.

There are several methodological shortcomings to the analysis, in my opinion. The following shortcomings are noted by the authors:

1. Cereal production is not equivalent to cereal consumption, so testing the hypothesis using cereal production data is problematic. Can the authors explain why 'consumption' data were not used, e.g. from the Comprehensive National Nutrition Survey?
2. Data are integrated at the District-level. This level of aggregation is likely to mask the true effect, if there is indeed an effect, between millet consumption and risk of malnutrition.

The following shortcomings are not reported in the paper:

- The authors propose a biochemical pathway through which consumption of millets and sorghum might negatively affect nutritional status of women and children. However, they ignore wider contextual factors that are likely to be related both to the likelihood of producing millets and to the risk of malnutrition. These factors include, for example, socioeconomic status and likelihood of drought. There are multiple pathways linking such environmental and socioeconomic factors to nutritional status, mostly operating outside specific biochemical pathways. The proposed pathways (Figure 6) should at most be included in the discussion (not results), with acknowledgement that it may explain a small proportion of the observed association...if at all.
- The relationship between millet production and prevalence of malnutrition outcomes (displayed in Figures 3 & 4) is messy. This is not necessarily inconsistent with the hypothesis, considering the District-level nature of the data and the multiple and complex factors underlying nutritional status. However, do the authors really find sufficient evidence with appropriate statistical certainty to reject the null hypothesis, that there is no association between the independent and dependent variables? The authors need to present a more

comprehensive assessment of the associations they find, including appropriate p-values. NB I found this hard to review as the statistical test of association was not specified in the methods. Relatedly, the axes in Figures 3 & 4 need units.

- As above, assuming production closely predicts consumption is a major limitation of the study. But this could be partially addressed through limiting the analysis to rural households only. This might avoid any associations being driven by highly-urban Districts where there is both low millet production and better-than-average nutritional status.
- The authors pick out particular micronutrients and amino acids which are apparently lower in concentration or bioavailability in millets than in other staple cereals, and suggest this offers a mechanism which might underlie the observed associations. I am not clear how the authors came to select these micronutrients and amino acids. There are others, for example calcium, which are present at much greater concentrations in millets (especially finger millet) than other staple cereals. The idea that a diet with greater millet consumption is less nutritious is not supported by the evidence.

Other comments:

- Are the estimates of malnutrition prevalence consistent with others? E.g. Hemalatha et al (2020), *The Lancet*¹.

From visual analysis of the figures, it appears there are some discrepancies. The authors could discuss this.

- The pattern of increased stunting and wasting prevalence around 12 mo of age (Figure 5) is consistent with other settings and is likely due to poor IYCF practices and high rates of diarrhoea when moving from exclusive breastfeeding to complementary feeding
- The authors could consider using a different base colour for Figure 1. The current colouring might not translate well to presentations or black and white printing. Also, the hatched areas represent areas with 'higher prevalence' of stunting or wasting, but it's not clear what they are higher than, i.e. what's the comparator.

Overall, I suggest the authors re-consider the framing of the paper and presentation of findings.

References

1. India State-Level Disease Burden Initiative CGF Collaborators: Mapping of variations in child stunting, wasting and underweight within the states of India: the Global Burden of Disease Study 2000-2017. *EClinicalMedicine*. 2020; **22**: 100317 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

No

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

No

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Dietary micronutrient assessment, population micronutrient assessment

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response 09 Oct 2020

Prashanth N Srinivas, Institute of Public Health Bengaluru, Bengaluru, India

Comments of Dr Joy

Thank you for your careful review of our submission and for raising important concerns. We have now comprehensively revised the paper and hope that the revised version has adequately responded to your concerns raised. We first provide a comprehensive overview of the main arguments underlying our revision and subsequently provide point-by-point response to the specific concerns raised.

On protein and amino acid content of cereals: Cecily D Williams, in her pathbreaking description of Kwashiorkor in 1933 commented¹: *"As maize is the only source of supplementary food, some amino acid or protein deficiency cannot be excluded as a cause."* The understanding about proteins being relevant to the causation of stunting and Intrauterine growth restriction(IUGR) through the MTOR pathway has been extensively studied^{2,3,4,5}. Since proteins are absorbed through the ileum after breakdown into amino acids or short peptide chains, and, MTOR being exquisitely sensitive to amino acids, we incorporate the MTOR pathway in hypothesizing about the purported differences in wasting prevalence between northern and peninsular India (see figure 1). Further, as per the 2011 FAO Expert Consultation on Protein Quality Evaluation in Human Nutrition⁶, it is recommended that *"dietary amino acids be treated as individual nutrients and that wherever possible data for digestible or bioavailable amino acids be given in food tables on an individual amino acid basis"*. Due to paucity of human data it was further stated that *"Where human data are lacking it is recommended that true ileal amino acid digestibility values from the growing pig be used, and where these data are not available from the growing laboratory rat."*⁶ This consideration is underlying our use of ileal digestibility data in pig (Table 5 in the revised version). In a recent paper Swaminathan, Vaz & Kurpad while analysing protein intakes in India⁷, highlight lower protein quality of a predominantly cereal based rural and tribal diet, particularly in pregnancy. There are small but significant differences in the Lysine contents across cereals in their analysis (table2 in Swaminathan S et al⁷). This is in line with earlier more global evidence-base⁸. These differences are germane to the differences in malnutrition patterns between northern and peninsular India.

On the matter of why some cereals such as millet and sorghum have lower digestibility, Millward implicates tougher plant cell walls prevalent in millets; others have implicated anti-nutrient factors as well^{8,9,10,11}. This is corroborated by measures such as the Digestible indispensable amino acid scores (DIAAS) of cooked cereals (including rice and wheat) which are the lowest for foxtail and proso millet¹². Other independent analyses to determine the standardised ileal digestibility (SID) among eight cereal grains using DIAAS scores also have shown highest values for rice but the lowest for maize, rye and Sorghum¹³. Furthermore, excess Leucine in Sorghum has been implicated in Pellagra reported among people who consume Sorghum as a staple in India¹⁴. There are few studies that compare ileal digestibility across various complementary foods in India. In a comparison of standardized ileal digestibility (%) of amino acids between Sorghum, pearl millet and different varieties of corn, Sorghum was the lowest followed by pearl millet. The different corn varieties were higher than both¹⁵. The digestibility of pearl millet protein has been suspected to be less than other major grains¹⁶. While evaluating the protein quality of complementary foods using dual isotope tracer method in comparing the true ileal digestibility of rice, finger millet (*ragi*) and egg, finger millet and mung dal had the lowest¹³. Studies are ongoing for other millets but are currently unavailable (email communication with corresponding author¹⁷).

Protein quality in the first 1000 days of conception. Pregnant women in fact need an additional 1, 9 and 31 gm per day in first, second and third trimester respectively¹⁸. This lack of protein (and energy quality) is linked to low maternal BMI and Intrauterine growth restriction¹⁸. As per the standard FAO reference source on millets and sorghum, Lysine amino acid score of pearl millet is variable (26-69)⁸. Barnyard millet, little millet and Sorghum had the lowest Lysine scores among millets⁸. The lower glycaemic index of millets vis-à-vis rice and wheat (which is indeed beneficial for elderly Diabetics), on the other hand for pregnant women on pre-dominant millet-based diet could contribute to low birth weight and wasting since glucose along with amino acids is an important upstream regulator for MTOR^{4,19}. In fact the FAO asserts that exclusive millet diets are not adequate to meet the growth requirements of infants and young children⁸. In table 5 (of the revised version of our paper) we provide Lysine scores of cereals and their references in studies conducted in University of Illinois, Urbana-Champaign and elsewhere. The table is extracted from a compendium curated by Hans-Henrik Stein from published data.

Based on these arguments and evidence presented in our paper further supported by extensive published data, we submit that monotonous cereal based diets of the rural poor wherever they are not supplemented with good quality protein from other sources could drive wasting, as is seen in monotonous millet-based diets.

On the micronutrient content in cereals: Micronutrient availability in rice and wheat, on the contrary, is worse off than in millets (see table 6 of revised paper). This could be underlying the higher prevalence of stunting in rice and wheat growing areas of northern India (possibly related to Zinc deficiency). Review evidence too corroborates the other reviewer's observation that high phytate and phytate to mineral molar ratios in plant based diets as contributing to deficiencies of Iron, zinc and Calcium^{20,21,22,23}. The higher Iron, Zinc and Calcium content in millets (especially pearl millet, Sorghum and other millets) could be offsetting the phytate and phenol inhibitors in comparison to rice and wheat (Table 6 data

from Indian Food Composition Table 2017²⁴). This underlies our assertion “Lower lysine, high phytates with lowered micronutrients like zinc and iron in cereal based diet in wheat and rice growing areas with poor dietary diversity could lead to stunting” (as per caption of pathway figure in version 1; now figure 11). Zinc has a known association with linear growth and supplementation in developing countries has been beneficial in marginal gain of length^{25,26}. There has been a decline in the zinc & Iron molar ratios with phytates, in India over the last four decades, attributed to lesser consumption of millets and sorghum^{21,22}. Characterising micronutrient availabilities *in vivo* (particularly Zinc, Vitamin D, Magnesium, Phosphate, Selenium) in different cereal based meals will require state-of-the-art laboratory techniques like stable isotope studies. Iodine is another micronutrient critical for foetal and child growth, but, unlikely to be of consequence in view of universal Iodization of common salt in India. Iodine availability of different cereals is however unavailable in IFCT 2017.

Malnutrition prevalence related to micronutrients are available only at state level and with no district level data and have been compiled by us from different sources in table 1 below. The state-level data do not reproduce intuitive patterns. For example, poorer states (higher poverty rates) are better off with respect to prevalence of low BMI among women in 10-19 years age group when compared with richer states which report higher coarse cereal cultivation (Rajasthan, Maharashtra, Karnataka and Gujarat highlighted in red). Similarly, lesser degree of low zinc prevalence was seen in 1-4 year age groups in Rajasthan and Maharashtra (see discussion above on better off micronutrient profile of coarse cereals) in comparison to Uttar Pradesh and Bihar. However, these patterns are not consistent (for example Karnataka which reports higher low zinc prevalence in 1-4 year age group despite having relatively high coarse cereal consumption). Nevertheless, a study comparing zinc levels among preschool children across five states of India too showed higher prevalence in Orissa(51.3%) followed by Uttar Pradesh(48.1%), Gujarat(44.2%), Madhya Pradesh(38.9%) and Karnataka(36.2%)²⁷. The latter three have a higher production of coarse cereals in comparison to others as seen in the table ([since tables are not allowed here, this table curated by us has been uploaded on figshare](#))

On the early onset of malnutrition in India and ecogeographic patterns: The timing of onset of malnutrition in India has been reported in earlier studies. The paper titled *The Asian Enigma* by UNICEF in 1996 brought it to the forefront with a discussion about higher proportion of low birth weights in India (and Bangladesh) with its linkages to women's health in general and maternal nutrition in particular²⁸. Cesar Vitoria et al²⁹ have demonstrated this with India having the lowest weight for age Z scores at 1 month of age. This aspect was emphasized by Martorell et al in the comparisons between stunting and wasting prevalence as well as their respective age of occurrence between India and Guatemala³⁰. [Figure 1 in the paper by Martorell](#) illustrates this comparison indicating earlier onset of wasting in India.

In our analysis the stunting prevalence comparison between the districts with high stunting and high wasting showed a similar pattern as seen in figures 9 and 10 (in the revised version), with higher and earlier onset of wasting in the 108 high wasting districts. That is likely to be attributable to maternal nutritional factors during pregnancy, lactation and low birth weight. However, granular data on low birth weight by district in India not being available, this remains a hypothesis to be tested when better data becomes available.

Further evidence of ecogeographic patterning of malnutrition is also available in the landmark Lancet 2008 series on malnutrition³¹ where it is asserted *"Furthermore, stunting and severe wasting are not necessarily associated on a geographical or ecological basis—ie, countries with a similar stunting prevalence can have a several-fold difference in the prevalence of severe wasting."* This phenomenon has also been written about by Cesar Victora in 1992³². What we are attempting to posit here is that, in a diverse country like India, such differences are likely to be important in addressing malnutrition.

Point-by-point response to review observations

- The authors assess whether staple cereal consumption patterns in India underlie malnutrition patterns. They test this through an association between District-level production of staple cereals, and prevalence of stunting/wasting in children or low BMI/short stature in adult women. The article is quite clearly written and I appreciate that the authors have made the datasets and analysis code available for reviewers/readers. This is commendable.

Thank you.

- The study title calls this an exploratory analysis, and if this is to be indexed I think the interpretation of the results and framing of the whole paper needs to be consistent with this. For example, the authors currently write "Policies and programs targeting malnutrition need to address type of cereal consumed in order to impact childhood malnutrition in parts of India where subsistence cultivation of millets for staple consumption is prevalent". I don't think such a statement is justified based on the findings of this paper, given the exploratory nature of the analysis, quite apart from the methodological shortcomings. The title also specifies wasting, where 4 different anthropometric outcomes were assessed.

There have been questions linked to our choice of the title by both reviewers. Overall, there are 108 districts each with higher prevalence of stunting and wasting, with an overlap of both only in 18 districts (see Junaid et. al.; reference number 33). Undeniably, cereals constitute upwards of 70% protein consumption in rural India and malnutrition has a strong rural preponderance worldwide⁷. Looking from the prism of predominantly cereal based diets, the wide range of protein (or amino acid) and micronutrient availabilities among cereals (as explained in our overarching response above) could well determine both wasting and stunting. Furthermore the results provided in the revised version strengthen the evidence-base for an association of wasting and low BMI with millet cultivation (as a proxy for consumption).

- There are several methodological shortcomings to the analysis, in my opinion. The following shortcomings are noted by the authors: Cereal production is not equivalent to cereal consumption, so testing the hypothesis using cereal production data is problematic. Can the authors explain why 'consumption' data were not used, e.g. from the Comprehensive National Nutrition Survey? Data are integrated at the District-level. This level of aggregation is likely to mask the true effect, if there is indeed an effect, between millet consumption and risk of malnutrition.

This is an important shortcoming of the paper. However, no consumption data are currently available. Detailed explanation for this is provided in response to reviewer 2 under the

heading *About validity of data for study purposes*.

- The following shortcomings are not reported in the paper: The authors propose a biochemical pathway through which consumption of millets and sorghum might negatively affect nutritional status of women and children. However, they ignore wider contextual factors that are likely to be related both to the likelihood of producing millets and to the risk of malnutrition. These factors include, for example, socioeconomic status and likelihood of drought. There are multiple pathways linking such environmental and socioeconomic factors to nutritional status, mostly operating outside specific biochemical pathways. The proposed pathways (Figure 6) should at most be included in the discussion (not results), with acknowledgement that it may explain a small proportion of the observed association...if at all.

We agree with the reviewer observation about the need to integrate other known co-variates of malnutrition which is now done in version 2. Furthermore as suggested the implications of our findings are moved to the discussion (figure 11 relating to plausible pathways is now moved to discussion).

- The relationship between millet production and prevalence of malnutrition outcomes (displayed in Figures 3 & 4) is messy. This is not necessarily inconsistent with the hypothesis, considering the District-level nature of the data and the multiple and complex factors underlying nutritional status. However, do the authors really find sufficient evidence with appropriate statistical certainty to reject the null hypothesis, that there is no association between the independent and dependent variables? The authors need to present a more comprehensive assessment of the associations they find, including appropriate p-values. NB I found this hard to review as the statistical test of association was not specified in the methods. Relatedly, the axes in Figures 3 & 4 need units.

Thank you for these observations. We have now reported the details of our statistical analysis. Corresponding changes have also been made in erstwhile figures 3 and 4 which have now been broken up into figures 3-5 & 6-8 in the revised version along with appropriate p-values.

- As above, assuming production closely predicts consumption is a major limitation of the study. But this could be partially addressed through limiting the analysis to rural households only. This might avoid any associations being driven by highly-urban Districts where there is both low millet production and better-than-average nutritional status.

The multivariate analysis has now taken into the account the effect of rural population proportion in the district.

- The authors pick out particular micronutrients and amino acids which are apparently lower in concentration or bioavailability in millets than in other staple cereals, and suggest this offers a mechanism which might underlie the observed associations. I am not clear how the authors came to select these micronutrients and amino acids. There are others, for example calcium, which are present at much greater concentrations in millets (especially finger millet) than other staple cereals.

This has now been comprehensively addressed in our overarching response above (see particularly discussion under sub-heading *On the matter of why some cereals such as millet*

and sorghum have lower digestibility and On micronutrient availability in cereals. Please see also section On the exclusion of ragi in response to reviewer 2).

- The idea that a diet with greater millet consumption is less nutritious is not supported by the evidence.

This has been addressed in the revisions made. Please also refer to section above titled *On micronutrient availability in cereals*.

- Are the estimates of malnutrition prevalence consistent with others? E.g. Hemalatha et al (2020), The Lancet⁶⁶.

Hemalatha et. al.⁶⁶ have used multiple data sources for their analysis, of which the only comprehensive dataset that comprises district-level malnutrition data for the entire country is NFHS4 (which we have used in our analysis). Other datasets used in their paper do not provide district-level data for the entire country. Our estimates of malnutrition prevalence at the district level are reproducible and are consistent with analysis of NFHS4.

- From visual analysis of the figures, it appears there are some discrepancies. The authors could discuss this.

The maps have been reproduced with better contrast and legends for clarity. The few instances of lack of overlaps between millet cultivation and wasting (e.g., bajra in north India) are now discussed in the revised version.

- The pattern of increased stunting and wasting prevalence around 12 mo of age (Figure 5) is consistent with other settings and is likely due to poor IYCF practices and high rates of diarrhoea when moving from exclusive breastfeeding to complementary feeding

We have extensively discussed the early onset of malnutrition in India above.

- The authors could consider using a different base colour for Figure 1. The current colouring might not translate well to presentations or black and white printing. Also, the hatched areas represent areas with 'higher prevalence' of stunting or wasting, but it's not clear what they are higher than, i.e. what's the comparator.

Thank you for these suggestions. The revised figures address these concerns. We took the levels of districts having higher prevalence of stunting >45% and wasting >27% based on cut-offs suggested by Junaid et al article³³. We used levels of > = 30% and >= 15% as high prevalence of women's BMI (<18.5) and short stature(< 145 cms) respectively.

References:

1. Williams CD: A nutritional disease of childhood associated with a maize diet. *Arch Dis Child*. 1933;8(48):423–433. 21031941 10.1136/adc.8.48.423 1975318
2. Semba RD: The Rise and Fall of Protein Malnutrition in Global Health. *Ann Nutr Metab*. 2016; **69**(2): 79–88. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Semba RD, Shardell M, Sakr Ashour FA, et al.: Child Stunting is Associated with Low Circulating Essential Amino Acids. *EBioMedicine*. 2016; **6**: 246–252. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Jansson T, Aye ILMH, Goberdhan DCI: The emerging role of mTORC1 signaling in placental nutrient-sensing. *Placenta*. 2012; **33**(Suppl 2): e23–e29. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

5. Millward DJ. Nutrition, infection and stunting: the roles of deficiencies of individual nutrients and foods, and of inflammation, as determinants of reduced linear growth of children. *Nutr Res Rev.* 2017;30(1):50-72. doi:10.1017/S0954422416000238.
6. Dietary protein quality evaluation in human nutrition. Report of an FAO Expert Consultation. *FAO Food Nutr Pap.* 2013;92:1-66.
7. Swaminathan S, Vaz M, Kurpad AV: Protein intakes in India. *Br J Nutr.* 2012; **108**(Suppl 2): S50–S58. [PubMed Abstract](#) | [Publisher Full Text](#)
8. Food and Agriculture Organization of the United Nations: Sorghum and millets in human nutrition. David Lubin Meml Libr Cat Publ Data FAO, Rome. 1995.
9. Millward DJ. The nutritional value of plant-based diets in relation to human amino acid and protein requirements. Proceedings of the Nutrition Society. Cambridge University Press; 1999;58(2):249–60.
10. Boncompagni E, Orozco-Arroyo G, Cominelli E, et al.: Antinutritional factors in pearl millet grains: Phytate and goitrogens content variability and molecular characterization of genes involved in their pathways. Kashkush K, ed. *PLoS One.* 2018;13(6):e0198394. 29856884 10.1371/journal.pone.0198394 5983567
11. Proietti I, Frazzoli C, Mantovani A: Exploiting Nutritional Value of Staple Foods in the World's Semi-Arid Areas: Risks, Benefits, Challenges and Opportunities of Sorghum. *Healthcare.* 2015;3(2):172–193. 27417755 10.3390/healthcare3020172 4939534
12. Han F, Han F, Wang Y, et al. Digestible indispensable amino acid scores of nine cooked cereal grains. *Br J Nutr.* 2019;121(1):30-41. doi:10.1017/S0007114518003033
13. Cervantes-Pahm SK, Liu Y, Stein HH. Digestible indispensable amino acid score and digestible amino acids in eight cereal grains. *Br J Nutr.* 2014;111(9):1663-1672. doi:10.1017/S0007114513004273
14. Gopalan C, Jaya Rao KS. Pellagra and amino acids imbalance. *Vitam Horm.* 1975;33:505-528. doi:10.1016/s0083-6729(08)60971-1 Yin, Y.-L., Gurung, Nar, K., Jeaurond, E. A., Sharpe, P. H. and de Lange, C. F. M. 2002.
15. Digestible energy and amino acid contents in Canadian varieties of sorghum, pearl millet, high-oil corn, high-oil-high-protein corn and regular corn samples for growing pigs. *Can. J. Anim. Sci.* **82**: 385–391
16. National Research Council: Lost Crops of Africa. Washington DC National Academies Press; 1996. [Reference Source](#)
17. Nirupama Shivakumar, Sindhu Kashyap, Satvik Kishore, Tinku Thomas, Aneesia Varkey, Sarita Devi, Thomas Preston, Farook Jahoor, M S Sheshshayee, Anura V Kurpad, Protein-quality evaluation of complementary foods in Indian children, *The American Journal of Clinical Nutrition*, Volume 109, Issue 5, May 2019, Pages 1319–1327, <https://doi.org/10.1093/ajcn/nqy265>
18. Ghosh S. Protein Quality in the First Thousand Days of Life. *Food Nutr Bull.* 2016;37 Suppl 1:S14-S21. doi:10.1177/0379572116629259
19. Atkinson FS, Foster-Powell K, Brand-Miller JC. International tables of glycemic index and glycemic load values: 2008. *Diabetes Care.* 2008;31(12):2281-2283. doi:10.2337/dc08-1239
20. Wessells KR, Brown KH. Estimating the global prevalence of zinc deficiency: results based on zinc availability in national food supplies and the prevalence of stunting. *PLoS One.* 2012;7(11):e50568. doi:10.1371/journal.pone.0050568
21. DeFries R, Chhatre A, Davis KF, et al. Impact of Historical Changes in Coarse Cereals Consumption in India on Micronutrient Intake and Anemia Prevalence. *Food Nutr Bull.*

- 2018;39(3):377-392. doi:10.1177/0379572118783492
22. Smith MR, DeFries R, Chhatre A, Ghosh-Jerath S, Myers SS. Inadequate Zinc Intake in India: Past, Present, and Future. *Food Nutr Bull.* 2019;40(1):26-40. doi:10.1177/0379572118825176
23. Gibson RS, Bailey KB, Gibbs M, Ferguson EL. A review of phytate, iron, zinc, and calcium concentrations in plant-based complementary foods used in low-income countries and implications for bioavailability. *Food Nutr Bull.* 2010;31(2 Suppl):S134-S146. doi:10.1177/15648265100312S206.
24. Indian Food Composition Tables 2017 T. Longvah, R. Ananthan, K. Bhaskarachary and K. Venkaiah
25. Imdad, A., Bhutta, Z.A. Effect of preventive zinc supplementation on linear growth in children under 5 years of age in developing countries: a meta-analysis of studies for input to the lives saved tool. *BMC Public Health* **11**, S22 (2011). <https://doi.org/10.1186/1471-2458-11-S3-S22>
26. Millward DJ. Nutrition, infection and stunting: the roles of deficiencies of individual nutrients and foods, and of inflammation, as determinants of reduced linear growth of children. *Nutr Res Rev.* 2017;30(1):50-72. doi:10.1017/S0954422416000238.
27. Kapil U, Jain K. Magnitude of zinc deficiency amongst under five children in India. *Indian J Pediatr* 2011; 78 : 1069-72.
28. Ramalingaswami V, Jonson U, Rohde J: The Asian Engima, the progress of nations. 1997.
29. Victora CG, de Onis M, Hallal PC, *et al.*: Worldwide Timing of Growth Faltering: Revisiting Implications for Interventions. *Pediatrics.* 2010; **125**(3): e473–e480. [PubMed Abstract](#) | [Publisher Full Text](#)
30. Martorell R, Young MF: Patterns of Stunting and Wasting: Potential Explanatory Factors. *Adv Nutr.* 2012; **3**(2): 227–233. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Black RE, Allen LH, Bhutta ZA, *et al.* Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet.* 2008;371(9608):243-260. doi:10.1016/S0140-6736(07)61690-0.
32. Victora CG. The association between wasting and stunting: an international perspective. *J Nutr* 1992; 122: 1105–10.
33. Khan J, Mohanty SK: Spatial heterogeneity and correlates of child malnutrition in districts of India. *BMC Public Health.* 2018;18(1):1027. 30119652 10.1186/s12889-018-5873-z 6098604

Competing Interests: Nil

Comments on this article

Version 3

Reader Comment 06 Jan 2022

Vidya Ramkumar, Sri Ramachandra Institute of Higher Education and Research (Deemed to be University), Chennai, India

The use of geomaps in this article was of interest to me. Upon reading I was intrigued by some of the concepts that forms the basis of the understanding of malnutrition.

Presumably, the authors chose to look at cereal cultivation owing to the large-scale nature of its production. Would things have been different if we grew more (proportionately), fruits, vegetables and nuts? Considering a plethora of micro and macro nutrients are available from these sources and can also increase overall sustainable eating and food habits. Nourishing food has to be easy, local, sustainable for it become a part of routine consumption. A study mapping consumption of natural, local foods and nutrition of children/mothers would be interesting.

Competing Interests: None