METHOD ARTICLE

REVISED Insecticide resistance selection and reversal in two strains of *Aedes aegypti* [version 2; peer review: 2 approved]

Jonathan Thornton^{1*}, Bruno Gomes^{id}1,2*, Constância Ayres³, Lisa Reimer^{id}1

¹Vector Biology Department, Liverpool School of Tropical Medicine, Liverpool, L35QA, UK

²Laboratório de Bioquímica e Fisiologia de Insetos, Oswaldo Cruz Institute (IOC-FIOCRUZ), Rio de Janeiro, 21040-360, Brazil

³Aggeu Magalhães Institute, Oswaldo Cruz Foundation (IAM-FIOCRUZ), Recife, Brazil

* Equal contributors

v2 First published: 31 Jul 2020, 5:183
<https://doi.org/10.12688/wellcomeopenres.15974.1>

Latest published: 16 Dec 2020, 5:183
<https://doi.org/10.12688/wellcomeopenres.15974.2>

Abstract

Background: Laboratory reared mosquito colonies are essential tools to understand insecticide action. However, they differ considerably from wild populations and from each other depending on their origin and rearing conditions, which makes studying the effects of specific resistance mechanisms difficult. This paper describes our methods for establishing multiple resistant strains of *Aedes aegypti* from two colonies as a new resource for further research on metabolic and target site resistance.

Methods: Two resistant colonies of *Ae. aegypti*, from Cayman and Recife, were selected through 10 generations of exposure to insecticides including permethrin, malathion and temephos, to yield eight strains with different profiles of resistance due to either target site or metabolic resistance. Resistance ratios for each insecticide were calculated for the selected and unselected strains. The frequency of *kdr* alleles (F1534C and V1016I) in the Cayman strains was determined using TaqMan assays. A comparative gene expression analysis among Recife strains was conducted using qPCR in larvae (CCae3A, CYP6N12, CYP6F3, CYP9M9) and adults (CCae3A, CYP6N12, CYP6BB2, CYP9J28a).

Results: In the selected strain of Cayman, mortality against permethrin reduced almost to 0% and *kdr* became fixated by 5 generations. A similar phenotype was seen in the unselected homozygous resistant colony, whilst mortality in the susceptible homozygous colony rose to 82.9%. The Recife strains showed different responses between exposure to adulticide and larvicide, with detoxification genes in the temephos selected strain staying similar to the baseline, but a reduction in detoxification genes displayed in the other strains.

Conclusions: These selected strains, with a range of insecticide resistance phenotypes and genotypes, will support further research on the effects of target-site and/or metabolic resistance mechanisms

Open Peer Review

Approval Status

	1	2
version 2		
(revision)
16 Dec 2020	view	view

version 1
31 Jul 2020	view	

1. **Leticia Smith** ^{id}, National Institutes of Health, Rockville, USA

2. **Qian Han** ^{id}, Laboratory of Tropical Veterinary Medicine and Vector Biology, School of Life and Pharmaceutical Sciences, Hainan University, Haikou, China

Any reports and responses or comments on the article can be found at the end of the article.

on various life-history traits, behaviours and vector competence of this important arbovirus vector.

Keywords

Aedes aegypti, insecticide resistance, mosquito, pyrethroid, permethrin, temephos, malathion, kdr,

Corresponding author: Lisa Reimer (lisa.reimer@lstm.ac.uk)

Author roles: **Thornton J:** Investigation, Methodology, Project Administration, Writing – Original Draft Preparation; **Gomes B:** Conceptualization, Formal Analysis, Methodology, Writing – Review & Editing; **Ayres C:** Methodology, Resources, Writing – Review & Editing; **Reimer L:** Conceptualization, Funding Acquisition, Methodology, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome [202137].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Thornton J *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Thornton J, Gomes B, Ayres C and Reimer L. **Insecticide resistance selection and reversal in two strains of *Aedes aegypti* [version 2; peer review: 2 approved]** Wellcome Open Research 2020, 5:183 <https://doi.org/10.12688/wellcomeopenres.15974.2>

First published: 31 Jul 2020, 5:183 <https://doi.org/10.12688/wellcomeopenres.15974.1>

REVISED Amendments from Version 1

We have made additions to the manuscript to clarify the scope, re-analyse the data and we have provided a few additional methodological details and rationale.

Any further responses from the reviewers can be found at the end of the article

Introduction

Aedes aegypti is one of the most significant mosquito species of public health concern due to its role as a vector of several arboviruses, including dengue, yellow fever, chikungunya and Zika. Over 4 million disability-adjusted life years worldwide were attributed to mosquito-borne viruses in 2013 (Moyes *et al.*, 2017). Dengue virus, the most ubiquitous arbovirus, is found in 128 countries across temperate and tropical regions, and 3.9 billion people currently live at risk of infection (Guzman & Harris, 2015; Pollett *et al.*, 2018). Yellow fever has re-emerged as an important disease in Africa and the Americas, with outbreaks occurring in regions that previously had low vaccination coverage and low-to-zero yellow fever incidence (Douam & Ploss, 2018; Monath & Vasconcelos, 2015). Large outbreaks of chikungunya virus have also been described since 2000 (Burt *et al.*, 2017). Meanwhile, the World Health Organization declared a “public health emergency of international concern” during the Zika virus epidemic of 2015 and following the discovery of its association with microcephaly (Kindhauser *et al.*, 2016).

Vector control is the primary strategy to prevent transmission of arboviruses due to the absence of prophylactic drugs or vaccines for most diseases (Silva *et al.*, 2018). Chemical insecticides, biological agents, and habitat management (George *et al.*, 2015; Horstick *et al.*, 2010) are three common methods of controlling *Aedes* spp. However, insecticide resistant *Ae. aegypti* are commonly reported in Latin America and southern Asia, and have been reported in Africa (Moyes *et al.*, 2017; Vontas *et al.*, 2012), threatening the efficacy of vector-borne disease control programs (Corbel *et al.*, 2017; Moyes *et al.*, 2017; Ranson *et al.*, 2010).

Experimental studies comparing the attributes of susceptible and resistant mosquito colonies are crucial to elucidate resistance mechanisms (Davies *et al.*, 2008; Feyereisen *et al.*, 2015; Ingham *et al.*, 2018; Li *et al.*, 2007; Melo-Santos *et al.*, 2010; Moyes *et al.*, 2017; Poupardin *et al.*, 2014; Ranson *et al.*, 2010; Rinkevich *et al.*, 2013; Stevenson *et al.*, 2012; Strode *et al.*, 2012; Vontas *et al.*, 2012; Weetman *et al.*, 2018), insecticide mode of action (Ffrench-Constant, 2013; Yunta *et al.*, 2016), and the fitness costs of resistance (Brito *et al.*, 2013; Diniz *et al.*, 2015). This information is necessary to develop new insecticide formulations and alternative control methods that avoid cross-resistance (de Lourdes Macoris *et al.*, 2018; Jankowska *et al.*, 2017). However, differences between susceptible and resistant colonies due to differing genetic backgrounds – caused by bottlenecks or genetic drift – may influence study outcomes (Ffrench-Constant, 2013). The ability to select sub-strains

from a single parent colony which exhibit phenotypically distinct insecticide susceptibility profiles could help address the limitations of using disparate colonies. The aim of our study was to select multiple strains of *Ae. aegypti*, through exposure to a range of insecticides, that vary in resistance phenotype. Here we present the resistant phenotype after ten generations of insecticide exposure using two parent colonies: CAYMAN, a pyrethroid-resistant colony conferred by two target-site mutations in the sodium channel gene (V1016I and F1534C) (Harris *et al.*, 2010), and RECIFE, a temephos-resistant colony conferred by over-expression of multiple detoxification genes (Diniz *et al.*, 2015). Additionally, we present differences in a select number of resistant alleles and metabolic genes which can be used to inform further research.

Methods

Summary of the study design

A Total of three colonies from *Aedes aegypti* were used in this study:

Colony CAYMAN (CAY) was originally established in 2008 with *Aedes aegypti* collected in Grand Cayman island (Caribbean). This colony is highly resistant to pyrethroids and DDT, attributed to *kdr* alleles (F1534C and V1016I), and has been routinely selected with 0.75 % permethrin for 1 hour in the Liverpool Insect Testing Establishment (LITE: <https://lite.lstmed.ac.uk/lite-facilities/lite-insectaries/aedes-aegypti-cayman>).

Colony RECIFE (REC) was originally established in 2004 with *Aedes aegypti* collected in Araripina, Brazil (7° 32' S and 40° 34' W; Melo-Santos *et al.*, 2010). This colony is resistant to temephos (OP), and biochemical assays indicate a higher activity of multiple detox enzyme families. Larvae from this colony have been routinely selected with 0.5 mg/L temephos for 24 hours.

Colony NEW ORLEANS is a susceptible colony established at LSTM in the 1970s. This colony originated in New Orleans (USA) and it is maintained in laboratory without insecticide exposure. Routine screening for target-site mutations associated with resistance indicates a lack of *kdr* alleles in the sodium channel and resistant alleles in Ace-1 (LITE: <https://lite.lstmed.ac.uk/lite-facilities/lite-insectaries/aedes-aegypti-new-orleans>).

We exposed mosquitoes from CAY and REC colonies to different insecticide selection pressures and monitored key indicators of resistance over time (Figure 1). We established the following eight strains of *Ae. aegypti*: CAY-P exposed to permethrin, CAY-RR unexposed and homozygous for resistance alleles (V1016I and F1534C), CAY-RS unexposed and heterozygous for resistance alleles, CAY-SS unexposed and homozygous for susceptible alleles, REC-R exposed to temephos, REC-M exposed to malathion, REC-P exposed to permethrin, and REC-U unexposed.

First, discriminating concentration assays were conducted to identify a suitable concentration of permethrin, malathion and temephos for the selection procedure. Second, mosquitoes were exposed to an insecticide selection regime as outlined

Figure 1. Experiment outline. Strains used were Cayman, which has target-site resistance to pyrethroids/DDT, and Recife, which has metabolic resistance to temephos. LC50s were determined by discriminating dose assays, then the LC50 was used to select the strain. Cayman was split into a strain selected with permethrin and an unselected strain, which was manually selected into R and S homozygotes and then crossed with reciprocal homozygotes to produce heterozygote strains. Recife was selected with malathion and permethrin, one strain was maintained with temephos exposure every three generations and one strain was left unselected. Each strain was subjected to bioassays using the WHO tube assay at the WHO recommended concentrations every three generations, and nucleic acid extraction and analysis of *kdr* alleles or detoxification genes was performed. At the end of 10 generations, each strain underwent bioassays to determine the LC50 and LC95 compared to the susceptible strain.

in Figure 1. Every three generations during the selection process, we monitored phenotypic resistance according to WHO diagnostic concentrations and we assessed the presence of *kdr* alleles (CAY) or the upregulation of a select panel of detox genes (REC). Finally, resistance ratios for the eight strains were compared against the fully susceptible colony (NO). Details of these procedures are described below.

Initial Discriminating Concentration assays in larvae and adults

Discriminating concentration assays using WHO tubes were conducted to determine an appropriate concentration of insecticide to expose each colony. The criteria is based on LC_{50} to provide insecticide pressure, but within concentrations that mosquitoes are likely to encounter in the field. Papers impregnated with

insecticide were prepared according to the standard WHO protocol (WHO, 2013). Filter papers of 12 cm × 15 cm were impregnated with a 1:1:1 volume mix of insecticide, acetone and corning oil and left to dry in a fume hood for 24 hours. Papers were stored wrapped in aluminium foil and placed in plastic bags at -20°C and used in up to five assays or within six months. The adult mosquitoes were selected with either permethrin (Sigma Aldrich, Pestanal®, CAS # 52645-53-1, >95.0% sum of cis+trans 97.8%) or malathion (Sigma Aldrich, Pestanal®, CAS # 121-75-5) and larvae were selected with temephos (CHEM SERVICE INC., CAS # 3383-96-8) (see “Mosquito selection regime”). Groups of 25 L3 larvae were exposed to different concentrations of temephos in 200 mL of water to confirm the current LC₅₀. The LC₅₀ identified for each colony-insecticide combination was used in subsequent selection procedures described below. However, we could not achieve an LC₅₀ in the CAY/permethrin combination that was relevant to concentrations that mosquitoes would typically encounter. We therefore decided to expose CAY to a high concentration (3%) that is similar to doses received in the wild.

Mosquito selection regime

Cayman colony. We exposed 2–5 day old adult female mosquitoes in one strain, CAY-P, to 3% permethrin every generation according to the standard WHO protocol (WHO, 2013). After a one-hour exposure and 24 hours recovery time, mortality was recorded and the surviving adults were allowed to mate and bloodfeed to create the next generation. We maintained another strain without insecticide exposure for ten generations (CAY-U). After five generations without insecticide exposure (G1–G6), we used molecular tools to manually split CAY-U (see “Manual selection of *kdr* alleles”) in generations 6 and 8 due to the high frequency of *kdr* alleles (I1016 and C1534). Four unselected strains were established: i) homozygous susceptible individuals (CAY-SS: V1016 and F1534), ii) homozygous resistant individuals (CAY-RR: I1016 and C1534), and iii) two heterozygote strains created by crossing resistant females with susceptible males (CAY-RS) or susceptible females with resistant males (CAY-SR) (see Figure 1).

Recife colony. We selected larvae with temephos every three generations to create REC-R (Melo-Santos *et al.*, 2010). Groups of L3 larvae were exposed to 0.5 mg/L temephos in plastic trays for 24 hours. At 24 hours, the mortality was recorded and surviving larvae were transferred to fresh water, provided with food, and allowed to pupate and emerge as usual. Adult females were allowed to mate and bloodfeed to create the next generation. Two additional strains were established by exposing 2–5 day old adult female mosquitoes to malathion (REC-M: 1% WHO papers for 6 generations and 1.5% WHO papers for 3 generations) or permethrin (REC-P: 0.4% WHO papers for 6 generations and 0.75% WHO papers for 3 generations) every generation according to the standard WHO protocol (WHO, 2013) (see Figure 1). The exposure concentrations were determined by the initial discriminating dose. Survivors were allowed to mate and bloodfeed to create the next generation. The concentration was increased at G7 following a decrease in mortality to 25% after exposure in G6 (see Extended data: Table S1 (Reimer, 2020)).

Evaluating phenotypic resistance

We performed WHO tube assays (WHO, 2013) pre-selection and every three generations (G3, G6 and G9) during the selection regime for permethrin or malathion. Assays were performed with standard WHO papers at diagnostic concentration (0.75% for permethrin, 5% for malathion), ordered from the Vector Control Research Unit, School of Biological Sciences, Universiti Sains Malaysia. Four exposure tubes and one negative control tube were filled with up to 25 mosquitoes each. Mosquitoes were exposed for one hour, returned to the holding tubes and provided with 10% sucrose solution. Mortality was recorded after 24 hours.

Detection of *kdr* alleles

For DNA analysis, 50 female mosquitoes were analysed every third generation of selection. Mosquitoes were killed at -20°C and stored on silica gel. We extracted DNA from individual mosquitoes using the Livak method (Livak, 1984). TaqMan® SNP Genotyping Assays for *Vgsc*-1016 and *Vgsc*-1534 alleles in *Aedes* (Extended data: Table S2 (Reimer, 2020)) were used to screen for *kdr* in the Cayman strains. The lack of resistant *kdr* alleles at *Vgsc*-1016 and *Vgsc*-1534 was confirmed in a small subsample of Recife pre-selection (G17). TaqMan reactions were performed in 10 µl volumes containing 1X TaqMan® Gene Expression Master Mix (Thermo Fisher Scientific, MA, USA), 800 nM of each primer, and 200 nM of each probe on an Mx3005P qPCR thermal cycler (Agilent Technologies, CA, USA) with initial denaturation of 10 min at 95°C followed by 40 cycles of 15 s at 92°C and 1 min at 60°C.

Detoxification gene expression

For RNA analysis, total RNA was extracted from pools of five female mosquitoes using Quick-RNA™ Miniprep (Zymo Research, CA, USA) and the purity and quantity of RNA were individually determined using a Nanodrop spectrophotometer (Nanodrop Technologies, DE, USA). SuperScript® III First-Strand Synthesis System performed cDNA synthesis from total RNA using oligo-dT20 primer (Thermo Fisher Scientific). Four genes associated with insecticide resistance were selected to screen expression profiles in larvae (CCae3A, CYP6N12, CYP6F3 and CYP9M9: all associated with temephos resistance) and adults (CCae3A and CYP6N12, associated with temephos resistance; and CYP6BB2 and CYP9J28a, associated with pyrethroid resistance; Moyes *et al.*, 2017). cDNA was diluted ten-fold and qPCR reactions were performed in 20 µl volumes containing 2 µl of cDNA, 1x PowerUp™ SYBR® Green Master Mix (Thermo Fisher Scientific) and 300 nM of each primer on an Mx3005P qPCR thermal cycler (Agilent Technologies) with initial denaturation of 10 min at 95°C followed by 40 cycles of 15 s at 92°C and 30 s at specific TA (Extended data: Table S3 (Reimer, 2020)). The specificity of the primers was verified by melting curve analyses.

Relative fold gene expression was calculated using the comparative CT method (2^{-ΔΔCT} method), taking into account PCR efficiency (Pfaffl, 2001). The genes coding for the 60S ribosomal protein L8 (RPL8) and the 40S ribosomal protein S7 (RPS7) were defined as reference genes. Between three and five biological replicates were performed for each strain and REC as baseline, respectively. All samples were run in duplicate.

Results were expressed as mean transcription ratio in each strain and life stage $\pm$ SD relative to the mean transcription ratio of the specific life-stage of REC. Mann–Whitney U tests from the R package “stats” (R version 4.01; Copyright (C) 2020 The R Foundation for Statistical Computing) were used to compare transcription ratios between the selected strains and REC. The sequential Holm–Bonferroni procedure (Holm, 1979) was used to adjust α to account for multiple comparisons.

Manual selection of *kdr* alleles

CAY-U was maintained without insecticide pressure for five generations (G1–G6) before starting the selection process to establish homozygous strains for *Vgsc*-1016 and *Vgsc*-1534, which over the course of the process showed complete linkage in all individuals. These strains were established by separating pupae by sex and removing a leg from each adult for genotyping, as described above, before returning the adults to a cage for mating. The process was divided into two steps: 1) in generation G6, all the isoleucine homozygote mosquitoes for *Vgsc*-1016 were removed, and only heterozygotes and valine homozygotes were allowed to mate; 2) in generation G8, two homozygous strains were established for *Vgsc*-1534 (CAY-SS: Phe/Phe; CAY-RR: Cys/Cys), and all the heterozygotes were removed. Both CAY-SS and CAY-RR were screened by TaqMan assays for *Vgsc*-1016 and *Vgsc*-1534 in generation G9 to confirm the genotype of each strain. An extra selection was repeated in G9 for CAY-SS to remove the few individuals with *kdr* alleles.

Determination of resistance ratio based on LC_{50} and LC_{95}
Standardised larval trays were prepared with 200 L1 larvae and provided a yeast tablet every other day until pupation. Adult two to five day old female mosquitoes were exposed to insecticide papers of a range of concentrations of malathion and permethrin, as described previously. Mortality was calculated 24 hours after exposure, and at least three replicates of each assay were performed. For larval bioassays, stock concentrations of temephos were prepared at 0.05 – 1 mg/ml. 1 ml of stock was added to 750 ml of water mixed in a 1:1 ratio from distilled water and larval water from the trays. The water was mixed and aliquoted into five pupae pots of 150 ml each and groups of 25 L3 larvae in 50 ml water were added to each. This process was repeated for each concentration in the assay. Larval mortality was recorded at 24 hours and Abbot’s formula (Abbott, 1925) was used to adjust mortality when necessary. The data from bioassays was organized by concentration in each biological replicate. These data were used to create General linear models (G.L.M.) with “logit” models for each strain using “glm” function in statistical software R-2.15.2. Lethal concentration and their 95% confidence intervals at 95% were calculated using the R-package “MASS”. Resistance ratios were calculated based on comparison of LC_{50} and LC_{95} from each strain against the reference colony New Orleans, which is fully susceptible to all three insecticides. Confidence intervals for ratios were calculated using the method MOVER-R (Newcombe, 2016) presented in the R-package “pairwiseCI”.

Mosquito colony maintenance

All mosquitoes were maintained in the insectary of the Liverpool School of Tropical Medicine under controlled temperature

($26 \pm 2^\circ\text{C}$), relative humidity ($75 \pm 20\%$), and photoperiod (12:12 L:D). Adult mosquitoes were housed in BugDorm cages (MegaView Science Co, Ltd, Taiwan) and provided with constant access to 10% sucrose solution on a cotton pad, which was changed weekly. Eggs were obtained by feeding mated adult females on blood using a Hemotek feeder (Hemotek Ltd, Blackburn, UK). Due to issues with our supplier, the blood source was changed from human to horse at the beginning of the selection procedures. However, issues with mosquito egg-laying performance forced us to switch back to human blood mixed from separate bags of red blood cells and plasma in a 50:50 ratio from supplier overstock. Larvae were reared in plastic trays and fed Brewer’s Yeast tablets (Nature’s Aid®).

Results

Cayman

In the unexposed strain CAY-U, we observed a slight increase in mortality over time for the standard WHO bioassay (0.75% permethrin) (Figure 2). In the permethrin selected CAY-P strain, mortality decreased to nearly 0% compared to the baseline of 4.6% mortality after only three generations of insecticide exposure at 3% permethrin. The *kdr* resistant alleles were still present in CAY-U at a high frequency (Table 1).

***Kdr* allele frequency.** Both *kdr* alleles (V1016I and F1534C) showed complete linkage in all 230 individuals. The allele frequency for both *kdr* alleles in the Cayman colony was 93% at baseline with a high frequency of resistant homozygotes (86%). Selection with 3% permethrin (CAY-P) led to *kdr* fixation within five generations. In CAY-U, *kdr* allele frequency was 68% split between a similar proportion of heterozygotes (46.4%) and resistant homozygotes (44.6%) (Table 1).

Only 1.27% of CAY-RR died in standard WHO tube bioassays with permethrin whereas CAY-SS had a mortality rate of 82.9%.

Resistance ratio to permethrin. CAY-P displayed a resistance ratio over 29x that of CAY-SS when compared to the reference colony New Orleans (Table 2). CAY-P and the unexposed CAY-RR had similar resistance ratios of over 190x resistance compared to New Orleans. Resistance ratios in the heterozygote strains CAY-RS and CAY-SR were similar to each other, around 2x higher than CAY-SS, and 13x lower than CAY-RR.

Recife

In the Recife colony, for both the malathion and permethrin-selected strains, a sharp drop in mortality to the WHO tube assay was observed, followed by a recovery then a shallower drop, rather than a gradual increase in resistance over selected generations (Figure 3). However, the mortality of REC-P for standard WHO bioassay (0.75% permethrin) on later generations (generations 6 and 9) is around 75%, while REC-M present mortality values for standard WHO bioassay (5% malathion) above 90% on later generations (Figure 3).

Differential expression of detoxification genes. The detoxification genes with differential expression among REC strains varied between larvae and adults (Table 3 and Table 4). In REC-R, we

Figure 2. Mortality of Cayman strains against WHO standard permethrin 0.75%. The mortality profile of the CAYMAN strain continuously exposed to permethrin (CAY-P) is compared to the unexposed strain after exposure to 0.75% permethrin in the WHO bioassay. By generation 21, CAY-U had been split into *kdr* homozygous individuals (CAY-RR) and homozygous susceptible (CAY-SS) and the bioassay was repeated in generation 22. Bars represent 95% confidence intervals.

Table 1. Frequency of *Kdr* alleles V1016I and F1534C in the Cayman strains across generations.

Strain	G	Details	V1016I				F1534C			
			N	Val/Val	Val/Ile	Ile/Ile	N	Phe/Phe	Phe/Cys	Cys/Cys
CAY	G1	Baseline	43	-	0.140	0.860	42	-	0.143	0.857
CAY-U	G3	unexposed	48	-	0.208	0.792	48	-	0.208	0.792
CAY-P	G3	PERM 3%	47	-	0.085	0.915	48	-	0.083	0.917
CAY-U	G6	unexposed	56	0.089	0.464	0.446	52	0.077	0.462	0.462
CAY-P	G6	PERM 3%	49	-	-	1.000	40	-	-	1.000
CAY-U	G6	1 st separation	351	0.046	0.313	0.641	NA	NA	NA	NA
CAY-U	G8	2 nd separation	NA	NA	NA	NA	416	0.269	0.510	0.221
CAY-RR	G10	screen	23	-	-	1.000	23	-	-	1.000
CAY-SS	G10	screen	23	0.978	-	0.022	23	0.978	-	0.022
CAY-SS	G10	cleaning	189	1.000	-	-	138	0.957	-	0.043

CAY-U: strain without insecticide exposure; CAY-P: strain selected with 3% permethrin; CAY-SS: homozygous for susceptible alleles without insecticide exposure; CAY-RR: homozygous for resistant alleles without insecticide exposure; CAY-RS: heterozygous by the cross between CAY-RR females and CAY-SS males; CAY-SR: heterozygous by the cross between CAY-SS females and CAY-RR males

Table 2. Lethal concentrations and resistance ratios of Cayman strains for permethrin.

Insecticide	Strain	LC ₅₀	RR ₅₀	LC ₉₅	RR ₉₅
Permethrin	New Orleans	0.076 (0.066–0.087)	N/A	0.14 (0.12–0.16)	N/A
	CAY-SS	0.50 (0.42–0.58)	6.58 (5.31–8.09)	0.98 (0.81–1.16)	7.00 (5.43–9.00)
	CAY-RR	15.53 (14.87–16.20)	203.31 (177.49–236.89)	25.75 (24.33–27.17)	183.18 (154.80–222.63)
	CAY-RS	1.16 (0.98–1.33)	15.16 (12.32–18.58)	2.31 (1.84–2.78)	16.43 (12.46–21.39)
	CAY-SR	1.07 (0.95–1.20)	14.01 (11.73–16.84)	1.84 (1.49–2.81)	13.06 (10.05–16.85)
	CAY-P	14.87 (14.16–15.57)	194.58 (169.59–226.99)	24.22 (22.75–25.68)	172.29 (145.31–209.70)

LC₅₀: Lethal concentration for 50% mortality; RR₅₀: resistance ratio for LC₅₀; LC₉₅: Lethal concentration for 95% mortality; RR₉₅: resistance ratio for LC₉₅; brackets: confidant intervals at 95%. CAY-SS: homozygous for susceptible alleles without insecticide exposure; CAY-RR: homozygous for resistant alleles without insecticide exposure; CAY-RS: heterozygous by the cross between CAY-RR females and CAY-SS males; CAY-SR: heterozygous by the cross between CAY-SS females and CAY-RR males; CAY-P: strain selected with 3% permethrin.

observed no changes in gene expression from baseline in larval stages (Table 3), but expression of CYP9J28 and CYP6BB2 in adults increased significantly. Some genes were significantly downregulated at this stage in other Recife strains: i) REC-M (CCae3A, CYP6F3, and CYP9M9; MW: $p < 0.000666$); ii) REC-U (CCae3A; MW: $p < 0.002664$); iii) REC-P (CYP6F3; MW: $p < 0.002664$) (Table 3). In adults of the REC-R strain, two genes were significantly upregulated in generation G45 (CYP9J28 and CYP6BB2) (MW: $p < 0.004329$). A significant upregulation in these genes was observed in REC-U (CYP9J28; MW: $p < 0.002498$) and REC-M (CYP9J28 and CYP6BB2; MW: $p < 0.002165$). Moreover, the CCae3A gene was upregulated in REC-M (MW: $p < 0.000250$) in adults (Table 4).

Resistance ratio to permethrin, malathion and temephos.

Nearly all strains were more resistant to all insecticides than New Orleans. REC-P was as much as 4x more resistant to permethrin than REC-U, and REC-R (Table 5). REC-R and REC-M were slightly more resistant to malathion (~1.4x) than REC-U. REC-R, REC-M and REC-P were more resistant to temephos (>2.5x) than REC-U.

Discussion

Inference of insecticide resistance in adults of *Ae. aegypti* is rarely performed by quantitative methods. A recent review (Moyes *et al.*, 2017) highlights the lack of literature that calculate resistance ratios based on dose-response, lethal concentration or lethal time (see S2 file in (Moyes *et al.*, 2017)). This practice limits comparative analysis of our results with other resistance studies. Resistance ratios for permethrin in CAY with homozygous resistance alleles (II/CC; CAY-RR and CAY-P) were of a similar magnitude to homozygous resistant *Ae. aegypti* from

Cayman Islands populations (Harris *et al.*, 2010). In other field populations, a significant positive correlation between the frequency of IICC individuals and resistance ratio for permethrin was also observed, which indicates a higher resistance for these double homozygotes (Estep *et al.*, 2018). The lack of variation in permethrin resistance ratio between IICC strains regardless of selection pressure indicates that this allele is primarily responsible for the phenotype observed. Moreover, the other CAY strains presented lower resistance ratios to permethrin. The susceptible double homozygous strain (VV/FF; CAY-SS) had a resistance ratio similar to susceptible homozygous field populations (RR: 0.8 – 7.0) in Asia (Bregues *et al.*, 2003) and lower than REC-P (the selected strain for permethrin in REC). This intermediate level of resistance in heterozygotes is consistent with the recessive nature of *kdr* alleles in mosquitoes and other dipterans (Gomes *et al.*, 2017; Huang *et al.*, 2004; Saavedra-Rodriguez *et al.*, 2007). However, the combination of multiple heterozygote *kdr* alleles in *Ae. aegypti* can present a stronger resistance phenotype in the future, as is observed in Thailand where triple heterozygotes (S/P989 + V/G1016 + F/C1534) had a higher resistance ratio to deltamethrin than *kdr* homozygotes at F1534C (Plernsub *et al.*, 2016).

Changes in temephos resistance in our REC strains differed from previous studies in two primary ways. First, the resistance level in our REC-R temephos selected strain was lower than the resistance level reported in previous studies where REC was put under similar selection pressure (Diniz *et al.*, 2015; Melo-Santos *et al.*, 2010). This is likely because we used the New Orleans colony as the denominator in calculating the resistance ratio, while the Rockefeller colony was used in other studies. Rockefeller has a lower LC₅₀ on average than New

Figure 3. Mortality of Recife strains against WHO standard malathion 5.0% (top) and permethrin 0.75% (below). Bars represent 95% confidence intervals.

Table 3. Mean fold change in gene expression for larvae, Recife colony.

Strain	G	CCae3A		CYP6F3		CYP6N12		CYP9M9	
		MF	sd	MF	sd	MF	sd	MF	sd
REC-Baseline	G1	1.06	0.39	1.13	0.54	1.12	0.52	1.07	0.43
REC-U	G9	0.34	0.28	0.65	0.34	0.66	0.56	0.78	0.47
REC-R	G9	0.73	0.23	0.92	0.38	1.81	0.7	1.70	0.82
REC-M	G9	0.38	0.12	0.41	0.07	0.59	0.10	0.20	0.05
REC-P	G9	1.77	0.39	0.41	0.13	0.95	0.38	0.57	0.19

G: Generation; MF: mean fold; sd: standard deviation. REC-U: strain without insecticide exposure; REC-R: strain with temephos exposure; REC-M: strain selected for malathion; REC-P: strain selected for permethrin. Bold: Expression values significantly different from Baseline (Holm-Bonferroni method)

Table 4. Mean fold change in gene expression for adults, Recife colony.

Strain	G	CCae3A		CYP9J28		CYP6N12		CYP6BB2	
		MF	sd	MF	sd	MF	sd	MF	sd
REC -Baseline	G1	1.35	0.79	1.04	0.29	1.05	0.31	1.39	0.82
REC-U	G9	1.75	0.43	2.63	0.47	0.63	0.25	1.41	0.37
REC- R	G9	1.7	0.52	2.56	0.31	0.74	0.20	3.34	0.75
REC-M	G9	2.89	0.37	3.37	2.11	1.73	0.34	3.99	0.90
REC-P	G9	1.56	1.00	0.72	0.38	0.74	0.22	1.95	0.65

G: Generation; MF: mean fold; sd: standard deviation. REC-U: strain without insecticide exposure; REC-R: strain with temephos exposure; REC-M: strain selected for malathion; REC-P: strain selected for permethrin. Bold: Expression values significantly different from Baseline (Holm-Bonferroni method).

Orleans (see S1 file in (Moyes *et al.*, 2017)). Second, in this study, temephos resistance in the REC-U unselected strain was not completely reversed while previous studies have documented reversal in a similar number of generations. This may be because our starting material had been under temephos selection pressure for longer prior to starting the experiments.

In the Recife strains, the response to the adulticides malathion and permethrin was different to the response to the larvicide temephos. REC P and REC M showed a similar LC50 to temephos as the REC-R colony and a slightly lower LC95. The gene expression within REC-R showed no significant variation compared to the baseline colony. However, REC-P, REC-M and REC-U showed downregulation of detoxification genes compared to the baseline, consistent with the lower tolerance to temephos displayed in the LC95s.

In contrast to larvicide exposure, resistance ratios for REC-P and REC-M showed a different pattern. Substantial differences between REC-P compared to either REC-R or REC-U suggest that exposure to permethrin increased the tolerance of this insecticide in the Recife colony. REC-M and REC-R present

similar resistance ratios to Malathion, while the unexposed strain REC-U presents a lower resistance ratio than REC-M/REC-R. Malathion exposure over more generations will be required to increase the divergence between REC-R and REC-M phenotypes. We have experienced three main limitations in the selection of these strains: i) inconsistent blood sources, ii) time required to reverse the resistance mechanism (Diniz *et al.*, 2015), iii) potential diversity loss associated with bottlenecks and/or high mortality due to aggressive artificial selection. We experienced difficulties in maintaining *Ae. aegypti* after a few generations using horse blood. Unfortunately, this was the only blood source available for our lab after the source of human blood was interrupted. Moreover, REC-M exhibited a drastic increase in mortality (over 90%) when we adjusted the malathion exposure concentration to 1.5% at generation 43. Mortality levels in malathion selection remained higher than 50% after this adjustment, and we had difficulties maintaining this strain post-generation 45. Future malathion selection will require a longer build-up to create a more viable resistant strain. Our method to calculate resistance ratios using WHO tubes is not able to define the individual dose received per mosquito, since mosquito weight and individual activity against the treated surface could not be measured. Topical assays, which don't

Table 5. Lethal concentrations and resistance ratios of Recife strains for three insecticides (i.e. permethrin, malathion and temephos).

Insecticide	Strain	LC ₅₀	RR ₅₀	LC ₉₅	RR ₉₅
Permethrin	New Orleans	0.076 (0.066–0.087)	N/A	0.14 (0.12–0.15)	N/A
	REC-U	0.18 (0.12–0.25)	2.41 (1.58–3.34)	0.36 (0.22–0.49)	2.53 (1.53–3.68)
	REC-R	0.23 (0.17–0.29)	3.01 (2.16–3.97)	0.44 (0.32–0.56)	3.11 (2.17–4.24)
	REC-P	0.91 (0.84–0.98)	11.09 (9.52–13.09)	1.73 (1.58–1.88)	12.32 (10.26–15.23)
Malathion	New Orleans	0.51 (0.44–0.57)	N/A	0.98 (0.85–1.11)	N/A
	REC-U	0.75 (0.68–0.82)	1.48 (1.27–1.74)	1.57 (1.44–1.71)	1.60 (1.37–1.89)
	REC-R	1.18 (1.08–1.27)	2.32 (2.02–2.70)	2.22 (2.01–2.43)	2.25 (1.92–2.67)
	REC-M	1.23 (1.12–1.35)	2.43 (2.09–2.85)	2.65 (2.37–2.93)	2.70 (2.28–3.21)
Temephos	New Orleans	0.012 (0.012–0.013)	N/A	0.024 (0.022–0.026)	N/A
	REC-U	0.15 (0.15–0.15)	12.27 (11.52–13.12)	0.26 (0.25–0.27)	11.03 (10.14–12.04)
	REC-R	0.40 (0.39–0.42)	32.74 (30.51–35.20)	0.90 (0.86–0.94)	37.65 (34.56–41.14)
	REC-M	0.42 (0.40–0.44)	34.20 (31.81–36.83)	0.76 (0.73–0.80)	31.92 (29.30–34.88)
	REC-P	0.40 (0.38–0.41)	32.13 (29.94–34.55)	0.70 (0.66–0.73)	29.12 (26.76–31.79)

LC₅₀: Lethal concentration for 50% mortality; RR₅₀: resistance ratio for LC₅₀; LC₉₅: Lethal concentration for 95% mortality; RR₉₅: resistance ratio for LC₉₅; brackets: confident intervals at 95%. REC-U: strain without insecticide exposure; REC-R: strain with temephos exposure; REC-M: strain selected for malathion; REC-P: strain selected for permethrin.

mimic natural exposure routes, do allow for controlled application and would further reduce the individual variation observed within each strain.

Conclusions

We generated strains of *Ae. aegypti* which differ in phenotypic resistance to permethrin, malathion and temephos. The selected CAY and REC strains will allow for further research on the effects of target-site and metabolic resistance, respectively, on the life-history traits, behaviour and vector competence of this important arbovirus vector. The strains can also be used to compare the efficacy of novel insecticide formulations in strains

with similar genetic backgrounds and different mechanisms of resistance.

Data availability

Underlying data

Open Science Framework: WT Seed project, <https://doi.org/10.17605/OSF.IO/8DQ9A> (Reimer, 2020). Project registered on 10th July 2020 (osf.io/f49jg).

This project contains the following underlying data:

- Raw values for insecticide selection mortality per generation

- Raw values for mortality in WHO bioassay
- CT values for detoxification genes in larvae
- CT values for detoxification genes in adults
- Raw values for mortality to a range of insecticide concentrations, used to calculate LC50, LC95 and resistance ratios

Extended data

Open Science Framework: WT Seed project, <https://doi.org/10.17605/OSF.IO/8DQ9A> (Reimer, 2020). Project registered on 10th July 2020 (osf.io/f49jg).

This project contains the following extended data:

- Table S1. Insecticide selection mortality (%) for Cayman and Recife strains against standard WHO tube concentrations.
- Table S2. Primers and probes for TaqMan® SNP Genotyping Assays for Vgsc-1016 and Vgsc-1534 alleles.
- Table S3. Primers for qPCR screening of genes associated with insecticide resistance and controls.

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

References

- Abbott WS: **A method of computing the effectiveness of an insecticide.** *J Econ Entomol.* The Oxford University Press. 1925; **8**(2): 265–267.
[Publisher Full Text](#)
- Brenques C, Hawkes NJ, Chandre F, *et al.*: **Pyrethroid and DDT cross-resistance in *Aedes aegypti* is correlated with novel mutations in the voltage-gated sodium channel gene.** *Med Vet Entomol.* Wiley/Blackwell (10.1111). 2003; **17**(1): 87–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Brito LP, Linss JG, Lima-Camara TN, *et al.*: **Assessing the Effects of *Aedes aegypti* kdr Mutations on Pyrethroid Resistance and Its Fitness Cost.** *PLoS One.* Public Library of Science. 2013; **8**(4): e60878.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Burt FJ, Chen W, Miner JJ, *et al.*: **Chikungunya virus: an update on the biology and pathogenesis of this emerging pathogen.** *Lancet Infect Dis.* Elsevier. 2017; **17**(4): e107–e117.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Corbel V, Fonseca DM, Weetman D, *et al.*: **International workshop on insecticide resistance in vectors of arboviruses, December 2016, Rio de Janeiro, Brazil.** *Parasit Vectors.* BioMed Central. 2017; **10**(1): 278.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Davies TE, O'Reilly AO, Field LM, *et al.*: **Knockdown resistance to DDT and pyrethroids: from target-site mutations to molecular modelling.** *Pest Manag Sci.* John Wiley & Sons, Ltd. 2008; **64**(11): 1126–1130.
[PubMed Abstract](#) | [Publisher Full Text](#)
- de Lourdes Macoris M, Martins AJ, Andrighetti MT, *et al.*: **Pyrethroid resistance persists after ten years without usage against *Aedes aegypti* in governmental campaigns: Lessons from São Paulo State, Brazil.** *PLoS Negl Trop Dis.* 2018; **12**(3): e0006390. Public Library of Science.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Diniz DF, de Melo-Santos MA, Santos EM, *et al.*: **Fitness cost in field and laboratory *Aedes aegypti* populations associated with resistance to the insecticide temephos.** *Parasit Vectors.* BioMed Central. 2015; **8**: 662.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Douam F, Ploss A: **Yellow Fever Virus: Knowledge Gaps Impeding the Fight Against an Old Foe.** *Trends Microbiol.* Elsevier. 2018; **26**(11): 913–928.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Estep AS, Sanscrainte ND, Waits CM, *et al.*: **Quantification of permethrin resistance and kdr alleles in Florida strains of *Aedes aegypti* (L.) and *Aedes albopictus* (Skuse).** *PLoS Negl Trop Dis.* Public Library of Science. 2018; **12**(10): e0006544.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Feyereisen R, Dermauw W, Van Leeuwen T: **Genotype to phenotype, the molecular and physiological dimensions of resistance in arthropods.** *Pestic Biochem Physiol.* Academic Press. 2015; **121**: 61–77.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ffrench-Constant RH: **The molecular genetics of insecticide resistance.** *Genetics.* 2013; **194**(4): 807–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- George L, Lenhart A, Toledo J, *et al.*: **Community-Effectiveness of Temephos for Dengue Vector Control: A Systematic Literature Review.** *PLoS Negl Trop Dis.* Public Library of Science. 2015; **9**(9): e0004006.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gomes B, Purkait B, Deb RM, *et al.*: **Knockdown resistance mutations predict DDT resistance and pyrethroid tolerance in the visceral leishmaniasis vector *Phlebotomus argentipes*.** *PLoS Negl Trop Dis.* 2017; **11**(4): e0005504.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Guzman MG, Harris E: **Dengue.** *Lancet.* Elsevier. 2015; **385**(9966): 453–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Harris AF, Rajatileka S, Ranson H: **Pyrethroid resistance in *Aedes aegypti* from Grand Cayman.** *Am J Trop Med Hyg.* The American Society of Tropical Medicine and Hygiene. 2010; **83**(2): 277–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Holm S: **A simple sequentially rejective multiple test procedure.** *Scand J Stat.* 1979; **6**(2): 65–70.
[Reference Source](#)
- Horstick O, Runge-Ranzinger S, Nathan MB, *et al.*: **Dengue vector-control services: how do they work? A systematic literature review and country case studies.** *Trans R Soc Trop Med Hyg.* 2010; **104**(6): 379–386. Oxford University Press.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Huang J, Kristensen M, Qiao CL, *et al.*: **Frequency of kdr gene in house fly field populations: correlation of pyrethroid resistance and kdr frequency.** *J Econ Entomol.* 2004; **97**(3): 1036–41. The Oxford University Press.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ingham VA, Wagstaff S, Ranson H: **Transcriptomic meta-signatures identified in *Anopheles gambiae* populations reveal previously undetected insecticide resistance mechanisms.** *Nat Commun.* 2018; **9**(1): 5282. Nature Publishing Group.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jankowska M, Rogalska J, Wyszowska J, *et al.*: **Molecular Targets for Components of Essential Oils in the Insect Nervous System—A Review.** *Molecules.* 2017; **23**(1): 34. Multidisciplinary Digital Publishing Institute.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kindhauser MK, Allen T, Frank V, *et al.*: **Zika: the origin and spread of a mosquito-borne virus.** *Bull World Health Organ.* 2016; **94**(9): 675–686C. World Health Organization.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Li X, Schuler MA, Berenbaum MR: **Molecular Mechanisms of Metabolic Resistance to Synthetic and Natural Xenobiotics.** *Annu Rev Entomol.* 2007; **52**: 231–253. Annual Reviews.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Livak KJ: **Organization and mapping of a sequence on the *Drosophila melanogaster* X and Y chromosomes that is transcribed during spermatogenesis.** *Genetics.* 1984; **107**(4): 611–34. Genetics Society of America.
[PubMed Abstract](#) | [Free Full Text](#)
- Melo-Santos MA, Varjal-Melo JJ, Araújo AP, *et al.*: **Resistance to the organophosphate temephos: Mechanisms, evolution and reversion in an**

***Aedes aegypti* laboratory strain from Brazil.** *Acta Trop.* 2010; **113**(2): 180–189. Elsevier.

[PubMed Abstract](#) | [Publisher Full Text](#)

Monath TP, Vasconcelos PF: **Yellow fever.** *J Clin Virol.* 2015; **64**: 160–173. Elsevier.

[PubMed Abstract](#) | [Publisher Full Text](#)

Moyes CL, Vontas J, Martins AJ, *et al.*: **Contemporary status of insecticide resistance in the major *Aedes* vectors of arboviruses infecting humans.** *PLoS Negl Trop Dis.* 2017; **11**(7): e0005625. Public Library of Science.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Newcombe RG: **MOVER-R confidence intervals for ratios and products of two independently estimated quantities.** *Stat Methods Med Res.* 2016; **25**(5): 1774–1778.

[PubMed Abstract](#) | [Publisher Full Text](#)

Pfaffl MW: **A new mathematical model for relative quantification in real-time RT-PCR.** *Nucleic Acids Res.* 2001; **29**(9): e45. Oxford University Press.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Plernsub S, Saingamsook J, Yanola J, *et al.*: **Additive effect of knockdown resistance mutations, S989P, V1016G and F1534C, in a heterozygous genotype conferring pyrethroid resistance in *Aedes aegypti* in Thailand.** *Parasit Vectors.* 2016; **9**(1): 417. BioMed Central.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Pollett S, Melendrez MC, Maljkovic Berry I, *et al.*: **Understanding dengue virus evolution to support epidemic surveillance and counter-measure development.** *Infect Genet Evol.* 2018; **62**: 279–295. Elsevier.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Poupardin R, Srisukontarat, W, Yunta, C, *et al.*: **Identification of Carboxylesterase Genes Implicated in Temephos Resistance in the Dengue Vector *Aedes aegypti*.** *PLoS Negl Trop Dis.* 2014; **8**: e2743. Public Library of Science.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Ranson H, Burhani J, Lumjuan N, *et al.*: **Insecticide resistance in dengue vectors.** *scielotropika.* 2010.

[Reference Source](#)

Reimer LJ: **WT Seed project.** 2020.

<http://www.doi.org/10.17605/OSF.IO/8DQ9A>

Rinkevich FD, Du Y, Dong K: **Diversity and Convergence of Sodium Channel Mutations Involved in Resistance to Pyrethroids.** *Pestic Biochem Physiol.* NIH Public Access, 2013; **106**: 93–100.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Saavedra-Rodriguez K, Urdaneta-Marquez L, Rajatileka S, *et al.*: **A mutation in the voltage-gated sodium channel gene associated with pyrethroid resistance in Latin American *Aedes aegypti*.** *Insect Mol Biol.* Blackwell Publishing Ltd, 2007; **16**(6): 785–798.

[PubMed Abstract](#) | [Publisher Full Text](#)

Silva JV, Lopes TR, Oliveira-Filho EF, *et al.*: **Current status, challenges and perspectives in the development of vaccines against yellow fever, dengue, Zika and chikungunya viruses.** *Acta Trop.* 2018; **182**: 257–263.

[PubMed Abstract](#) | [Publisher Full Text](#)

Stevenson BJ, Pignatelli P, Nikou D, *et al.*: **Pinpointing P450s Associated with Pyrethroid Metabolism in the Dengue Vector, *Aedes aegypti*: Developing New Tools to Combat Insecticide Resistance.** *PLoS Negl Trop Dis.* 2012; **6**(3): e1595. Public Library of Science.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Strode C, de Melo-Santos M, Magalhães T, *et al.*: **Expression Profile of Genes during Resistance Reversal in a Temephos Selected Strain of the Dengue Vector, *Aedes aegypti*.** *PLoS One.* 2012; **7**(8): e39439.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Vontas J, Kioulos E, Pavlidi N, *et al.*: **Insecticide resistance in the major dengue vectors *Aedes albopictus* and *Aedes aegypti*.** *Pestic Biochem Physiol.* 2012; **104**: 126–131.

[Publisher Full Text](#)

Weetman D, Djogbenou LS, Lucas E: **Copy number variation (CNV) and insecticide resistance in mosquitoes: evolving knowledge or an evolving problem?** *Curr Opin Insect Sci.* 2018; **27**: 82–88.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

World Health Organization (WHO): **Test procedures for insecticide resistance monitoring in malaria vector mosquitoes.** 2013; Geneva.

[Reference Source](#)

Yunta C, Grisales N, Nász S, *et al.*: **Pyriproxyfen is metabolized by P450s associated with pyrethroid resistance in *An. gambiae*.** *Insect Biochem Mol Biol.* 2016; **78**: 50–57.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 18 January 2021

<https://doi.org/10.21956/wellcomeopenres.18162.r42088>

© 2021 Han Q. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Qian Han

Laboratory of Tropical Veterinary Medicine and Vector Biology, School of Life and Pharmaceutical Sciences, Hainan University, Haikou, China

The manuscript describes a study to develop multiple *Aedes aegypti* strains with resistance to different insecticides. The study includes two resistant colonies of *Ae. aegypti*, selection of eight strains with different profiles of resistance after exposure to different insecticides, calculation of resistance ratios for each insecticide for the selected and unselected strains, frequency of *kdr* alleles and some gene expression analysis. The manuscript is in an important area of methodology study understanding mechanisms underlying insecticide resistance.

After revision based on the comments from Reviewer 1, the manuscript was well organized and written.

Minors

1. Mosquito name typing. Keywords: *Aedes aegypti* in italic; Introduction, the second paragraph: *Aedes spp.* *Aedes* in italic, spp. in normal font; Methods: *Aedes aegypti* should be written in *Ae. aegypti* in italic.
2. Subscript fonts. Discussion, the third paragraph: "LC50, LC95", here 50 and 95 should be written in subscript. Also correct other similar errors in the sections of Method and Data availability.
3. To be consistent in typing "Cayman": "CAYMAN". "Cayman" or "CAY"? Same to "RECIFE".

Is the rationale for developing the new method (or application) clearly explained?

Yes

Is the description of the method technically sound?

Yes

Are sufficient details provided to allow replication of the method development and its use by others?

Yes

If any results are presented, are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions about the method and its performance adequately supported by the findings presented in the article?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Vector Biology, Biochemistry and Molecular Biology.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 07 January 2021

<https://doi.org/10.21956/wellcomeopenres.18162.r41855>

© 2021 Smith L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Leticia Smith

National Institutes of Health, Rockville, MD, USA

The authors addressed all of my previous comments and I have no new comments on the contents of the article. There were a few typos in the "Determination of resistance ratio..." section in the Methods. Other than that, this article provides a clear description of the methods used to create these insecticide resistant mosquito strains.

Is the rationale for developing the new method (or application) clearly explained?

Yes

Is the description of the method technically sound?

Yes

Are sufficient details provided to allow replication of the method development and its use by others?

Yes

If any results are presented, are all the source data underlying the results available to

ensure full reproducibility?

Yes

Are the conclusions about the method and its performance adequately supported by the findings presented in the article?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Insecticide resistance, mosquito biology, insect physiology.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 17 August 2020

<https://doi.org/10.21956/wellcomeopenres.17519.r39800>

© 2020 Smith L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Leticia Smith

National Institutes of Health, Rockville, MD, USA

This study used two insecticide resistant parental colonies of *Aedes aegypti* to develop multiple strains with resistance to different insecticides. The aim was to create these strains as a tool so that the mechanisms of resistance can be studied without interference from varying genetic backgrounds. Understanding insecticide resistance in mosquitoes is as important now as ever and because insecticides will continue to be used to control disease vectors in the foreseeable future, understanding the mechanisms underlying insecticide resistance is critical for vector control. For this reason, having multiple strains with the same genetic background but different resistance mechanisms and resistance to different insecticides is a great value to the insecticide resistance research community.

Below are my suggestions and specific points related to the questions above that I hope will improve this publication.

The rationale for developing the new method wasn't clearly explained, but perhaps part of my confusion is that there didn't seem to be anything new about the method. The rationale for developing the strains was clearly articulated, but the method of generating strains with the same genetic background through exposing versus not exposing sub-samples of the population to insecticides has been done numerous times already. Also selecting strains based on resistance

genotype has been done before. I may have simply missed it, but perhaps the novelty of this method needs to be elaborated and/or more clearly articulated. As it is now, it seems like the goal was to simply select strains from a parental colony.

The description of the selection method was technically sound, but the evaluation of resistance part of the method could use some improvement. For the detection of *kdr* mutations, is there a reason why V410L was not checked? This is a common mutation in *Ae. aegypti* in the Americas and was only first detected after the initial study that collected the parental strain was published. How were the metabolic genes selected? Are they known to be responsible for resistance in the Recife region where the colony came from? These genes can vary across populations and by insecticide selection, so some clear explanation of how these particular ones were tested is needed. Also, why was this assay done only on the REC strains and not CAY? Some of these genes are also associated with pyrethroid resistance and are likely involved in the resistance observed in the CAY strain. Perhaps insecticide bioassays using CYP, carboxylesterase, or GST synergists would be a better evaluation of metabolic resistance than gene expression? For determining the resistance ratios, why use the WHO assay?

As far as details provided to allow replication of the method development and its use by others, some important information is missing or needs clarification. More information on the parental colonies would be helpful. Although the citations provide information on where the strains originated from, they do not say how many generations the strains have been in the lab prior to this study, or whether they were previously selected for insecticide resistance in the lab or are the original field collected colony. The source and stereochemistry (when applicable) for the insecticides used needs to be provided. The generation numbers used are a bit confusing. It would be helpful if only the generations from this study were used and not a mix between generations selected and what might be total number of generations in the lab. Also, in the mosquito selection regime section, the number of generations don't seem to add up. After 5 generations without exposure, wouldn't that be generation 5? If the authors mean after five additional generations without exposure, wouldn't that be generation 15 (10 + 5)? Figure 1 shows the selection process, but that doesn't seem to match the selection process described in the text. CAY-P was selected at every generation, not every third generation. REC-R was selected as larvae, so not using the WHO bioassay, and REC-M and REC-P seems to have been selected a total of nine generations (the figure shown only three selections). What was the calculation or method used to determine LC_{50} and LC_{95} ? R is a programming language and the software is only an environment to perform computations, this gives no information of how the LC_{50} was calculated. Maybe a link to a GitHub project or profile with the code used? Or a citation for the calculation methodology used? The authors also need to be more specific about how the resistance ratio was calculated. How were they compared? How was significance determined? How was control mortality accounted for? For figures 2 and 3, change "time point" to "generations" and to make it clearer and add a description of what the bars represent (error, st. dev., 95%CI?).

The authors provide a great statement in the beginning for the discussion about the need to perform more dose-response curves in insecticide resistance studies, but sadly this study also didn't exactly provide a dose-response curve. The issue with using a concentration, such as in the WHO assay, is that the exact amount of insecticide the individuals were exposed to remains unknown. Some animals may be more active and therefore be exposed to more by walking around the tube more than others. Also, an individual that is being affected may fall on its side more frequently and in the process get a higher exposure than a healthy animal that is standing

straight. While LC assays are great for certain purposes, they are not optimal for determining resistance ratios and dose-response curves. Would it be possible to do topical assays on the final strains to get a dose response RR instead of a concentration response RR?

In regard to using the New Orleans colony as the denominator in calculating the resistance ratio, a recent study showed that the NO strain has multiple *kdr* mutations¹, just something to keep in mind. In regard to the potential loss of diversity associated with bottlenecks limitations mentioned by the authors in the discussion section, perhaps backcrossing back to the parental colony after each selection could reduce this problem. With enough selections and backcrossing you should retain only the resistance mutations (if they were present in the original population) and any genes that are linked to the resistance alleles (this one is harder to separate out).

On a very minor note, the citation for Harris *et al.* 2010 was listed twice.

References

1. Fan Y, O'Grady P, Yoshimizu M, Ponlawat A, et al.: Evidence for both sequential mutations and recombination in the evolution of *kdr* alleles in *Aedes aegypti*. *PLoS Negl Trop Dis*. **14** (4): e0008154 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the rationale for developing the new method (or application) clearly explained?

No

Is the description of the method technically sound?

Partly

Are sufficient details provided to allow replication of the method development and its use by others?

Partly

If any results are presented, are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions about the method and its performance adequately supported by the findings presented in the article?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Insecticide resistance, mosquito biology, insect physiology.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 02 Dec 2020

Lisa Reimer, Liverpool School of Tropical Medicine, Liverpool, UK

Thank you to Reviewer 1 for the helpful and insightful comments. We have made additions to the manuscript to clarify the scope, re-analysed the data and provided a few additional methodological details and rationale. There were quite a few suggestions for additional research or using different methods than what we have (such as tarsal assays, synergist assays, sequencing). While these are all valid suggestions they were beyond the scope of this methods article. We have hopefully clarified the aims of this paper, which is to describe the methods underlying the selection and creation of the 8 mosquito strains. These strains have been provided to other researchers to address some of the key questions about insecticide resistance and the purpose of this article is to provide the complete background on the process. Our primary outcome was the resistance ratio (compared to a susceptible lab strain). Where we chose a few markers (kdr alleles, metabolic resistance genes, knockdown in WHO assays to select insecticides) the purpose was to track the impact the process of selection was having not to provide a complete characterisation of all possible mechanisms. We have provided a response to the below comments and have uploaded a new version of the manuscript to reflect the changes.

Comment 1

The rationale for developing the new method wasn't clearly explained, but perhaps part of my confusion is that there didn't seem to be anything new about the method. The rationale for developing the strains was clearly articulated, but the method of generating strains with the same genetic background through exposing versus not exposing sub-samples of the population to insecticides has been done numerous times already. Also selecting strains based on resistance genotype has been done before. I may have simply missed it, but perhaps the novelty of this method needs to be elaborated and/or more clearly articulated. As it is now, it seems like the goal was to simply select strains from a parental colony.

This is correct, what we present here is not completely novel methodology but it is a record of the methods used to develop these mosquito resources and it provides an initial characterisation of resistance status. These strains have been shared with other researchers who are currently investigating different aspects of insecticide resistance and we wanted to have a published record of how the resources were established. While the methods are known amongst the scientific community, these selected strains are still a unique resource which allows for the study of multiple resistant phenotypes. We request editorial advice on the most appropriate classification for this manuscript.

Comment 2

The description of the selection method was technically sound, but the evaluation of resistance part of the method could use some improvement. For the detection of kdr mutations, is there a reason why V410L was not checked? This is a common mutation in *Ae. aegypti* in the Americas and was only first detected after the initial study that collected the parental strain was published. How were the metabolic genes selected? Are they known to be responsible for resistance in the Recife region where the colony came from? These genes can vary across populations and by insecticide selection, so some clear explanation of how these particular ones were tested is needed. Also, why was this assay done only on the REC strains and not CAY? Some of these genes are also associated with pyrethroid resistance

and are likely involved in the resistance observed in the CAY strain. Perhaps insecticide bioassays using CYP, carboxylesterase, or GST synergists would be a better evaluation of metabolic resistance than gene expression? For determining the resistance ratios, why use the WHO assay?

CAYMAN has been maintained in laboratory since 2008 by LITE and is screened for two *kdr* alleles (F1534C and V1016I). Moyes et al. (2017) was an important reference in our rationale for the choice to continue screening F1534C and V1016I since they are considered key mutations associated with pyrethroid resistance in *Aedes aegypti* when compared with other *kdr* alleles. We understand that V410L is among the many *kdr* alleles of *Aedes aegypti* in the Americas, and was detected approximately 10 years after our colony was established (though was found in historic samples). We chose to continue the tracking of F1534C and V1016I to perform our selection process. Our results indicate that these two resistance alleles are linked in CAYMAN and it is possible that other *kdr* alleles exist in this strain. The information about alternative resistant alleles which may be contributing further to the resistant phenotype can be explored through sequencing studies which are currently underway by other researchers. We did not screen detox genes in CAYMAN for characterization, since resistant phenotypes suggest similar resistance for both homozygotic resistant strains (CAY-P, CAY-RR).

RECIFE: We selected a battery of detox genes associated with metabolic resistance in *Aedes aegypti* from Americas (based on the literature) to perform an initial screen in our strains. The purpose of the initial screen was to determine whether the selection process was having an impact and to supplement the primary outcome measure which were the resistance ratios. However, as above, further research studies based on RNAseq analysis will be conducted by other researchers. We agree that transcriptomic analyses will be very valuable but they are beyond the scope of this methods article.

We have added some information in the text to clarify genes used and the rationale:
Line 75-81: "Here we present the resistant phenotypes after ten generations of insecticide exposure... Additionally, we present differences in a select number of resistant alleles and metabolic genes which can be used to inform further research."

Line 188-192: "Four genes associated with insecticide resistance were selected to screen expression profiles in larvae (CCae3A, CYP6N12, CYP6F3 and CYP9M9: all associated with temephos resistance in multiple studies) and adults (CCae3A and CYP6N12 associated with temephos resistance; and CYP6BB2 and CYP9J28a associated with pyrethroid resistance in multiple studies, Moyes et al. 2017)."

We also updated the abstract to clarify which alleles were tracked during the selection.

Bioassays with synergist: We agree that studies with synergists and antagonists are interesting for determining metabolic resistance, but this was not a priority during our selection process.

Quantitative approach with WHO assays: we address this concern under comment 4 below

Comment 3

As far as details provided to allow replication of the method development and its use by others, some important information is missing or needs clarification. More information on the parental colonies would be helpful. Although the citations provide information on where the strains originated from, they do not say how many generations the strains have been in the lab prior to this study, or whether they were previously selected for insecticide resistance in the lab or are the original field collected colony. The source and stereochemistry (when applicable) for the insecticides used needs to be provided. The generation numbers used are a bit confusing. It would be helpful if only the generations from this study were used and not a mix between generations selected and what might be total number of generations in the lab. Also, in the mosquito selection regime section, the number of generations don't seem to add up. After 5 generations without exposure, wouldn't that be generation 5? If the authors mean after five additional generations without exposure, wouldn't that be generation 15 (10 + 5)? Figure 1 shows the selection process, but that doesn't seem to match the selection process described in the text. CAY-P was selected at every generation, not every third generation. REC-R was selected as larvae, so not using the WHO bioassay, and REC-M and REC-P seems to have been selected a total of nine generations (the figure shown only three selections). What was the calculation or method used to determine LC50 and LC95? R is a programming language and the software is only an environment to perform computations, this gives no information of how the LC50 was calculated. Maybe a link to a GitHub project or profile with the code used? Or a citation for the calculation methodology used? The authors also need to be more specific about how the resistance ratio was calculated. How were they compared? How was significance determined? How was control mortality accounted for? For figures 2 and 3, change "time point" to "generations" and to make it clearer and add a description of what the bars represent (error, st. dev., 95%CI?).

Colony: We include description for all three colonies used in this study (lines 85-100).

"A Total of three colonies from *Aedes aegypti* were used in this study:

Colony CAYMAN (CAY) was originally established in 2008 with *Aedes aegypti* collected in Grand Cayman island (Caribbean). This colony is highly resistant to pyrethroids and DDT, attributed to *kdr* alleles (F1534C and V1016I), and has been routinely selected with 0.75 % permethrin for 1 hour in the Liverpool Insect Testing Establishment (LITE: <https://lite.lstmed.ac.uk/lite-facilities/lite-insectaries/aedes-aegypti-cayman>).

Colony RECIFE (REC) was originally established in 2004 with *Aedes aegypti* collected in Araripina, Brazil (7° 32' S and 40° 34' W; Melo-Santos et al. 2010). This colony is resistant to temephos (OP), and biochemical assays indicate a higher activity of multiple detox enzyme families. Larvae from this colony have been routinely selected with 0.5 mg/L temephos for 24 hours.

Colony NEW ORLEANS is a susceptible colony established at LSTM in the 1970s. This colony originated in New Orleans (USA) and it is maintained in laboratory without insecticide exposure. Routine screening for target-site mutations associated with resistance indicates a lack of *kdr* alleles in the sodium channel and resistant alleles in Ace-1 (LITE: <https://lite.lstmed.ac.uk/lite-facilities/lite-insectaries/aedes-aegypti-new-orleans>)."

Stereochemistry: We include more information about the insecticides (lines 129-132)

"The adult mosquitoes were selected with either permethrin (Sigma Aldrich, Pestanal®, CAS # 52645-53-1, >95.0% sum of cis+trans 97.8%) or malathion (Sigma Aldrich, Pestanal®, CAS # 121-75-5) and larvae were selected with temephos (CHEM SERVICE INC., CAS # 3383-96-8) (see "*Mosquito selection regime*")."

Generations: We agree with the reviewer about the generation information in the text. To avoid confusion, we use the generation number starting in the beginning of our experiment for both Recife (G1-G10) and Cayman (G1-G11). The information about colony age at the beginning of our experiment was only included in the figure 1.

We updated:

- The text of the manuscript (mainly methods);
- Tables 1, 3, and 4;
- Figures 1 to 3;
- supplementary information: CT values adults.csv; Mortality after insecticide selection.csv; WHO bioassay results.csv; Table S1. Insecticide selection total mortality.csv.

Information about analysis:

Thank for your comment. We recognize that the information about analysis was incomplete.

- We include more details about the analysis and the R script.

Lines 235-244: "This process was repeated for each concentration in the assay. Larval mortality was recorded at 24 hours and Abbot's formula (Abbot 1925) was used to adjust mortality when necessary. The data from bioassays was organized by concentration in each biological replicate. These data were used to create general linear models (G.L.M.) with "logit" models for each strain using "glm" function in statistical software R-2.15.2. Lethal concentration and 95% confidence intervals were calculated using the r-package "MASS".

Resistance ratios were calculated based on comparison of LC_{50} and LC_{95} from each strain against the reference colony New Orleans, which is fully susceptible to all three insecticides. Confidence intervals for ratios were calculated using the method MOVER-R (Newcombe 2016) presented in the R-package "pairwiseCI."

- We update the Table 2 and 5
- We update the information about resistance ratios in Results (lines 273-277, and lines 297-300).
- We also include new references:

Abbott, W.S. 1925. A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.* **18**: 265–267. The Oxford University Press.

Newcombe RG. 2016. MOVER-R confidence intervals for ratios and products of two independently estimated quantities. *Stat Methods Med Res* **25**:1774–1778, SAGE Publications

Figure 2 and 3:

We include "Bars represent 95% CI."

Comment 4

The authors provide a great statement in the beginning for the discussion about the need

to perform more dose-response curves in insecticide resistance studies, but sadly this study also didn't exactly provide a dose-response curve. The issue with using a concentration, such as in the WHO assay, is that the exact amount of insecticide the individuals were exposed to remains unknown. Some animals may be more active and therefore be exposed to more by walking around the tube more than others. Also, an individual that is being affected may fall on its side more frequently and in the process get a higher exposure than a healthy animal that is standing straight. While LC assays are great for certain purposes, they are not optimal for determining resistance ratios and dose-response curves. Would it be possible to do topical assays on the final strains to get a dose response RR instead of a concentration response RR?

We understand the critique on using WHO assays to calculate lethal concentrations and resistance ratios. The calculation of doses based on mosquito weight may refine a resistance ratio calculation reducing certain limitations of WHO assays (or similar assays such as CDC bottle assay). However, topical assays also present disadvantages to study insecticide resistance since they require immobilization (e.g. using cold temperature, CO₂) and insecticide is normally apply in thorax or head (not the most common area of entry in the field). For this reason, we believe that WHO assays are a valid method for this manuscript, and they are a clear improvement over qualitative assays with reference concentrations.

We have included a discussion on the limitations of WHO assays for quantitative approach:

Lines 359-364: "Our method to calculate resistance ratios using WHO tubes is not able to define the individual dose received per mosquito, since mosquito weight and individual activity against the treated surface could not be measured. Topical assays, which don't mimic natural exposure routes, do allow for controlled application and would further reduce the individual variation observed within each strain."

We also added to the discussion:

Lines 303-306: "Inference of insecticide resistance in adults of *Ae. aegypti* is rarely performed by quantitative methods. A recent review (Moyes *et al.*, 2017) highlights the lack of literature that calculate resistance ratios based on dose-response, lethal concentration or lethal time (see S2 file in (Moyes *et al.*, 2017))."

Comment 5

In regard to using the New Orleans colony as the denominator in calculating the resistance ratio, a recent study showed that the NO strain has multiple kdr mutations¹, just something to keep in mind.

1. Fan Y, O'Grady P, Yoshimizu M, Ponlawat A, et al.: Evidence for both sequential mutations and recombination in the evolution of kdr alleles in *Aedes aegypti*. PLoS Negl Trop Dis. 14 (4): e0008154 PubMed Abstract | Publisher Full Text

Thank you for the concern. However, our susceptible New Orleans colony is an old colony without *Kdr* alleles such as "Liverpool", and "Rockefeller" (see comment 3). Moreover, Fan et

al. is focused on recent collections from Americas and Africa that includes recent field collection from New Orleans.

We include the description of all colonies used in the study (Lines 88-105).

In regard to the potential loss of diversity associated with bottlenecks limitations mentioned by the authors in the discussion section, perhaps backcrossing back to the parental colony after each selection could reduce this problem. With enough selections and backcrossing you should retain only the resistance mutations (if they were present in the original population) and any genes that are linked to the resistance alleles (this one is harder to separate out).

Thank you for this suggestion.

On a very minor note, the citation for Harris et al. 2010 was listed twice.

References were corrected to remove this duplication.

Competing Interests: No competing interests were disclosed.