RESEARCH ARTICLE

Synchrony between daily rhythms of malaria parasites and hosts is driven by an essential amino acid [version 1; peer review: 1 approved, 1 approved with reservations]

Kimberley F. Prior ^{1,2}, Benita Middleton³, Alíz T.Y. Owolabi^{1,2},
Mary L. Westwood ^{1,2}, Jacob Holland^{1,2}, Aidan J. O'Donnell^{1,2},
Michael J. Blackman^{4,5}, Debra J. Skene ³, Sarah E. Reece ^{1,2}

¹Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, UK

²Institute of Immunology & Infection Research, University of Edinburgh, Edinburgh, UK

³School of Biosciences and Medicine, University of Surrey, Surrey, UK

⁴Malaria Biochemistry Laboratory, Francis Crick Institute, London, UK

⁵Faculty of Infectious Diseases, London School of Hygiene & Tropical Medicine, London, UK

V1 First published: 22 Jul 2021, 6:186
<https://doi.org/10.12688/wellcomeopenres.16894.1>
Latest published: 20 Oct 2021, 6:186
<https://doi.org/10.12688/wellcomeopenres.16894.2>

Abstract

Background: Rapid asexual replication of blood stage malaria parasites is responsible for the severity of disease symptoms and fuels the production of transmission forms. Here, we demonstrate that the *Plasmodium chabaudi*'s schedule for asexual replication can be orchestrated by isoleucine, a metabolite provided to the parasite in periodic manner due to the host's rhythmic intake of food.

Methods: We infect female C57BL/6 and Per1/2-null TTFL clock-disrupted mice with 1×10^5 red blood cells containing *P. chabaudi* (DK genotype). We perturb the timing of rhythms in asexual replication and host feeding-fasting cycles to identify nutrients with rhythms that match all combinations of host and parasite rhythms. We then test whether perturbing the availability of the best candidate nutrient *in vitro* elicits changes their schedule for asexual development.

Results: Our large-scale metabolomics experiment and follow up experiments reveal that only one metabolite - the amino acid isoleucine - fits criteria for a time-of-day cue used by parasites to set the schedule for replication. The response to isoleucine is a parasite strategy rather than solely the consequences of a constraint imposed by host rhythms, because unlike when parasites are deprived of other essential nutrients, they suffer no apparent costs from isoleucine withdrawal.

Conclusions: Overall, our data suggest parasites can use the daily rhythmicity of blood-isoleucine concentration to synchronise asexual development with the availability of isoleucine, and potentially other resources, that arrive in the blood in a periodic manner due to the

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
20 Oct 2021	view	
version 1		
22 Jul 2021	view	view

1. **Luis Larrondo** , Millennium Institute for Integrative Biology (iBIO), Santiago, Chile
Pontificia Universidad Católica de Chile, Santiago, Chile

2. **Xiaodong Zhuang** , University of Oxford, Oxford, UK

Any reports and responses or comments on the article can be found at the end of the article.

host's daily feeding-fasting cycle. Identifying both how and why parasites keep time opens avenues for interventions; interfering with the parasite's time-keeping mechanism may stall replication, increasing the efficacy of drugs and immune responses, and could also prevent parasites from entering dormancy to tolerate drugs.

Keywords

Plasmodium, periodicity, circadian rhythm, metabolism, metabolomics, intraerythrocytic development cycle, isoleucine, asexual replication

Corresponding author: Kimberley F. Prior (priorkimberley@gmail.com)

Author roles: Prior KF: Formal Analysis, Investigation, Visualization, Writing – Original Draft Preparation; Middleton B: Methodology, Resources, Software; Owolabi ATY: Investigation, Methodology; Westwood ML: Formal Analysis, Investigation; Holland J: Investigation; O'Donnell AJ: Investigation, Software, Visualization; Blackman MJ: Conceptualization, Resources, Supervision; Skene DJ: Methodology, Resources, Software; Reece SE: Conceptualization, Project Administration, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Wellcome Trust [202769, <https://doi.org/10.35802/202769>]; 108905; FC001043] and the Royal Society [UF110155; NF140517].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Prior KF *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Prior KF, Middleton B, Owolabi ATY *et al.* Synchrony between daily rhythms of malaria parasites and hosts is driven by an essential amino acid [version 1; peer review: 1 approved, 1 approved with reservations] Wellcome Open Research 2021, 6:186 <https://doi.org/10.12688/wellcomeopenres.16894.1>

First published: 22 Jul 2021, 6:186 <https://doi.org/10.12688/wellcomeopenres.16894.1>

Introduction

Circadian rhythms are assumed to have evolved to coordinate organisms' activities with daily rhythms in the environment (Hozer *et al.*, 2020; Ouyang *et al.*, 1998; Spoelstra *et al.*, 2016). The value of organising sleep/wake cycles according to whether it is light or dark, whether predators or prey are active, *etc.*, is clear, but how parasites cope with rhythmicity in the within-host environment has been neglected (Martinez-Bakker & Helm, 2015; Reece *et al.*, 2017; Westwood *et al.*, 2019). Many of the processes underlying interactions between hosts and parasites have a circadian basis, yet why these rhythms exist and what their consequences are for hosts and parasites remain mysterious. For example, parasites are confronted with rhythms in host behaviours and physiologies, including immune responses and metabolism. Thus, some host rhythms offer opportunities for parasites to exploit, whilst other rhythms impose constraints parasites must cope with. Some parasites use their own circadian clocks to control metabolism (Rijo-Ferreira *et al.*, 2017), and virulence (Hevia *et al.*, 2015), suggesting that host rhythms are a selective (evolutionary) driver of parasite rhythms. Host rhythms have fitness consequences for malaria (*Plasmodium*) parasites (O'Donnell *et al.*, 2011; O'Donnell *et al.*, 2013), whose rhythmicity in development during blood-stage replication is aligned with the timing of host feeding-fasting cycles (Hirako *et al.*, 2018; O'Donnell *et al.*, 2020; Prior *et al.*, 2018).

Explaining the timing and synchrony exhibited by malaria parasites during successive cycles of blood-stage replication has been a puzzle since the Hippocratic Era, when the duration of these cycles (24-, 48, or 72-hours, depending on the *Plasmodium* spp.) were used to diagnose malaria infection (Garcia *et al.*, 2001; Mideo *et al.*, 2013). Blood-stage asexual replication is orchestrated by the intraerythrocytic development cycle (IDC), which is characterised by parasite invasion of red blood cells (RBC), growth, division, and finally bursting of RBC to release the next cohort of asexually replicating parasites. Given that asexual replication is responsible for the disease symptoms of malaria and fuels the production of transmission forms, explaining how and why the vast majority of *Plasmodium* species progress through the IDC in synchrony, and why transitions between these stages occur at particular times of day, may unlock novel interventions and improve the efficacy of existing treatments. However, such endeavours rely on identifying the precise host rhythm(s) that parasites align to and explaining why this matters for their fitness (Prior *et al.*, 2020).

Malaria parasites are able to keep time (Rijo-Ferreira *et al.*, 2020; Subudhi *et al.*, 2020) and organise their IDC schedule to coordinate with an aspect(s) of host feeding-fasting rhythms (Hirako *et al.*, 2018; Prior *et al.*, 2018), but not the act of eating itself (Rijo-Ferreira *et al.*, 2020), or processes directly controlled by canonical host circadian clocks (O'Donnell *et al.*, 2020). Coordination with host feeding-fasting rhythms may allow parasites to couple the nutritional needs of each IDC stage with circadian fluctuations in the concentrations of nutrients in the blood (Prior *et al.*, 2020). Whilst parasites are able to

scavenge most amino acids from haemoglobin digestion, and can biosynthesise some metabolites themselves, other nutrients are essential and must be taken up from the RBCs or blood plasma. Haemoglobin digestion and biosynthesis can occur at any time-of-day but essential nutrients that both host and parasite must acquire from the host's food (including glucose and the amino acid, isoleucine) follow circadian rhythms in the blood (Skene *et al.*, 2018). Hosts forage during their active phase and fast during the rest phase, which generally generates peaks in the blood concentrations of metabolites shortly after the onset of feeding and their lowest point (nadir) occurs during the fasting phase.

Nutritional needs vary across IDC stages (Figure 1A), with for example, requirements for glycerophospholipids and amino acids to fuel biogenesis, increasing as parasites progress through the IDC, and the release of parasite progeny at the end of the IDC consumes a lot of glucose (Déchamps *et al.*, 2010; Olszewski *et al.*, 2009). Whether essential nutrients are available around-the-clock at sufficient concentrations to satisfy the needs of all parasite cells within an infection is unknown, but in culture, IDC progression is affected by nutrient limitation. For example, parasites experiencing glucose limitation quickly mount starvation responses (Daily *et al.*, 2007) and isoleucine starvation rapidly induces dormancy (Babbitt *et al.*, 2012). Moreover, the nadir of blood glucose rhythms in the rest phase can be exacerbated by malaria infection (Hirako *et al.*, 2018). Thus, to avoid the problems of temporal resource limitation, we hypothesise that parasites match transitions between IDC stages to rhythmicity in the availability of essential nutrients in the blood, ensuring the most demanding IDC stages (trophozoites and schizonts) coincide with when resources are most abundant. Parasites could synchronise with resource rhythms by using an essential nutrient itself as a time-cue to set the schedule for IDC transitions, and/or allow the host to impose the IDC schedule by, for example, selectively killing IDC stages that are not "on time". By getting the timing of the IDC schedule right, parasites garner fitness in two ways. They maximise asexual replication, which underpins within-host survival (O'Donnell *et al.*, 2011; O'Donnell *et al.*, 2013), and benefit from coordinating the production of sexual transmission stages with the time-of-day their vectors forage for blood (Pigeault *et al.*, 2018; Schneider *et al.*, 2018).

We integrate evolutionary ecology and parasitology with chronobiology to, first, undertake a hypothesis-driven screen of several metabolite classes and glucose to identify nutrients with daily fluctuations in the blood of malaria-infected hosts. Second, we determine which metabolites follow rhythms set by the timing of different perturbations of host feeding-fasting rhythms as well as matching the timing of the IDC schedules in all treatment groups. We find that the IDC schedule cannot be explained by fluctuations in blood glucose concentrations and almost all metabolites in the screen, but experiment does reveal a single candidate; the amino acid, isoleucine. Third, we test if the timing of isoleucine provision and withdrawal affects IDC progression in the manner expected if parasites use it as a time-of-day cue to schedule their

Figure 1. Rhythms in the intraerythrocytic development cycle (IDC) and host rhythms. (A) The role of each IDC stage during cycles of asexual replication, and the resources known to be essential to each stage. RG=ring stage, ET=early stage trophozoite, MT=mid stage trophozoite, LT=late stage trophozoite, SZ=schizont. (B), the experimental design used wild type mice housed in a 12h light: 12h dark regime (indicated by the light-dark bar) with unrestricted access to food for 12 hours either during the night-times (DF, dark feeding) or the day times (LF, light feeding) as indicated by the position of the cheeses. *Per1/2*-null mice were housed in continuous darkness (DD) and either experienced cycles of 12-hours with food followed by 12 h without access to food (TRF, time restricted feeding,) or given constant access to food (ALF, *ad libitum* feeding). The parasite IDC is rhythmic in DF, LF and TRF mice but not ALF mice (O'Donnell *et al.*, 2020) causing the IDC rhythm to be substantially dampened. (C) Metabolites that were significantly rhythmic in DF, LF and TRF mice (highlighted in red), but not in ALF mice, were sought. Treatment groups colour coded throughout: DF=orange, LF=light blue, TRF=green, ALF=dark blue.

IDC or in the manner expected if the IDC schedule is imposed by starvation of mis-timed IDC stages. We capitalise on the rodent malaria *P. chabaudi* model system in which *in vivo* experiments exploit ecologically relevant host-parasite interactions and short-term *in vitro* studies allow within-host ecology to be probed in-depth.

Results

Metabolites that associate with host feeding rhythms and the IDC schedule

To identify metabolites whose rhythms correspond to the timing of host feeding and the IDC schedule, we compared four groups of malaria infections in mice that were either wild type (WT) C57BL/6J strain or *Per1/2*-null (Period 1 and Period 2) circadian clock-disrupted mice. *Per1/2*-null mice have an impaired canonical clock (transcription translation feedback loop, TTFL) and exhibit no circadian rhythms in physiology or behaviour when kept in constant darkness (Bae *et al.*, 2001; Maywood *et al.*, 2014; O'Donnell *et al.*, 2020). We generated three different groups of hosts whose feeding-fasting rhythms differed with respect to the light:dark cycle and whether they had an intact TTFL clock, and a 4th group of hosts which lacked both feeding rhythms and an intact TTFL clock (Figure 1B). All infections were sampled every two hours from day 5 post infection (when infections are at a low parasitaemia, ~10%, to minimise the contribution of parasite molecules to the dataset, Olszewski *et al.*, 2009) for 26 hours. We hypothesised that a time-cue/time-dependent resource must vary or have rhythmicity across 24 hours, with a peak concentration that associates with the timing of host feeding-fasting across the three treatment groups with rhythmic feeding, yet also be arrhythmic in the 4th group (Figure 1B). In addition, the parasite IDC must be rhythmic in the three treatment groups with rhythmic feeding, with the same IDC stage present at the time of feeding, and be arrhythmic in the 4th group. Thus, we identified candidate metabolites by intersecting rhythmic metabolites in the treatment groups (Figure 1C).

IDC schedules followed the expected patterns for each treatment group (Figure 2, and O'Donnell *et al.*, 2020). Specifically, parasites displayed opposite IDC schedules in dark (i.e. night, DF, *n*=18) and light (i.e. day, LF, *n*=17) fed wild type (WT) mice because their feeding-fasting rhythms are inverted (Figure 2, Table 1). Parasites in *Per1/2*-null mice (kept in constant darkness), fed during a time-restricted 12 h window (TRF mice, *n*=17; Table 2) that coincided with when LF mice were fed, followed the same IDC schedule as parasites in LF mice. Finally, parasites in *Per1/2*-null mice (kept in constant darkness) with constant access to food (ALF mice, *n*=16; Table 2) exhibited dampened IDC rhythms because their hosts have no feeding-fasting rhythm (O'Donnell *et al.*, 2020). Across the entire data set, 110 metabolites exhibited variation in their blood concentrations during the 26-hour sampling window (101 in DF, 91 in LF, 50 in TRF and one in ALF; Table 3). That only one metabolite (lysoPC a C20:3) exhibited a rhythm in ALF hosts demonstrates that host TTFL clocks and timed feeding are required to generate metabolite rhythms (Figure 3A, Reinke & Asher, 2019). Further, that approximately half of the metabolites

rhythmic in hosts with TTFL clocks (DF and LF) were also rhythmic in TRF hosts reveals that feeding rhythms alone can maintain metabolic rhythms during infection (Figure 3A). Of all metabolites, only 42 were rhythmic in all three groups of infections with both feeding-fasting and IDC rhythms (i.e. the red area in Figure 1C), consisting of three acylcarnitines, 11 amino acids, nine biogenic amines and 19 glycerophospholipids (Figure 3A). Next, we identified 33 metabolites that exhibited a peak phase (timing) that corresponded to the timing of the host's feeding-fasting rhythm and the parasite's IDC (Figure 3B; metabolites clustering in the blue and purple areas).

Glucose does not associate with the IDC schedule

Previous work suggests that blood glucose rhythms are responsible for the IDC schedule (Hirako *et al.*, 2018; Prior *et al.*, 2018) but glucose could not be measured by the UPLC/MS-MS metabolomics platform. Therefore, we set up an additional set of DF, LF, TRF and ALF infections (*n*=5/group), which followed the same feeding-fasting and IDC schedules as infections in the metabolomics screen, Figure 2, Table 2), to quantify blood-glucose rhythms. Mean blood-glucose concentration differed between the groups, being higher in DF and TRF mice (DF=8.55±0.14 mmol/L, TRF=8.59±0.13 mmol/L) than in LF and ALF mice (LF=7.90±0.14 mmol/L, ALF=7.68±0.11 mmol/L). We found that two competitive models, including only time-of-day (Zeitgeber time, ZT/h) or both time-of-day and treatment (DF, LF, TRF and ALF) as main effects, can explain patterns of glucose concentration (Figure 4, Table 4). Specifically, glucose concentration varied throughout the day in DF mice but much less in LF mice, and glucose was invariant in both groups of *Per1/2*-null mice (TRF and ALF) (Table 4). The rhythmic IDC schedule in the DF, LF and TRF groups but the lack of significant rhythmicity in blood glucose in the TRF (and possibly LF) infections demonstrates that glucose does not explain the connection between feeding rhythms and the IDC schedule across all groups.

Linking metabolites to the IDC schedule

Having ruled out a role for blood glucose, we explored whether the metabolites from our screen could explain why the IDC schedule follows feeding-fasting rhythms. The metabolites whose peak associated with both feeding and late IDC stages, are alanine, asparagine, isoleucine, leucine, methionine, phenylalanine, proline, threonine, valine, methionine-sulfoxide and serotonin. Whereas 20 acylcarnitines and glycerophospholipids (with the exception of two biogenic amines) associated with fasting and early IDC stages (ADMA, SDMA, C14:1, C16, C18:1, lysoPCaC16:1, lysoPCaC18:1, lysoPCaC18:2, PCaaC32:1, PCaaC34:4, PCaaC38:3, PCaaC38:4, PCaaC38:5, PCaaC38:6, PCaaC40:4, PCaaC40:5, PCaeC34:3, PCaeC38:0, PCaeC38:3 and PCaeC42:1; Table 5 and Table 6). We asked if these metabolites could schedule the IDC through the following non-mutually exclusive mechanisms. First, by being sufficiently limiting at a certain time-of-day to enforce a rhythm on the IDC because mis-timed IDC stages starve and die. This scenario requires that parasites are unable to overcome resource limitation by synthesising the metabolite itself or scavenging it from a source (such as haemoglobin) that is available around

Figure 2. Intraerythrocytic development cycle (IDC) schedule and host feeding rhythms. The proportion of parasites at ring stage (phase marker) for: **(A)** DF: dark feeding WT mice food access ZT 12-24 (12 h at night) in 12h:12h light:dark; **(B)** LF: light feeding WT mice food access ZT 0-12 (12 h in day) in 12h:12h light:dark; **(C)** TRF: time restricted feeding Per1/2-null mice food access 0-12 hours (12 h at the same time (GMT) as LF mice) in constant darkness (DD). **(D)** ALF: *ad libitum* feeding Per1/2-null mice access in constant darkness (see Figure 1 for experimental design and more details). Feeding-fasting rhythms are indicated by cheeses, the white panel denotes lights on (Zeitgeber Time=0-12 h) for DF and LF, 0-24 h for TRF and ALF). Model fits for each group are plotted on the raw data (n= 2-5 infections per time point). The fitted sine/cosine curve for DF **(A)** is distorted due to a large amplitude which exceeds the bounds possible for proportion data (between 0 and 1) so is truncated for visualisation. The patterns for groups are explained better by sine/cosine curves than by a straight line (see Table 1).

the clock. Second, by acting as a time-of-day cue to which certain IDC stages respond by for example, initiating the transition to the next stage (called a “just-in-time” strategy) or by acting as a Zeitgeber to entrain an endogenous circadian clock that allows parasites to anticipate when IDC transitions should occur. For a metabolite to be a reliable time-of-day cue/Zeitgeber, it should be something the parasite cannot synthesise/scavenge to avoid the challenge of differentiating between inaccurate endogenous and accurate exogenous time information leading to the risk of mistakenly responding to an endogenous signal. Of the 33 metabolites on the shortlist, only isoleucine fulfils these criteria (Table 7, Figure 5), suggesting that parasites could use isoleucine both as a resource and a time-of-day cue.

Timing and completion of the parasite IDC depends on the availability of isoleucine

That malaria parasites rely on exogenous isoleucine as a both a resource and time-cue to complete the IDC is supported by previous observations across species of *Plasmodium*. First, when the human malaria *P. falciparum* is deprived of isoleucine, parasites immediately and dramatically slow cell cycle progression akin to dormancy (Liu *et al.*, 2006) yet are able to recover even after long periods of starvation (~4 IDCs) (Babbitt *et al.*, 2012). Second, in contrast to isoleucine, if other amino acids are removed from culture media, parasites switch to scavenging them from haemoglobin with only minor effects on IDC completion (Babbitt *et al.*, 2012). Third, not only is

Table 1. Degrees of Freedom (df), log-Likelihood (logLik), AICc, $\Delta AICc$ ($AICc_i - \Delta AICc_{min}$) and AICc w ($AICc$ weight) for each linear model in the parasite stage proportion analysis ordered in descending fit (best-fitting model at the top). The response variable for each model is proportion of ring stages (Ring.prop), with "sine" and "cosine" terms being the sine or cosine function of ($2\pi \times$ time of day)/24 with a fixed 24-hour period fitted for each treatment group (DF, LF, TRF or ALF). AICc is a form of the Akaike Information Criteria corrected for smaller sample sizes to address potential overfitting, used for model selection. DF=dark fed wild type mice, LF=light fed wild type mice, TRF=time restricted fed Per1/2-null mice, ALF=ad libitum fed Per1/2-null mice.

	Model description: Ring.prop ~	df	logLik	AICc	$\Delta AICc$	AICc w
DF	sine + cosine	4	13.30	-16.9	0.00	1.000
	sine	3	-5.77	18.5	35.44	0.000
	cosine	3	-8.69	24.3	41.27	0.000
	null	2	-14.60	33.7	50.60	0.000
LF	sine + cosine	4	8.57	-7.5	0.00	0.788
	sine	3	5.92	-4.9	2.63	0.212
	cosine	3	-1.03	9.0	16.52	0.000
	null	2	-2.68	9.8	17.34	0.000
TRF	sine + cosine	4	21.85	-33.5	0.00	1.000
	sine	3	10.98	-14.7	18.77	0.000
	cosine	3	-0.14	7.5	41.01	0.000
	null	2	-2.54	9.7	43.15	0.000
ALF	sine + cosine	4	32.12	-53.6	0.00	0.995
	sine	3	24.51	-41.5	12.05	0.002
	cosine	3	24.49	-41.5	12.08	0.002
	null	2	19.90	-35.1	18.47	0.000

Table 2. Mouse numbers for experimental treatment groups. A) Mice in the metabolomics experiment were sampled in blocks (A–D) with 4/5 mice per block sampled every eight hours. Totals for each treatment group: DF=18, LF=17, TRF=17, ALF=16. B) Mice in the glucose experiment were sampled every two hours. Totals for each treatment group: DF=5, LF=5, TRF=5, ALF=5. For each experiment repeated measures from mice were controlled for during the analysis. DF=dark fed wild type mice, LF=light fed wild type mice, TRF=time restricted fed Per1/2-null mice, ALF=ad libitum fed Per1/2-null mice.

Time (ZT/hour)	0	2	4	6	8	10	12	14	16	18	20	22	24/0	2
A Metabolomics experiment														
Block	A	B	C	D	A	B	C	D	A	B	C	D	A	B
DF	5	5	4	4	5	5	4	4	5	5	4	4	5	5
LF	5	4	4	4	5	4	4	4	5	4	4	4	5	4
TRF	5	4	4	4	5	4	4	4	5	4	4	4	5	4
ALF	4	4	4	4	4	4	4	4	4	4	4	4	4	4

Time (ZT/hour)	0	2	4	6	8	10	12	14	16	18	20	22	24/0	2
B Glucose experiment														
DF	5	5	5	5	5	5	5	5	5	5	5	5	5	5
LF	5	5	5	5	5	5	5	5	5	5	5	5	5	5
TRF	5	5	5	5	5	5	5	5	5	5	5	5	5	5
ALF	5	5	5	5	5	5	5	5	5	5	5	5	5	5

Table 3. Metabolite numbers that significantly fluctuate every 24 hours in the mouse blood for three methods. A) Data for all metabolites were run through three circadian programmes to find those following a 24h rhythm using ECHO (Benjamini-Hochberg (BH) adjusted p value of 0.05), CircWave (standard p value of 0.05) and JTK_Cycle (BH adjusted p value of 0.05). Metabolites that are excluded using the Surrey LC/MS assay criteria were removed from the analysis (Figure 9). **B)** Rhythmic metabolites in each programme (ECHO, CircWave and JTK) were intersected, with a metabolite counted as rhythmic if it is significantly rhythmic in at least two programmes (ECHO=CW, ECHO=JTK, CW=JTK). **C)** The metabolites not fulfilling these criteria were analysed using ANOVA (including time-of-day as a factor), with BH adjusted p values at the 5% level. Metabolites from both methods (circadian programmes and ANOVA) were then combined to perform a final intersection between DF, LF and TRF to find common metabolites. Metabolites rhythmic in ALF mice were then removed. DF=dark fed wild type mice, LF=light fed wild type mice, TRF=time restricted fed Per1/2-null mice, ALF=ad libitum fed Per1/2-null mice.

A	ECHO		CircWave (CW)		JTK	
	Rhythmic with 24h period	Not-rhythmic with 24h period	Rhythmic with 24h period	Not-rhythmic with 24h period	Rhythmic with 24h period	Not-rhythmic with 24h period
DF	75	59	67	67	47	87
LF	106	28	83	51	70	64
TRF	55	77	41	93	23	111
ALF	2	127	6	128	0	134
B	ECHO==CW	ECHO==JTK	CW==JTK		Unique	
DF	62	47	46		63	
LF	79	68	63		88	
TRF	37	23	21		39	
ALF	1	0	0		1	
C	Not-rhythmic with 24h period	Time-of-day effect	No time-of-day effect	Time-of-day effect + rhythmic with 24h period	Total candidates	
DF	71	38	33	101	42	
LF	46	3	43	91		
TRF	95	11	84	50		
ALF	133	0	133	1		

isoleucine crucial for the development of *P. knowlesi* in culture, parasites only incorporate isoleucine for part of the IDC (until the point that schizogony starts (Butcher & Cohen, 1971; Polet & Conrad, 1968; Polet & Conrad, 1969; Sherman, 1979)). Fourth, isoleucine is one of few amino acids that exhibits a daily rhythm in the blood of mice and humans. Specifically,

isoleucine was inverted with a 12-hour shift in simulated day and night shift working humans (a phase difference of $11:49 \pm 02:10$ h between the day and night shift conditions) and follows the timing of food intake (Skene *et al.*, 2018). Therefore, our next experiments tested whether an exogenous supply of isoleucine is capable of scheduling the IDC.

Figure 3. Rhythmic metabolites that associate with intraerythrocytic development cycle (IDC) timing and host feeding-fasting cycles. (A) The numbers of rhythmic intersecting metabolites out of a total of 109. (B) Peak timing (phase) concentration of the 42 metabolites that are rhythmic in DF, LF and TRF infections. Top left panel (light blue) denotes peaks during the fasting phase and the bottom right panel (purple) denotes peaks in the feeding phase, yielding 33 candidates linked to the feeding-fasting cycle. Metabolite classes: acylcarnitines=green circles, amino acids/biogenic amines=orange triangles, glycerophospholipids=purple squares. See Table 6 for peak times of each metabolite. Ring stages (RG), late trophozoites (LT) and schizonts (SZ) peak during the hosts feeding period (purple) when some amino acids/biogenic amines peak. Early trophozoites (ET) and mid trophozoites (MT) peak during host fasting (light blue) when glycerophospholipids and acylcarnitines peak.

We carried out two experiments in parallel to quantify how *P. chabaudi*'s IDC progression is affected when deprived of isoleucine, and whether the IDC is then completed (defined as the proportion of parasites that reach the schizont stage) when isoleucine is restored. Isoleucine cannot be perturbed *in vivo* without causing confounding off-target effects on the host or interference by host homeostasis, so we carried these

experiments *in vitro* where conditions can be controlled. First, parasites cultured in the absence of isoleucine (n=32 cultures from the blood of eight mice, which were split equally across both treatments) develop extremely slowly with approximately three-fold fewer completing the IDC compared to parasites with isoleucine (50 mg/L, which is the same concentration as standard culture media; RPMI 1640) (n=32 cultures) (Figure 6A).

Figure 4. Blood glucose concentrations (mmol/L). (A) DF: dark feeding WT mice food access ZT 12-24 (12 h at night) in 12h:12h light: dark, (B) LF: light feeding WT mice food access ZT 0-12 (12 h in day) in 12 h:12 h light:dark. (C) TRF: time restricted feeding Per1/2-null mice food access 0-12 hours (12 h at the same time (GMT) as LF mice) in constant darkness (DD). (D) ALF: *ad libitum* feeding Per1/2-null mice access in constant darkness (see Figure 1 for experimental design). Feeding-fasting rhythms are indicated by cheeses, the white panel denotes lights on (Zeitgeber Time=0-12 h), dark grey panel denotes lights off (Zeitgeber Time=12-24 h for DF and LF, 0-24 h for TRF and ALF). The lines and shading are mean $\pm$ SEM at each time point and the dots are the raw data. At time points 0.5 and 2.5 for DF, TRF and ALF, and 12.5 and 14.5 for LF the points are stacked because the time course lasted ~26 hours and the data are plotted on a 0-24 hour axis.

The best fitting model contained only “treatment” (parasites cultured with or without isoleucine) as a main effect ($\Delta\text{AICc}=0$, Table 8A). The reduction in schizonts in isoleucine-free conditions was not due to a higher death rate because the density of parasites remains constant during the experiment and did not differ between the treatments (Figure 6B) (best fitting model is the null model $\Delta\text{AICc}=0$, Table 8B). Further, incorporating either “treatment” or “hours in culture” into the model did not improve the model fit (treatment: $\Delta\text{AICc}=4.16$, hours in culture: $\Delta\text{AICc}=5.50$, Table 8B). This experiment extends previous findings for *P. falciparum* (Babbitt *et al.*, 2012; Liu *et al.*, 2006;

McLean & Jacobs-Lorena, 2020) across species as well as provides greater resolution on the parasite phenotypes involved.

The substantial slowing of IDC progression in the absence of isoleucine is further supported by our second experiment. This experiment revealed that like for *P. falciparum*, *P. chabaudi* completes IDC development when isoleucine deprivation ends. Parasites ($\sim 10^7$ per culture) were added to isoleucine-free media and incubated for seven, 14, or 18 hours, after which isoleucine (50 mg/L) was added to their cultures ($n=16$ cultures per treatment). Parasites completed development when

Table 4. Degrees of Freedom (df), log-Likelihood (logLik), AICc, $\Delta AICc$ ($AICc_i - AICc_{min}$) and AICc w (AICc weight) for each linear model in the glucose concentration analysis ordered in descending fit (best-fitting model at the top). The response variable for each model is glucose concentration (Glucose.conc) and the random effect is "mouseID". "Treatment" refers to the treatment group (DF, LF, TRF or ALF) and "time" refers to the time-of day (ZT 0-24h or 0-24h) which was fitted as a factor. DF=dark fed wild type mice, LF=light fed wild type mice, TRF=time restricted fed Per1/2-null mice, ALF=ad libitum fed Per1/2-null mice.

	Model description: Glucose.conc ~ + (1 mouseID)	df	logLik	AICc	$\Delta AICc$	AICc w
	time	14	-369.71	769.0	0.00	0.555
	treatment + time	17	-366.55	769.4	0.44	0.445
	treatment*time	50	-331.38	785.1	16.12	0.000
	null	3	-390.80	787.7	18.68	0.000
	treatment	6	-387.75	787.8	18.80	0.000
DF	time	14	-90.06	215.7	0.00	0.996
	null	3	-110.28	226.9	11.17	0.004
LF	null	3	-94.83	196.0	0.00	0.550
	time	14	-80.39	196.4	0.40	0.450
TRF	null	3	-85.23	176.8	0.00	0.999
	time	14	-77.54	190.7	13.9	0.001
ALF	null	3	-88.02	182.4	0.00	0.994
	time	14	-78.43	192.6	10.21	0.006

Table 5. Highlighted in green are the metabolites rhythmic in each treatment group (according to the circadian programmes and ANOVA). DF: 101 y, 33 n; LF: 91 y, 43 n; TRF: 50 y, 84 n; ALF: 1 y, 133 n.

Metabolite	Class	DF	LF	TRF	ALF
Ala	Amino.acid				
Arg	Amino.acid				
Asn	Amino.acid				
Asp	Amino.acid				
Cit	Amino.acid				
Gln	Amino.acid				
Glu	Amino.acid				
Gly	Amino.acid				
His	Amino.acid				
Ile	Amino.acid				
Leu	Amino.acid				
Lys	Amino.acid				

Metabolite	Class	DF	LF	TRF	ALF
Met	Amino.acid				
Orn	Amino.acid				
Phe	Amino.acid				
Pro	Amino.acid				
Ser	Amino.acid				
Thr	Amino.acid				
Trp	Amino.acid				
Tyr	Amino.acid				
Val	Amino.acid				
ADMA	Biogenic_amine				
alpha.AAA	Biogenic_amine				
Carnosine	Biogenic_amine				
Histamine	Biogenic_amine				
Kynurenine	Biogenic_amine				
Met.SO	Biogenic_amine				
Putrescine	Biogenic_amine				
Sarcosine	Biogenic_amine				
SDMA	Biogenic_amine				
Serotonin	Biogenic_amine				
Spermidine	Biogenic_amine				
t4.OH.Pro	Biogenic_amine				
Taurine	Biogenic_amine				
C0	Acylcarnitines				
C2	Acylcarnitines				
C3	Acylcarnitines				
C4	Acylcarnitines				
C14.1	Acylcarnitines				
C16	Acylcarnitines				
C18.1	Acylcarnitines				
lysoPC.a.C16.0	Glycerophospholipids				
lysoPC.a.C16.1	Glycerophospholipids				
lysoPC.a.C17.0	Glycerophospholipids				
lysoPC.a.C18.0	Glycerophospholipids				
lysoPC.a.C18.1	Glycerophospholipids				
lysoPC.a.C18.2	Glycerophospholipids				
lysoPC.a.C20.3	Glycerophospholipids				
lysoPC.a.C20.4	Glycerophospholipids				

Metabolite	Class	DF	LF	TRF	ALF
lysoPC.a.C24.0	Glycerophospholipids				
lysoPC.a.C26.0	Glycerophospholipids				
lysoPC.a.C26.1	Glycerophospholipids				
lysoPC.a.C28.0	Glycerophospholipids				
lysoPC.a.C28.1	Glycerophospholipids				
PC.aa.C24.0	Glycerophospholipids				
PC.aa.C28.1	Glycerophospholipids				
PC.aa.C30.0	Glycerophospholipids				
PC.aa.C32.0	Glycerophospholipids				
PC.aa.C32.1	Glycerophospholipids				
PC.aa.C32.2	Glycerophospholipids				
PC.aa.C32.3	Glycerophospholipids				
PC.aa.C34.1	Glycerophospholipids				
PC.aa.C34.2	Glycerophospholipids				
PC.aa.C34.3	Glycerophospholipids				
PC.aa.C34.4	Glycerophospholipids				
PC.aa.C36.1	Glycerophospholipids				
PC.aa.C36.2	Glycerophospholipids				
PC.aa.C36.3	Glycerophospholipids				
PC.aa.C36.4	Glycerophospholipids				
PC.aa.C36.5	Glycerophospholipids				
PC.aa.C36.6	Glycerophospholipids				
PC.aa.C38.0	Glycerophospholipids				
PC.aa.C38.3	Glycerophospholipids				
PC.aa.C38.4	Glycerophospholipids				
PC.aa.C38.5	Glycerophospholipids				
PC.aa.C38.6	Glycerophospholipids				
PC.aa.C40.2	Glycerophospholipids				
PC.aa.C40.3	Glycerophospholipids				
PC.aa.C40.4	Glycerophospholipids				
PC.aa.C40.5	Glycerophospholipids				
PC.aa.C40.6	Glycerophospholipids				
PC.aa.C42.0	Glycerophospholipids				
PC.aa.C42.1	Glycerophospholipids				
PC.aa.C42.2	Glycerophospholipids				
PC.aa.C42.4	Glycerophospholipids				
PC.aa.C42.5	Glycerophospholipids				
PC.aa.C42.6	Glycerophospholipids				

Metabolite	Class	DF	LF	TRF	ALF
PC.ae.C30.1	Glycerophospholipids				
PC.ae.C30.2	Glycerophospholipids				
PC.ae.C32.1	Glycerophospholipids				
PC.ae.C32.2	Glycerophospholipids				
PC.ae.C34.0	Glycerophospholipids				
PC.ae.C34.1	Glycerophospholipids				
PC.ae.C34.2	Glycerophospholipids				
PC.ae.C34.3	Glycerophospholipids				
PC.ae.C36.0	Glycerophospholipids				
PC.ae.C36.1	Glycerophospholipids				
PC.ae.C36.2	Glycerophospholipids				
PC.ae.C36.3	Glycerophospholipids				
PC.ae.C36.4	Glycerophospholipids				
PC.ae.C36.5	Glycerophospholipids				
PC.ae.C38.0	Glycerophospholipids				
PC.ae.C38.1	Glycerophospholipids				
PC.ae.C38.2	Glycerophospholipids				
PC.ae.C38.3	Glycerophospholipids				
PC.ae.C38.4	Glycerophospholipids				
PC.ae.C38.5	Glycerophospholipids				
PC.ae.C38.6	Glycerophospholipids				
PC.ae.C40.1	Glycerophospholipids				
PC.ae.C40.2	Glycerophospholipids				
PC.ae.C40.3	Glycerophospholipids				
PC.ae.C40.4	Glycerophospholipids				
PC.ae.C40.5	Glycerophospholipids				
PC.ae.C40.6	Glycerophospholipids				
PC.ae.C42.1	Glycerophospholipids				
PC.ae.C42.2	Glycerophospholipids				
PC.ae.C42.3	Glycerophospholipids				
PC.ae.C44.3	Glycerophospholipids				
PC.ae.C44.5	Glycerophospholipids				
PC.ae.C44.6	Glycerophospholipids				
SM.OH..C14.1	Sphingolipids				
SM.OH..C16.1	Sphingolipids				
SM.OH..C22.1	Sphingolipids				
SM.OH..C22.2	Sphingolipids				

Metabolite	Class	DF	LF	TRF	ALF
SM.OH..C24.1	Sphingolipids				
SM.C16.0	Sphingolipids				
SM.C16.1	Sphingolipids				
SM.C18.0	Sphingolipids				
SM.C18.1	Sphingolipids				
SM.C20.2	Sphingolipids				
SM.C24.0	Sphingolipids				
SM.C24.1	Sphingolipids				
SM.C26.0	Sphingolipids				
SM.C26.1	Sphingolipids				

Table 6. Timing of peak of each final candidate metabolite in the blood for each treatment group. For LF and TRF groups ZT 0/ 0 hours is the start of the feeding window and the time of lights on for LF, while for DF ZT0 is the start of the fasting window and the time of lights on. LF and DF groups are in 12h:12h light:dark although feeding in the day and night, respectively. TRF are in constant darkness although feed for the same 12h window (0-12 hours) as the LF group (same experimenter time). DF=dark fed wild type mice, LF=light fed wild type mice, TRF=time restricted fed Per1/2-null mice.

Metabolite	Class	DF (ZT, hours)	LF (ZT, hours)	TRF (hours)
C14.1	Acylcarnitines	7.12	20.85	19.13
C16	Acylcarnitines	6.83	19.97	20.44
C18.1	Acylcarnitines	6.65	19.92	20.18
Ala	Amino.acid	19.17	7.99	7.38
Asn	Amino.acid	18.47	6.39	6.03
Asp	Amino.acid	17.46	20.42	18.01
Glu	Amino.acid	17.59	20.08	18.38
Ile	Amino.acid	21.03	7.70	0.40
Leu	Amino.acid	20.36	6.65	2.85
Met	Amino.acid	19.90	6.20	5.90
Phe	Amino.acid	19.78	6.99	5.91
Pro	Amino.acid	19.08	6.95	6.34
Thr	Amino.acid	20.06	7.93	6.33
Val	Amino.acid	20.50	7.95	5.05
ADMA	Biogenic_amine	9.74	20.85	19.43
Carnosine	Biogenic_amine	16.39	20.41	19.03
Histamine	Biogenic_amine	15.27	19.06	19.21
Met.SO	Biogenic_amine	19.37	8.04	6.54
Putrescine	Biogenic_amine	15.80	19.05	16.13
SDMA	Biogenic_amine	9.83	20.78	20.54

Metabolite	Class	DF (ZT, hours)	LF (ZT, hours)	TRF (hours)
Serotonin	Biogenic_amine	13.31	6.29	18.74
Spermidine	Biogenic_amine	16.54	19.78	19.13
Taurine	Biogenic_amine	16.07	20.22	19.60
lysoPCaC16:1	Glycerophospholipids	6.50	18.17	18.20
lysoPCaC18:1	Glycerophospholipids	6.97	18.57	16.64
lysoPCaC18:2	Glycerophospholipids	5.90	15.27	12.05
PCaaC32:1	Glycerophospholipids	1.41	16.26	18.89
PCaaC32:2	Glycerophospholipids	23.87	16.15	18.57
PCaaC34:4	Glycerophospholipids	6.71	17.92	0.54
PCaaC38:3	Glycerophospholipids	7.48	19.01	22.51
PCaaC38:4	Glycerophospholipids	8.18	21.07	1.18
PCaaC38:5	Glycerophospholipids	7.70	20.03	0.89
PCaaC38:6	Glycerophospholipids	6.81	20.36	23.97
PCaaC40:4	Glycerophospholipids	5.80	18.63	23.57
PCaaC40:5	Glycerophospholipids	7.01	19.68	23.88
PCaeC34:1	Glycerophospholipids	23.59	16.45	18.99
PCaeC34:3	Glycerophospholipids	0.48	15.22	14.65
PCaeC36:2	Glycerophospholipids	22.36	12.23	10.04
PCaeC38:0	Glycerophospholipids	7.80	19.97	0.71
PCaeC38:2	Glycerophospholipids	20.45	12.78	9.58
PCaeC38:3	Glycerophospholipids	2.18	16.49	19.91
PCaeC42:1	Glycerophospholipids	8.78	20.70	2.72

isoleucine became available, regardless of the duration of deprivation (seven, 14, or 18 hours), with the best fitting model containing main effects of “treatment” and “hours since isoleucine added” ($\Delta AIC_c=0$, Table 8C). Importantly, including the interaction did not improve the model fit ($\Delta AIC_c=13.65$, Table 8C), demonstrating that IDC completion proceeds at the same rate despite different durations of isoleucine starvation. Specifically, the rate of IDC completion in the 6–9 hours following isoleucine replenishment was approximately 50% for all groups (Figure 6C). Again, higher death rates in cultures deprived of isoleucine for the longest time period did not give the appearance of equal IDC rates because the model incorporating “hours since isoleucine” was competitive with the null model ($\Delta AIC_c=0.56$, Figure 6D, Table 8D), revealing parasites were still as viable after 18 hours of deprivation as those deprived for seven and 14 hours. Furthermore, cultures deprived the longest achieved the most schizonts (18 hours, mean $\pm$ SEM: $1.61 \times 10^6 \pm 0.20$ Figure 6C), while the least schizonts were observed in cultures deprived for the shortest period (seven hours, mean $\pm$ SEM: $0.83 \times 10^6 \pm 0.14$ Figure 6C). Unlike previous

studies, by comparing rates of completion across treatments (i.e. statistically testing the roles of the duration of deprivation and its interaction with treatment), our experiment demonstrates that isoleucine deprivation is not costly to parasites (Figure 7A). The variation in the intercepts of Figure 6C (<1% schizonts after seven hours, 3% after 14 hours and 5% after 18 hours deprivation) is likely explained by the 18 hour deprivation cultures accumulating a higher proportion of schizonts at the time of isoleucine provision simply as a product of developing very slowly during a longer window of deprivation (in line with Liu *et al.*, 2006 and Babbitt *et al.*, 2012). The lack of costs and ability to recover development when isoleucine is replenished - which differs from the response to the loss of other essential nutrients - is consistent with isoleucine being sufficient to act as a time-cue as well as being an essential resource (Figure 7B).

Discussion

Our large-scale metabolomics screening experiment (Figure 3–Figure 5, Table 7) and follow-up experiments (Figure 6), demonstrate that isoleucine is sufficient to explain

Table 7. Potential for acylcarnitines, glycerophospholipids and amino acids/amines candidates that associate with the feeding-fasting cycle to explain the intraerythrocytic development cycle (IDC) schedule. Requirements for these metabolites is likely to increase as each parasite cell progresses through its IDC.

Metabolite class & general function	Role in the IDC?	Putative time-cue?
Acylcarnitines <ul style="list-style-type: none"> • Energy metabolism • Mitochondrial function • Fatty acid transport 	<p>No evidence a lack of acylcarnitines affects either the IDC schedule or replication rate.</p> <p>Peak during fasting when parasites are in their least energetically/ metabolically demanding stages.</p>	<p>Very unlikely.</p>
Glycerophospholipids <ul style="list-style-type: none"> • Cellular signalling and trafficking • Membrane neogenesis • Haemozoin formation 	<p>Phospholipid composition of the host RBC increases during infection, especially oleic acid (18:1) (Déchamps <i>et al.</i>, 2010).</p> <p>Parasites scavenge fatty acids and also lysoPCs from host plasma, which compete with each other as a source of the acyl components required for lipid synthesis. Lysophosphatidylcholine a C18:1 (lysoPC a 18:1), which contains an oleic acid side chain, is associated with IDC progression.</p> <p>Majority peak during host fasting when parasites are in their least energetically/ metabolically demanding stages. Thus, to be a time-of day cue</p>	<p>Unlikely. Not essential; unlike the liver stage, IDC stages can synthesise fatty acids de novo via type II FA synthase (FASII), (Iarun <i>et al.</i>, 2009).</p>
Biogenic amines and amino acids: <ul style="list-style-type: none"> • Nucleic acid and protein synthesis 	<p>Parasites must scavenge several amino acids from the host, including six host-essential' (isoleucine, leucine, methionine, phenylalanine, threonine and valine) and one host-'non-essential' (alanine) amino acid (Payne & Loomis, 2006) that are rhythmic.</p> <p>Parasites can scavenge most amino acids from catabolism of host haemoglobin (Babbitt <i>et al.</i>, 2012; Liu <i>et al.</i>, 2006; Martin & Kirk, 2007). Isoleucine is the only amino acid absent from human haemoglobin and is one of the least abundant amino acids in rodent haemoglobin (1-3%, Figure 8) yet makes up 9% of both <i>P. falciparum</i>'s and <i>P. chabaudi</i>'s amino acids (Yadav & Swati, 2012).</p> <p>The response of <i>P. falciparum</i> to isoleucine withdrawal is not influenced by whether they were previously cultured in a high or low isoleucine concentration environment and results in dormancy that can be broken upon isoleucine replenishment (Babbitt <i>et al.</i>, 2012).</p> <p>Coinciding with the rise in isoleucine concentration during the feeding window, trophozoites transition to schizonts before bursting and beginning development as ring stages at the end of the feeding window (Figure 5, Figure 7).</p>	<p>Likely. Isoleucine is the only essential amino acid for <i>P. falciparum</i>, it cannot be stored, and dormancy has no apparent ill-effects on subsequent development. <i>P. falciparum</i> deprived of other amino acids develop as normal (Liu <i>et al.</i>, 2006).</p>

Figure 5. Isoleucine is rhythmic and coincides with the intraerythrocytic development cycle (IDC) schedule and host feeding-fasting cycles. Model fit (best fit line and 95% prediction interval) for DF, LF and TRF infections combined, from the time since feeding commences, with feeding-fasting windows overlaid.

how malaria parasites schedule their IDC in line with host feeding-fasting rhythms. A key challenge was differentiating to what extent the absence of isoleucine creates a non-permissive environment, imposing a constraint on parasites that simply forces IDC progression to stop, versus parasites responding in a strategic manner and ‘deciding’ to stop development (Figure 7A). There are several reasons why our findings are not a case of revealing the obvious, that parasites cannot develop when resource-limited. First, when deprived of other essential resources such as glucose, *P. falciparum* rapidly displays stress responses at the transcriptional level yet when deprived of isoleucine, its normal transcriptional pattern persists, albeit very slowly (Babbitt *et al.*, 2012), suggesting it is waiting for a signal to proceed (a “gate”). Second, parasites cannot cope with deprivation of other essential resources, for example dying within hours of glucose deprivation (Babbitt *et al.*, 2012), yet the IDC restarts with no apparent ill-effects when isoleucine is replenished (Figure 6). Third, that parasites cannot create isoleucine stores, nor generate it endogenously, and mount rapid transcriptional responses to changes in exogenous isoleucine concentration (Babbitt *et al.*, 2012), are all hallmarks of an informative cue and a sensitive response system. Thus, we propose that either parasites are so well adapted to isoleucine starvation they cope just fine without it,

Figure 6. Isoleucine provision and withdrawal drives intraerythrocytic development cycle (IDC) completion but does affect parasite mortality. (A) IDC completion defined as the proportion of parasites that are schizonts, in cultures with (orange triangles, Ile +, 50 mg/L) or without (green circles, Ile -) isoleucine. (B) Density of all parasite stages when parasites are cultured with or without isoleucine. Density of (C) schizonts and (D) all parasite stages, after the addition of isoleucine into cultures following isoleucine deprivation for seven (green circles), 14 (orange triangles) and 18 hours (purple squares). The proportion of schizonts in the blood seeding the cultures was ~0.005.

Table 8. Degrees of Freedom (df), log-Likelihood (logLik), AICc, Δ AICc ($AICc_i - AICc_{min}$) and AICc w (AICc weight) for each linear model in the schizont/parasite proportion/density analysis ordered in descending fit (best-fitting model at the top). The response variable for each model is either schizont proportion (Schizont.prop), parasite density (Parasite.dens) or schizont density (Schizont.dens) and the random effect is "mouseID". "Treatment" refers to the treatment group (DF, LF, TRF or ALF), "hours in culture" refers to the number of hours spent in culture since being extracted from the mice, and "hours since isoleucine" refers to the number of hours since isoleucine was added to the cultures. Treatment, hours in culture and hours since isoleucine were all fitted as factors. DF=dark fed wild type mice, LF=light fed wild type mice, TRF=time restricted fed Per1/2-null mice, ALF=ad libitum fed Per1/2-null mice.

Model description: A) Schizont.prop ~ + (1 mouseID)	df	logLik	AICc	Δ AICc	AICcw
treatment	4	106.26	-203.8	0.00	0.996
treatment + hours in culture	7	104.29	-192.6	11.26	0.004
treatment * hours in culture	10	98.81	-173.5	30.37	0.000
null	3	80.96	-155.5	48.31	0.000
hours in culture	6	74.89	-136.3	67.54	0.000
B) Parasite.dens ~ + (1 mouseID)					
null	3	0.14	6.6	0.00	0.836
treatment	4	-0.63	10.7	4.16	0.104
hours in culture	4	-1.30	12.1	5.50	0.053
treatment + hours in culture	5	-2.07	16.4	9.88	0.006
treatment * hours in culture	6	-2.79	20.9	14.36	0.001
C) Schizont.dens ~ + (1 mouseID)					
treatment + hours since isoleucine	6	69.67	-125.3	0.00	0.887
hours since isoleucine	4	65.04	-121.2	4.14	0.112
treatment * hours since isoleucine	8	65.67	-111.6	13.65	0.001
null	3	56.87	-107.2	18.10	0.000
treatment	5	56.67	-101.9	23.39	0.000
D) Parasite.dens ~ + (1 mouseID)					
hours since isoleucine	4	12.96	-17.0	0.00	0.568
null	3	11.49	-16.4	0.56	0.430
treatment + hours since isoleucine	6	9.10	-4.1	12.85	0.001
treatment	5	7.76	-4.1	12.90	0.001
treatment * hours since isoleucine	8	10.47	-1.2	15.75	0.000

unlike their ability to cope with other forms of starvation, or that isoleucine exerts its effects on IDC progression as both a time-cue and resource (Figure 7B).

Most studies of isoleucine uptake and use in malaria parasites focus on *P. falciparum*. This parasite uses several channels and

receptors (both parasite- and host-derived) to acquire resources from the host. Uninfected human RBC take up both isoleucine and methionine via the saturable L-system (Cobbold *et al.*, 2011), which supplies 20% of the necessary isoleucine (Martin & Kirk, 2007). When parasitised, there is a five-fold increase in isoleucine entering RBC which is attributable to

Figure 7. Schematic for isoleucine's role in the intraerythrocytic development cycle (IDC) schedule. (A) In the absence of isoleucine (Ile), but in the presence of all other essential components of culture media, *P. chabaudi*'s IDC progression stops or continues very slowly, as observed for *P. falciparum* (Liu *et al.*, 2006 and Babbitt *et al.*, 2012). This observation is consistent with isoleucine being an essential resource and/or a time-cue. This means the time-of-day cue may also be a direct selective agent (i.e. the reason why a timing strategy has evolved). In such situations it is very difficult to unravel the consequences of cue loss. Here, the challenge is differentiating to what extent the absence of isoleucine (i) creates a non-permissive environment, imposing a constraint on parasites that simply forces IDC progression to stop, versus (ii) parasites responding in a strategic manner and 'deciding' to stop development. Under scenario (i) parasites experience resource limitation, and starvation by definition should have negative consequences. For example, the longer parasites are starved, the more die, or experience increasingly lower or slower rates of recovery upon isoleucine replenishment. Under scenario (ii) there are no such costs because parasites avoid starvation by stopping/slows development and waiting for a signal that it is time to expect resources to be replenished. That parasite number is not affected by isoleucine withdrawal, even for prolonged periods, and that parasites recover IDC completion at the same rate regardless of the duration of withdrawal, is consistent with scenario (ii) not scenario (i). **(B)** It is now well established that *P. chabaudi*'s IDC schedule is synchronised to host feeding-fasting cycles. Our findings suggest this coordination is achieved by parasites responding to daily rhythmicity in the blood concentration of isoleucine for the purpose of maximising exploitation of the host's isoleucine, and potentially other nutrients, they can more easily acquire from the host's digestion of its food than by biosynthesis or scavenging from haemoglobin.

Figure 8. Frequency of amino acids in alpha (A) and beta (B) chains of human (orange) and mouse (blue) haemoglobin. Amino acid codes: A - alanine, C - cysteine, D - aspartic acid, E - glutamic acid, F - phenylalanine, G - glycine, H - histidine, I - isoleucine, K - lysine, L - leucine, M - methionine, N - asparagine, P - proline, Q - glutamine, R - arginine, S - serine, T - threonine, V - valine, W - tryptophan, Y - tyrosine.

new permeability pathways (NPPs) introduced into the RBC membrane by the parasite (Martin & Kirk, 2007; McLean & Jacobs-Lorena, 2020). NPPs supply 80% of the necessary isoleucine (Martin & Kirk, 2007) and are active only in the host membrane at the trophozoite and schizont stages of the IDC, suggesting this influx of isoleucine occurs only at certain times of day (e.g. after host feeding) (Kutner *et al.*, 1985). Assuming *P. chabaudi* has analogous mechanisms, we propose that elevated isoleucine is used by the parasite as a marker for a sufficiently nutrient-rich environment to traverse cell cycle checkpoints and complete the IDC (McLean & Jacobs-Lorena, 2020; O'Neill *et al.*, 2020).

Glucose has previously been suggested as a time-cue or scheduling force for the IDC (Hirako *et al.*, 2018; Prior *et al.*, 2018). Our experiment - which is the first to quantify glucose and IDC rhythms across different feeding-fasting rhythms - does not support a direct effect of glucose but it may be indirectly involved. Parasites that are glucose-limited fail to concentrate isoleucine (Martin & Kirk, 2007), likely due to a lack of glycolysis and ATP production needed to operate isoleucine transporters. High concentrations of isoleucine in the blood are also associated with uptake of glucose by tissues, potentially contributing to the hypoglycaemia associated with TNF-stimulation of immune cells during malaria infection (Elased & Playfair, 1994; Hirako *et al.*, 2018). Additionally, rodent models and humans with obesity and type 2 diabetes like pathologies have elevated levels of isoleucine and dampened glucose rhythms in the blood (Lynch & Adams, 2014;

Isherwood *et al.*, 2017). Thus, if glucose limitation or elevation interferes with the parasite's ability to acquire time-of-day information from isoleucine, the IDC schedule will be disrupted, as observed in diabetic mice (Hirako *et al.*, 2018). Connections between isoleucine and glucose might also explain why the parasite protein kinase 'KIN' is involved in nutrient sensing (Mancio-Silva *et al.*, 2017).

Given that daily variation in the concentration of isoleucine in the blood of infected mice appears modest (55 μ M to 80 μ M from nadir to peak; Figure 5), we suggest that like *P. falciparum*, *P. chabaudi* is very sensitive to changes in isoleucine levels. This observation could be interpreted in two non-mutually exclusive ways. Perhaps this seemingly damp rhythm means that sufficient isoleucine is available around-the-clock to meet the parasites resource-needs (from the blood and scavenging from the very low level (<3%) of isoleucine in rodent haemoglobin; Figure 8), so isoleucine acts on the IDC schedule only as a time-cue (to align with resources that are limited at certain times of day) rather than as a developmental-rate limiting resource as well. For instance, the appearance of isoleucine may signal a window during which vitamins, cofactors, purines, folic acid, pantothenic acid, and glucose, that are also required for successful replication are available (Müller & Kappes, 2007; Müller *et al.*, 2010; Sherman, 1979). Furthermore, isoleucine is also reliable cue for the timing of other nutrients that are perhaps easier for parasites to acquire from the host's digestion of food than from biosynthesis or scavenging from haemoglobin. In particular, that the expression of genes associated with

Figure 9. Concentration of all metabolites at all time points during the time series, those included and excluded from the analysis. DF=dark fed wild type mice, LF=light fed wild type mice, TRF=time restricted fed Per1/2-null mice, ALF=ad libitum fed Per1/2-null mice. Each x-axis mark is a different metabolite with concentration (μM) on the y-axis. In black are metabolites excluded since they failed set LC/MS assay criteria and those metabolites taken forward into the analysis are red. Metabolites are plotted in alphabetical order: Ala, Arg, Asn, Asp, Cit, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Orn, Phe, Pro, Ser, Thr, Trp, Tyr, Val, Ac.Orn, ADMA, alpha.AAA, c4.OH.Pro, Carnosine, Creatinine, DOPA, Dopamine, Histamine, Kynurenine, Met.SO, Nitro.Tyr, PEA, Putrescine, Sarcosine, SDMA, Serotonin, Spermidine, Spermine, t4.OH.Pro, Taurine, C0, C2, C3, C3.DC.C4.OH., C3.OH, C3.1, C4, C4.1, C5, C5.DC..C6.OH., C5.M.DC, C5.OH..C3.DC.M., C5.1, C5.1.DC, C6.C4.1.DC., C6.1, C7.DC, C8, C9, C10, C10.1, C10.2, C12, C12.DC, C12.1, C14, C14.1, C14.1.OH, C14.2, C14.2.OH, C16, C16.OH, C16.1, C16.1.OH, C16.2, C16.2.OH, C18, C18.1, C18.1.OH, C18.2, lysoPC.a.C14.0, lysoPC.a.C16.0, lysoPC.a.C16.1, lysoPC.a.C17.0, lysoPC.a.C18.0, lysoPC.a.C18.1, lysoPC.a.C18.2, lysoPC.a.C20.3, lysoPC.a.C20.4, lysoPC.a.C24.0, lysoPC.a.C26.0, lysoPC.a.C26.1, lysoPC.a.C28.0, lysoPC.a.C28.1, PC.aa.C24.0, PC.aa.C26.0, PC.aa.C28.1, PC.aa.C30.0, PC.aa.C32.0, PC.aa.C32.1, PC.aa.C32.2, PC.aa.C32.3, PC.aa.C34.1, PC.aa.C34.2, PC.aa.C34.3, PC.aa.C34.4, PC.aa.C36.0, PC.aa.C36.1, PC.aa.C36.2, PC.aa.C36.3, PC.aa.C36.4, PC.aa.C36.5, PC.aa.C36.6, PC.aa.C38.0, PC.aa.C38.3, PC.aa.C38.4, PC.aa.C38.5, PC.aa.C38.6, PC.aa.C40.1, PC.aa.C40.2, PC.aa.C40.3, PC.aa.C40.4, PC.aa.C40.5, PC.aa.C40.6, PC.aa.C42.0, PC.aa.C42.1, PC.aa.C42.2, PC.aa.C42.4, PC.aa.C42.5, PC.aa.C42.6, PC.aa.C30.0, PC.aa.C30.1, PC.aa.C30.2, PC.aa.C32.1, PC.aa.C32.2, PC.aa.C34.0, PC.aa.C34.1, PC.aa.C34.2, PC.aa.C34.3, PC.aa.C36.0, PC.aa.C36.1, PC.aa.C36.2, PC.aa.C36.3, PC.aa.C36.4, PC.aa.C36.5, PC.aa.C36.6, PC.aa.C38.0, PC.aa.C38.1, PC.aa.C38.2, PC.aa.C38.3, PC.aa.C38.4, PC.aa.C38.5, PC.aa.C38.6, PC.aa.C40.1, PC.aa.C40.2, PC.aa.C40.3, PC.aa.C40.4, PC.aa.C40.5, PC.aa.C40.6, PC.aa.C42.0, PC.aa.C42.1, PC.aa.C42.2, PC.aa.C42.3, PC.aa.C42.4, PC.aa.C42.5, PC.aa.C44.3, PC.aa.C44.4, PC.aa.C44.5, PC.aa.C44.6, SM.OH, C14.1, SM.OH.C16.1, SM.OH.C22.1, SM.OH.C22.2, SM.OH.C24.1, SM.C16.0, SM.C16.1, SM.C18.0, SM.C18.1, SM.C20.2, SM.C24.0, SM.C24.1, SM.C26.0, SM.C26.1.

translation are the most commonly disrupted when *P. chabaudi*'s rhythms are perturbed away from the host's rhythms (Subhudi *et al.*, 2020) suggests parasites align the IDC schedule with

the resources required for proteins. However, if isoleucine is not a limiting resource at certain times of day, this may only be the case early infections.

Like blood glucose concentration, the isoleucine rhythm may become exacerbated as infections develop high parasite burdens and hosts become sick. In this case, by aligning the IDC schedule correctly early in infections, parasites minimise the costs of resource limitation later on.

Our findings offer a route into identifying the molecular pathways involved in transducing environmental signals to the IDC schedule. To date, the only gene known to be involved in the IDC schedule is SR10, which modulates IDC duration in response to perturbations of host time-of-day (Subudhi *et al.*, 2020). Identifying KIN (Mancio-Silva *et al.*, 2017), regulated pathways whether they, along with pathways associated with SR10 (Subudhi *et al.*, 2020), and PK7/MORC (which is involved in IDC stage transitions; Singh *et al.*, 2021), are sensitive to isoleucine might reveal how the parasites' time-keeping mechanism operates. Whilst by no means conclusive, our results feed the debate about whether malaria parasites keep time by an endogenous circadian oscillator or a simpler mechanism. The hallmarks of an endogenous circadian clock are (i) temperature compensation, (ii) free running in constant conditions, and (iii) entrainment to a Zeitgeber (Pittendrigh, 1960). Recent observations are consistent with (ii) (Rijo-Ferreira *et al.*, 2020; Smith *et al.*, 2020; Subudhi *et al.*, 2020) and our results now allow entrainment to isoleucine to be tested for (iii), as well as free-running in isoleucine-constant conditions (ii). However, an endogenous clock may exist to allow parasites to schedule activities other than the IDC. Overall, the rapid stop-start responses to isoleucine withdrawal and replenishment that we observe, and that is known for *P. falciparum* (Babbitt *et al.*, 2012; Liu *et al.*, 2006; McLean & Jacobs-Lorena, 2020), appear more parsimoniously explained by a "just-in-time reactionary strategy" than an endogenous oscillator. This is because clocks generally require several cycles to catch up with a large change in the phase (timing) of their Zeitgeber (which is why jet lag occurs). Thus, in the absence of isoleucine or a phase-shift in its availability, if the ticking of an endogenous clock cannot be readily de-coupled from IDC progression, development would have continued as best as possible, not stopped. Furthermore, most clocks are set by a Zeitgeber (usually light) that is a good proxy for something else (e.g. predation risk) that is the ultimate driver of selection for a timing mechanism. Yet, isoleucine appears to be both cue and selective driver, further suggesting a just-in-time response. This strategy has the advantage of cueing in to the most accurate information about resource availability, and providing a more fast-responding strategy than a clock when the bulk of host foraging could occur anytime within the broad window of the active phase, and whose timing may be disrupted if hosts become severely ill.

However the IDC is scheduled, coordinating development according to the availability of the resources needed to produce progeny intuitively seems like a good strategy to maximise fitness. Yet, the costs/benefits of the IDC schedule demonstrated by parasites may be mediated by parasite density. At low parasite densities (e.g. at the start of infection), resources may be sufficient to support IDC completion of all parasites at any time-of-day, but at intermediate densities, parasites may need

to align their IDC needs with rhythms in resource availability. Finally, at very high densities and/or when hosts become sick, resources could be limiting so a synchronous IDC leads to deleterious competition between related parasites. Quantifying whether the costs and benefits of a synchronous IDC vary during infections in line with parasite density and resource availability is complicated by the connection between the IDC schedule and the timing of transmission stages. By aligning the IDC schedule with the feeding-fasting rhythm which is set by light-dark rhythms in the environment, parasites benefit from the knock-on alignment of transmission stage maturation with vector biting activity (O'Donnell *et al.*, 2011; Pigeault *et al.*, 2018; Schneider *et al.*, 2018). Therefore, a key question is to what extent does the IDC schedule contribute to parasite fitness via transmission benefits versus variation during infections in the benefits and costs of aligning asexual replication with rhythmic resources in the blood? More broadly, by integrating evolutionary and mechanistic insight, it may be possible to improve antimalarial treatments. For example, impairing the parasites ability to detect or respond to isoleucine may stall the IDC, reducing virulence and buy time for host immune responses to clear the infection, as well as preventing ring stage dormancy from providing tolerance to antimalarials (Codd *et al.*, 2011).

Methods

Ethics statement

All procedures complied with the UK Home Office regulations (Animals Scientific Procedures Act 1986; project 483 licence number 70/8546) and were approved by the University of Edinburgh. All efforts were made to ameliorate harm to animals through monitoring of animal health by twice daily visual inspections and daily routine health checks, such as weight and RBC count.

Experimental designs and terminology

The same four perturbations of host and parasite rhythms were used in the metabolomics experiment and the glucose monitoring experiment. The *Per1/2*-null mice (non-functional proteins Period 1 & 2, backcrossed onto a C57BL/6 background for over 10 generations) were donated by Michael Hastings (MRC Laboratory of Molecular Biology, Cambridge, UK) and generated by David Weaver (UMass Medical School, Massachusetts, USA). Wild type C57BL/6 mice were housed in a 12h light: 12h dark regime (12 hours of light followed by 12 hours of darkness) and the *Per1/2*-null mice housed in constant darkness (DD) for two weeks prior to the start of infections and throughout sampling. We randomly separated mice into their treatment groups. We refer to time-of-day using Zeitgeber Time (ZT) for mice housed under entrained conditions (light: dark cycles) and hours when *Per1/2*-null mice are housed under constant conditions (dark:dark). WT mice either had access to food at night (dark feeding, DF) or in the day (light feeding, LF). *Per1/2*-null mice either had access to food for 12 hours (time restricted feeding, TRF) or constant access to food (*ad libitum* feeding, ALF). Every 12 hours, food was added/removed accordingly from the DF, LF and TRF cages and the cages were checked for evidence of hoarding, which

was never observed. ALF cages were also disturbed during food removal/provision of the other groups. Investigators could not be blinded to the mouse strain due the difference in coat colour, or the lighting and feeding regime due to the need to add and remove food manually.

Sampling. All mice were infected intravenously with *P. chabaudi* DK genotype by injection of 1×10^5 infected RBC at ring stage. Sampling started on day 5 post infection and occurred every two hours for both the metabolomics and glucose experiments. We made a thin blood smear each time a mouse was sampled to quantify parasite rhythms by counting ~100 parasites per blood smear using microscopy. Smears were read without knowledge of the treatment group, and excluded where smear quality was too poor to count RBCs. Following [Prior et al. \(2018\)](#); [O'Donnell et al. \(2020\)](#) and [Rijo-Ferreira et al. \(2020\)](#), we used the proportion of ring stages as a phase marker (an estimate of the timing of parasite development in the blood) of parasite rhythms. We calculated amplitude and time-of-day of peak for each treatment group using sine and cosine curves in a linear model to confirm the IDC schedules for each group as used in [O'Donnell et al. \(2020\)](#).

Metabolomics experiment. We infected 68 eight-week-old female mice: 35 C57BL/6 wild type animals (DF and LF mice) and 33 *Per1/2*-null TTFL clock-disrupted mice (TRF and ALF) ([Table 2](#)). Sample sizes were chosen based on previous data collected from our lab using our study system. We sampled mice in blocks (A-D), meaning each individual mouse was sampled every eight hours during the 26-hour sampling window, with 14 time points in total. We did not sample each mouse at each sampling point to minimise the total volume of blood being taken. At each sampling point for each designated host, 20 μ l blood was taken from the tail vein to provide 10 μ l blood plasma for snap freezing using dry ice.

Glucose experiment. We infected 20 eight-week-old male mice: 10 C57BL/6 wild type animals and 10 *Per1/2*-null circadian clock-disrupted mice (as described above, [Table 2](#)). Sample sizes were chosen based on previous data collected using our study system. We recorded blood glucose concentration from all mice every two hours by taking 1 μ l blood using an [Accu-Chek Performa Nano glucometer](#).

Targeted metabolomics

We quantified metabolites by analysing plasma samples using the AbsoluteIDQ p180 targeted metabolomics kit (Biocrates Life Sciences AG, Innsbruck, Austria) and a Waters Xevo TQ-S mass spectrometer coupled to an Acquity UPLC system (Waters Corporation, Milford, MA, USA) ([Isherwood et al., 2017](#); [Skene et al., 2018](#)). We prepared the plasma samples (10 μ l) according to the manufacturer's instructions, adding several stable isotope-labelled standards to the samples prior to the derivatization and extraction steps. Using UPLC/MS (ultra performance liquid chromatography/mass spectrometry), we quantified 185 metabolites from five different compound classes (acylcarnitines, amino acids, biogenic amines, glycerophospholipids, and sphingolipids). We ran the samples on two 96-well plates, randomised the sample order and ran three levels of

quality control (QC) on each plate. We normalised the data between the plates using the results of quality control level 2 (QC2) repeats across the plate ($n=4$) and between plates ($n=2$) using Biocrates METIDQ software (QC2 correction). Metabolites were excluded if the CV% of QC2 was $>30\%$ or if all four groups contained $>25\%$ of samples that were below the limit of detection, below the lower limit of quantification, or above the limit of quantification or blank out of range. The remaining 134 quantified metabolites comprised of seven acylcarnitines, 19 amino acids, 15 biogenic amines, 79 glycerophospholipids and 14 sphingolipids (see [Figure 9](#)).

Isoleucine response experiments

To test the effect of the amino acid isoleucine on the IDC, we compared parasite developmental progression in cultures with and without isoleucine (50 mg/L), as well as after different durations (seven, 14, 18 hours) of isoleucine starvation. We set up $N = 112$ cultures (from eight mice) so that for each time point within each treatment, an independent culture was sampled, avoiding any bias associated with repeat-sampling individual cultures. We used eight-week-old wild type female mice, MF1 strain, housed in a 12h:12h light:dark regime before and during infection. We infected mice intraperitoneally with 1×10^6 *P. chabaudi* genotype DK infected red blood cells at ring stage and terminally bled them on day 6 post infection, when the parasitaemia was around 15%. Mice were bled at ZT4, when parasites were late rings/ early trophozoites (see [Prior et al., 2018](#)). Approximately 1 ml of blood was collected from each mouse, which was then split equally across cultures in all five treatment groups.

Culturing. We washed infected blood twice with buffered RPMI containing no amino acids (following [Spence et al., 2011](#)), before being reconstituted in the RPMI medium corresponding to each treatment (containing isoleucine, or not) ([dx.doi.org/10.17504/protocols.io.buuqnwvw](#)). We cultured parasites in 96-well round bottom plates with total culture volumes of 200–250 μ l, at ~3% haematocrit and kept the culture plates inside a gas chamber which was gassed upon closing with 88% nitrogen 7% carbon dioxide and 5% oxygen, and then placed inside a 37°C incubator. The culture medium was custom made RPMI from [Cell Culture Technologies](#), Switzerland.

Sampling. We sampled parasites in the first experiment (comparing IDC completion in isoleucine rich versus isoleucine free media) at 13–14, 16–18, 20–21, 24–26 and 27 hours after culture initiation. We also sampled parasites that had been isoleucine deprived for seven, 14 or 18 hours at six and nine hours after isoleucine addition. Samples consisted of a thin blood smear from each culture fixed with methanol and Giemsa stained. We measured the proportion of parasites in the schizont stage (as an indicator of parasites having completed their IDC) by counting ~300 parasites per blood smear. Smears were read without knowledge of the treatment group, and excluded where smear quality was too poor to count RBCs.

Statistical analysis

We defined daily rhythmicity in the concentrations of the 134 metabolites detected by the UPLC/MS-MS platform as

the detection of rhythmicity in at least two of the following circadian analysis programmes: **ECHO**, **CircWave** and **JTK_Cycle**. For metabolites not found to be rhythmic by at least two of the circadian analysis programmes, we also carried out Analysis of Variance to identify metabolites that varied across the day but without a detectable 24-hour rhythm (Table 3 for breakdown of which candidate metabolites were rhythmic in each programme). To calculate the acrophase (timing of peak), we used linear mixed-effects regression models containing sine and cosine terms on all metabolites that varied across the day. Metabolites whose acrophase fell in the same 12-hour feeding or fasting window (ZT0-12 or ZT12-24) in LF and TRF infections but fell in the opposite window for the DF group were shortlisted as potential candidates to connect host feeding rhythms and the IDC schedule. We compared schizont proportion and the density of combined IDC stages using linear regression models. To avoid overfitting due to small sample sizes, “Akaike information criterion-corrected” (AICc) values were calculated for each model, and a change in two AICc ($\Delta\text{AICc}=2$) was chosen to select the most parsimonious model (Brewer *et al.*, 2016).

An earlier version of this article can be found on bioRxiv (doi: <https://doi.org/10.1101/2020.08.24.264689>).

Data availability

Underlying data

Figshare: Plasmodium chabaudi genotype DK parasite stages during time series. <https://doi.org/10.6084/m9.figshare.14842695.v1> (Prior *et al.*, 2021a).

Figshare: Targeted metabolomics on malaria-infected mouse blood. <https://doi.org/10.6084/m9.figshare.14842638.v1> (Prior *et al.*, 2021b).

Figshare: Plasmodium schizont proportion in culture grown with and without isoleucine. <https://doi.org/10.6084/m9.figshare.14842713.v1> (Prior *et al.*, 2021c).

Figshare: Plasmodium schizont densities in culture grown with and without isoleucine, as well as isoleucine starvation for different durations of time. <https://doi.org/10.6084/m9.figshare.14842716.v1> (Prior *et al.*, 2021d).

Figshare: Blood glucose concentration time series in mice infected with malaria parasites <https://doi.org/10.6084/m9.figshare.14912709.v1> (Prior *et al.*, 2021e).

Reporting guidelines

Figshare: ARRIVE checklist for “Synchrony between daily rhythms of malaria parasites and hosts is driven by an essential amino acid”. <https://doi.org/10.6084/m9.figshare.14912703.v1> (Prior *et al.*, 2021f).

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Acknowledgements

We thank Daan van der Veen, Sam Rund, Petra Schneider, Alejandra Herbert-Mainero and Ronnie Mooney for practical help and discussion.

References

- Babbitt SE, Altenhofen L, Cobbold SA, *et al.*: **Plasmodium falciparum** responds to amino acid starvation by entering into a hibernatory state. *Proc Natl Acad Sci U S A*. 2012; **109**(47): E3278–E3287.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bae K, Jin X, Maywood ES, *et al.*: Differential functions of *mPer1*, *mPer2*, and *mPer3* in the SCN circadian clock. *Neuron*. 2001; **30**(2): 525–536.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Brewer MJ, Butler A, Cooksley SL: The relative performance of AIC, AICc, and BIC in the presence of unobserved heterogeneity. *Methods Ecol Evol*. 2016; **7**(6): 679–692.
[Publisher Full Text](#)
- Butcher GA, Cohen S: Short-term culture of *Plasmodium knowlesi*. *Parasitology*. 1971; **62**(2): 309–320.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cobbold SA, Martin RE, Kirk K: Methionine transport in the malaria parasite *Plasmodium falciparum*. *Int J Parasitol*. 2011; **41**(1): 125–135.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Codd A, Teuscher F, Kyle DE, *et al.*: Artemisinin-induced parasite dormancy: a plausible mechanism for treatment failure. *Malar J*. 2011; **10**: 56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Daily JP, Scanfeld D, Pochet N, *et al.*: Distinct physiological states of *Plasmodium falciparum* in malaria-infected patients. *Nature*. 2007; **450**(7172): 1091–1095.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Déchamps S, Shastri S, Wengelnik K, *et al.*: Glycerophospholipid acquisition in *Plasmodium* - a puzzling assembly of biosynthetic pathways. *Int J Parasitol*. 2010; **40**(12): 1347–1365.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Elased K, Playfair JH: Hypoglycemia and hyperinsulinemia in rodent models of severe malaria infection. *Infect Immun*. 1994; **62**(11): 5157–5160.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Garcia CR, Markus RP, Madeira L: Tertian and quartan fevers: temporal regulation in malarial infection. *J Biol Rhythms*. 2001; **16**(5): 436–443.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hevia MA, Canessa P, Müller-Esparza H, *et al.*: A circadian oscillator in the fungus *Botrytis cinerea* regulates virulence when infecting *Arabidopsis thaliana*. *Proc Natl Acad Sci USA*. 2015; **112**(28): 8744–8749.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hirako IC, Assis PA, Hojo-Souza NS, *et al.*: Daily Rhythms of TNF α Expression and Food Intake Regulate Synchrony of *Plasmodium* Stages with the Host Circadian Cycle. *Cell Host Microbe*. 2018; **23**(6): 796–808.e6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hozer C, Perret M, Parvard S, *et al.*: Survival is reduced when endogenous period deviates from 24 h in a non-human primate, supporting the circadian resonance theory. *Sci Rep*. 2020; **10**(1): 18002.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Isherwood CM, van der Veen DR, Johnston JD, *et al.*: Twenty-four-hour rhythmicity of circulating metabolites: effect of body mass and type 2 diabetes. *FASEB J*. 2017; **31**(12): 5557–5567.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kutner S, Breuer WV, Ginsburg H, *et al.*: Characterization of permeation pathways in the plasma membrane of human erythrocytes infected with early stages of *Plasmodium falciparum*: association with parasite development. *J Cell Physiol*. 1985; **125**(3): 521–527.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Liu J, Istvan ES, Gluzman IY, *et al.*: *Plasmodium falciparum* ensures its amino

acid supply with multiple acquisition pathways and redundant proteolytic enzyme systems. *Proc Natl Acad Sci U S A*. 2006; **103**(23): 8840–8845.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Lynch CJ, Adams SH: **Branched-chain amino acids in metabolic signalling and insulin resistance.** *Nat Rev Endocrinol*. 2014; **10**(12): 723–736.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Mancio-Silva L, Slavic K, Ruivo MTG, et al.: **Nutrient sensing modulates malaria parasite virulence.** *Nature*. 2017; **547**(7662): 213–216.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Martin RE, Kirk K: **Transport of the essential nutrient isoleucine in human erythrocytes infected with the malaria parasite *Plasmodium falciparum*.** *Blood*. 2007; **109**(5): 2217–2224.

[PubMed Abstract](#) | [Publisher Full Text](#)

Martinez-Bakker M, Helm B: **The influence of biological rhythms on host-parasite interactions.** *Trends Ecol Evol*. 2015; **30**(6): 314–26.

[PubMed Abstract](#) | [Publisher Full Text](#)

Maywood ES, Chesham JE, Smyllie NJ, et al.: **The *Tau* mutation of casein kinase 1ε sets the period of the mammalian pacemaker via regulation of Period1 or Period2 clock proteins.** *J Biol Rhythms*. 2014; **29**(2): 110–118.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

McLean KJ, Jacobs-Lorena M: **The response of *Plasmodium falciparum* to isoleucine withdrawal is dependent on the stage of progression through the intraerythrocytic cell cycle.** *Malar J*. 2020; **19**(1): 147.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Mideo N, Reece SE, Smith AL, et al.: **The Cinderella syndrome: why do malaria-infected cells burst at midnight?** *Trends Parasitol*. 2013; **29**(1): 10–16.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Müller IB, Hyde JE, Wrenger C: **Vitamin B metabolism in *Plasmodium falciparum* as a source of drug targets.** *Trends Parasitol*. 2010; **26**(1): 35–43.

[PubMed Abstract](#) | [Publisher Full Text](#)

Müller S, Kappes B: **Vitamin and cofactor biosynthesis pathways in *Plasmodium* and other apicomplexan parasites.** *Trends Parasitol*. 2007; **23**(3): 112–121.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

O'Donnell AJ, Mideo N, Reece SE: **Disrupting rhythms in *Plasmodium chabaudi*: costs accrue quickly and independently of how infections are initiated.** *Malar J*. 2013; **12**: 372.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

O'Donnell AJ, Prior KF, Reece SE: **Host circadian clocks do not set the schedule for the within-host replication of malaria parasites.** *Proc Biol Sci*. 2020; **287**(1932): 20200347.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

O'Donnell AJ, Schneider P, McWatters HG, et al.: **Fitness costs of disrupting circadian rhythms in malaria parasites.** *Proc Biol Sci*. 2011; **278**(1717): 2429–2436.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

O'Neill JS, Hoyle NP, Robertson JB, et al.: **Eukaryotic cell biology is temporally coordinated to support the energetic demands of protein homeostasis.** *Nat Commun*. 2020; **11**(1): 4706.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Oliszewski KL, Morrissey JM, Wilinski D, et al.: **Host-parasite interactions revealed by *Plasmodium falciparum* metabolomics.** *Cell Host Microbe*. 2009; **5**(2): 191–199.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Ouyang Y, Andersson CR, Kondo T, et al.: **Resonating circadian clocks enhance fitness in cyanobacteria.** *Proc Natl Acad Sci USA*. 1998; **95**(15): 8660–8664.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Payne SH, Loomis WF: **Retention and loss of amino acid biosynthetic pathways based on analysis of whole-genome sequences.** *Eukaryotic Cell*. 2006; **5**(2): 272–276.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Pigeault R, Caudron Q, Nicot A, et al.: **Timing malaria transmission with mosquito fluctuations.** *Evol Lett*. 2018; **2**(4): 378–389.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Pittendrigh CS: **Circadian rhythms and the circadian organization of living systems.** *Cold Spring Harb Symp Quant Biol*. 1960; **25**: 159–184.

[PubMed Abstract](#) | [Publisher Full Text](#)

Polet H, Conrad ME: **Malaria: extracellular amino acid requirements for *in vitro* growth of erythrocytic forms of *Plasmodium knowlesi*.** *Proc Soc Exp Biol Med*. 1968; **127**(1): 251–253.

[PubMed Abstract](#) | [Publisher Full Text](#)

Polet H, Conrad ME: **The influence of three analogs of isoleucine on *in vitro* growth and protein synthesis of erythrocytic forms of *Plasmodium knowlesi*.** *Proc Soc Exp Biol Med*. 1969; **130**(2): 581–586.

[PubMed Abstract](#) | [Publisher Full Text](#)

Prior K, Middleton B, Owolabi A, et al.: ***Plasmodium chabaudi* genotype DK**

parasite stages during time series. *figshare*. Dataset. 2021a.

<http://www.doi.org/10.6084/m9.figshare.14842695.v1>

Prior K, Middleton B, Owolabi A, et al.: **Targeted metabolomics on malaria-infected mouse blood.** *figshare*. Dataset. 2021b.

<http://www.doi.org/10.6084/m9.figshare.14842638.v1>

Prior K, Middleton B, Owolabi A, et al.: ***Plasmodium* schizont proportion in culture grown with and without isoleucine.** *figshare*. Dataset. 2021c.

<http://www.doi.org/10.6084/m9.figshare.14842713.v1>

Prior K, Middleton B, Owolabi A, et al.: ***Plasmodium* schizont densities in culture grown with and without isoleucine, as well as isoleucine starvation for different durations of time.** *figshare*. Dataset. 2021d.

<http://www.doi.org/10.6084/m9.figshare.14912709.v1>

Prior K, Middleton B, Owolabi A, et al.: **Blood glucose concentration time series in mice infected with malaria parasites.** *figshare*. Dataset. 2021e.

<http://www.doi.org/10.6084/m9.figshare.14912709.v1>

Prior K, Middleton B, Owolabi A, et al.: **ARRIVE Checklist.pdf.** *figshare*. Journal contribution. 2021f.

<http://www.doi.org/10.6084/m9.figshare.14912703.v1>

Prior KF, Rijo-Ferreira F, Assis PA, et al.: **Periodic parasites and daily host rhythms.** *Cell Host Microbe*. 2020; **27**(2): 176–187.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Prior KF, van der Veen DR, O'Donnell AJ, et al.: **Timing of host feeding drives rhythms in parasite replication.** *PLoS Pathog*. 2018; **14**(2): e1006900.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Reece SE, Prior KF, Mideo N: **The Life and Times of Parasites: Rhythms in Strategies for Within-host Survival and Between-host Transmission.** *J Biol Rhythms*. 2017; **32**(6): 516–533.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Reinke H, Asher G: **Crosstalk between metabolism and circadian clocks.** *Nat Rev Mol Cell Biol*. 2019; **20**(4): 227–241.

[PubMed Abstract](#) | [Publisher Full Text](#)

Rijo-Ferreira F, Pinto-Neves D, Barbosa-Morais NL, et al.: ***Trypanosoma brucei* metabolism is under circadian control.** *Nat Microbiol*. 2017; **2**: 17032.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Rijo-Ferreira F, Acosta-Rodriguez VA, Abel JH, et al.: **The malaria parasite has an intrinsic clock.** *Science*. 2020; **368**(6492): 746–753.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Schneider P, Rund SSC, Smith NL, et al.: **Adaptive periodicity in the infectivity of malaria gametocytes to mosquitoes.** *Proc Biol Sci*. 2018; **285**(1888): 20181876.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sherman IW: **Biochemistry of *Plasmodium* (malarial parasites).** *Microbiol Rev*. 1979; **43**(4): 453–495.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Singh MK, Tessarin-Almeida G, Dias BKM, et al.: **A nuclear protein, PFMORC confers melatonin dependent synchrony of the human malaria parasite *P. falciparum* in the asexual stage.** *Sci Rep*. 2021; **11**(1): 2057.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Skene DJ, Skorniyakov E, Chowdhury NR, et al.: **Separation of circadian- and behavior-driven metabolite rhythms in humans provides a window on peripheral oscillators and metabolism.** *Proc Natl Acad Sci U S A*. 2018; **115**(30): 7825–7830.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Smith LM, Motta FC, Chopra G, et al.: **An intrinsic oscillator drives the blood stage cycle of the malaria parasite *Plasmodium falciparum*.** *Science*. 2020; **368**(6492): 754–759.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Spence PJ, Cunningham D, Jarra W, et al.: **Transformation of the rodent malaria parasite *Plasmodium chabaudi*.** *Nat Protoc*. 2011; **6**(4): 553–561.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Spoelstra K, Wikelski M, Daan S, et al.: **Natural selection against a circadian clock gene mutation in mice.** *Proc Natl Acad Sci USA*. 2016; **113**(3): 686–691.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Subudhi AK, O'Donnell AJ, Ramaprasad A, et al.: **Malaria parasites regulate intra-erythrocytic development duration via serpentine receptor 10 to coordinate with host rhythms.** *Nat Commun*. 2020; **11**(1): 2763.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tarun AS, Vaughan AM, Kappe SHI: **Redefining the role of de novo fatty acid synthesis in *Plasmodium* parasites.** *Trends Parasitol*. 2009; **25**(12): 545–550.

[PubMed Abstract](#) | [Publisher Full Text](#)

Westwood ML, O'Donnell AJ, de Bekker C, et al.: **The evolutionary ecology of circadian rhythms in infection.** *Nat Ecol Evol*. 2019; **3**(4): 552–560.

[PubMed Abstract](#) | [Publisher Full Text](#)

Yadav MJ, Swati D: **Comparative genome analysis of six malarial parasites using codon usage bias based tools.** *Bioinformation*. 2012; **8**(24): 1230–1239.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 06 October 2021

<https://doi.org/10.21956/wellcomeopenres.18637.r45928>

© 2021 Zhuang X. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Xiaodong Zhuang

Nuffield Department of Medicine, University of Oxford, Oxford, UK

Prior *et al.* performed an interesting study that identified a key nutrient mediator responsible for the synchrony between plasmodium blood-stage replication and the host feeding-fasting rhythm. In this study, circadian entrained wild type and circadian defect *per1/2* knockout mice with different feeding schedules were infected with *P. chabaudi* followed by metabolomics analysis of blood samples at 2h intervals. The data suggest the feeding rhythm alone can override the host clock defect to regulate the rhythmic intraerythrocytic development cycle (IDC) and metabolites. Isoleucine was identified as the 'circadian gate' for parasites to set their time-of-day schedule for replication. Their findings provide important evidence to increase our understanding of how parasites align their lifecycle with the host's daily rhythm. Future mechanistic studies to define the role of isoleucine in regulating rhythmic IDCs are certainly warranted. The manuscript is nicely written and the results clearly demonstrated, which supports their major conclusions.

Specific comments:

- Abstract - Define TTFL.
- Figure 1 - Legend (B) "The experimental...".
- Figure 3 - Please label the top right and bottom left boxes.
- Figure 5 - Recommend to show ALF as a negative control.
- Figure 6c - Please provide slopes in text or normalized data to allow comparison.
- Figure 8 - Bar 'T' is blue for both human and mouse.
- Figure 9 - This figure is less informative; suggest move to supplementary material.
- Table 2 - It would be helpful to explain the rationale of block sampling for the metabolomics study but not for the glucose experiment.

- The page layout can be improved by re-sizing figures (for example fig.2, 3, 4) and moving some long tables (table 5 and 6) to supplementary materials.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Infectious disease; Circadian biology; Immunology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 08 Oct 2021

Kimberley Prior, University of Edinburgh, Edinburgh, UK

We are very grateful to Reviewer 2, Xiaodong (Alan) Zhuang, for making such constructive comments and we have revised the manuscript as suggested, detailed in the following responses.

Specific comments:

1) Abstract - Define TTFL.

RESPONSE: Now included.

2) Figure 1 - Legend (B) "The experimental...".

RESPONSE: Now fixed.

3) Figure 3 - Please label the top right and bottom left boxes.

RESPONSE: We removed the background colours from these portions of the Fig (in response to Reviewer 1 and explained what these regions are in the legend

4) Figure 5 - Recommend to show ALF as a negative control.

RESPONSE: Great idea, because the best fitting pattern for this group is linear, so we have indicated intercept in the legend and on the revised figure (also, this has become Figure 6).

5) Figure 6c - Please provide slopes in text or normalized data to allow comparison.

RESPONSE: This now Figure 7. The statistical analysis revealing a significant interaction confirms the slopes are different and the aim is to simply compare the groups rather than parameterise developmental patterns

6) Figure 8 - Bar 'T' is blue for both human and mouse.

RESPONSE: Thanks for spotting that – now fixed

7) Figure 9 - This figure is less informative; suggest move to supplementary material.

RESPONSE: We agree but the journal no long supports supplementary material so our understanding is that has be included in the vicinity of where it is cited.

8) Table 2 - It would be helpful to explain the rationale of block sampling for the metabolomics study but not for the glucose experiment.

RESPONSE: We've included a pointer to the legend to where this is explained in the methods section.

9) The page layout can be improved by re-sizing figures (for example fig.2, 3, 4) and moving some long tables (table 5 and 6) to supplementary materials.

RESPONSE: We agree but our understanding is that the location and size of elements is determined by the journal and outwith author's control.

Competing Interests: The authors declare no competing interests

Reviewer Report 25 August 2021

<https://doi.org/10.21956/wellcomeopenres.18637.r45125>

© 2021 Larrondo L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Luis Larrondo

¹ ANID - Millennium Science Initiative, Millennium Institute for Integrative Biology (iBIO), Santiago, Chile

² Pontificia Universidad Católica de Chile, Santiago, Chile

The work by Prior *et al.*, is a nicely written MS that reports extremely interesting data on how malaria parasites may be sensing and responding to circulating metabolic host cues to synchronize their biology. Surprisingly, under time restricted feeding protocol the synchronizing

cue is not glucose levels (as it could have been suspected from other studies) but instead, is isoleucine. Importantly, this appears to be an active strategy by *Plasmodium* as opposed to a constraint imposed by host rhythms, as isoleucine deprivation does not appear to have a major cost compared to other essential nutrients. While the MS does not go into the mechanistic details on how the sensing of isoleucine occurs or on how it gates rhythmic asexual development, it does describe an important set of original and valuable observations that will help to advance the field.

Something that needs to be clarified for all readers (without having to go back to prior publications) is why this particular experimental design was selected. In particular, one would have expected the experimental groups to also include WT in constant darkness both under ALF and TRF conditions. Yet, the only animals kept under DD were mice devoid of a functional clock (*per1/2* DKO). While one could try to interpret the logic based on O'Donnell *et al.*, 2020, it is crucial for general readers to clearly grasp the underlying reasons of the design (i.e why there was no ALF for WT animals). This information is important since other reports (Rijo-Ferreira *et al.*, 2020) have shown that the parasite rhythms persist even under ad libitum conditions in WT animals (something that was not observed in O'Donnell *et al.*, 2020) yet, was such condition (WT ad libitum feeding) was not analyzed in the current MS. In addition, the data by Rijo-Ferreira *et al.*, 2020 show that in arrhythmic animals the parasite rhythms persist for about 4 days before becoming arrhythmic.

Hirako *et al.* found that rhythmic oscillation in blood glucose levels was associated with *P. chabaudi* cell cycle synchronization. Yet, in light of the results of Prior *et al.*, it appears that this would be an association and not causation. Nevertheless, the same study by Hirako *et al.* implicated TNFR in this mechanism, and sure enough $-/-$ animals showed no rhythms in the parasite proliferation. These observations should be further discussed.

Fig 1, legend: "cycles of 12-hours with food followed by 12 h without access to food (TRF, time-restricted feeding)".

To use TRF only here is somehow weird as it is equivalent to the other protocols (light or dark feeding). Thus, the other protocols are also equivalent to TRF. Indeed, in several of the papers cited in the MS (as well as in many other publications) what is presented here as light or dark feeding would be called TRF.

Figur3 3B: Gray color is not explained

"Mean blood-glucose concentration differed between the groups, being higher in DF and TRF mice (DF=8.55±0.14 mmol/L, TRF=8.59±0.13 mmol/L)"

These numbers are hard to recognize in the data (Fig 4), where the lines and shading are mean ± SEM glucose concentrations. Indeed, in most cases, the SEM in the plots span at least 1 unit (quite different from the +/- 0.1 units referred to in the text). Are those glucose concentration average values, or values at a particular moment?

"Isoleucine cannot be perturbed in vivo without causing confounding off-target effects on the host or interference by host homeostasis".

References and a brief explanation are needed.

Fig 8. The orange bar for I (human) is missing and instead, the blue bar (rodent) has twice the width. Since Ile is absent from the human hemoglobin it should be replaced by a Y-axis 0 orange

bar, instead of being replaced by a broader blue one.

Fig 9 contains important data, although it is plotted in a way that is not useful. Each of the columns in the x axis should have a number that allows identifying (from the legend) the corresponding metabolite. Likewise, changing the size of each sphere (dot) would allow better visual resolution between one set and the other

Ideally one would like to see experiments that would compare the qualitative effect of diets on the parasite rhythms. In particular, it would be interesting to compare existing feeding paradigms (i.e high fat diet, TRF vs *ad libitum*) versus high protein diets (TRF vs *ad libitum*). Yet, conducting and analyzing these experiments would be another paper in itself. Nevertheless, there may be existing literature that allows establishing interesting correlations between protein (Ile) content and the characteristics of the disease. Likewise, one would expect that this effect would be clearer in humans, particularly in cases where dietary interventions/ food supplements modify isoleucine circulating levels. While such association would establish only correlation (and not causation), it would enrich the discussion.

"Wild type C57BL/6 mice were housed" Specify sex of the animals.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: chronobiology, molecular genetics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 09 Oct 2021

Kimberley Prior, University of Edinburgh, Edinburgh, UK

We are very grateful to Reviewer 1, Luis Larrondo, for making such constructive comments and we have revised the manuscript as suggested, detailed in the following responses.

COMMENT 1) Something that needs to be clarified for all readers (without having to go back to prior publications) is why this particular experimental design was selected. In particular, one would have expected the experimental groups to also include WT in constant darkness both under ALF and TRF conditions. Yet, the only animals kept under DD were mice devoid of a functional clock (per1/2 DKO). While one could try to interpret the logic based on O'Donnell et al., 2020, it is crucial for general readers to clearly grasp the underlying reasons of the design (i.e why there was no ALF for WT animals). This information is important since other reports (Rijo-Ferreira et al., 2020) have shown that the parasite rhythms persist even under ad libitum conditions in WT animals (something that was not observed in O'Donnell et al., 2020) yet, was such condition (WT ad libitum feeding) was not analyzed in the current MS. In addition, the data by Rijo-Ferreira et al., 2020 show that in arrhythmic animals the parasite rhythms persist for about 4 days before becoming arrhythmic.

RESPONSE: As noted, other studies have focused on asking what IDC schedules occur under different combinations of light/dark cycles, feeding-fasting treatments and host TTFL manipulations. Rijo-Ferreira et al., 2020 reveals that the process simply of food intake does not schedule the IDC, thus O'Donnell et al., 2020 and others (Hirako et al., 2018, Prior et al., 2018) are best interpreted as finding process(es) involved in food digestion/fasting schedule the IDC. The aim of the present study was to leverage these observations – which have proved largely robust across studies – to compare and contrast treatments in which different IDC schedules occur. As such, a fully factorial design including WT mice in DD (whether TRF or ALF) would not provide a different IDC schedule to the treatments we selected. This is because (i) the IDC schedule is not directly scheduled by LD cycles (Prior et al., 2018, Hirako et al., 2018); (ii) WT mice in DD would free run (with a period very close to 24 hr in this strain (Schwartz and Zimmerman, 1990)) and maintain the same feeding rhythms under ALF as WT ALF mice in LD cycles (O'Donnell et al., 2020); (iii) if we applied TRF to WT mice in DD in subjective day, this would mirror the conditions in WT LF mice; and (iv) we focused on using TRF treatments (i.e. light-fed, LF, and dark-fed, DF) for the WT mice to avoid the potentially confounding effects of ALF mice being able to take in more food across the circadian cycle on metabolite concentrations. Further, adding more treatment groups would have traded-off against the number of replicates within treatments, due to the time constraints on completing sample collection within an appropriate window. We have added an explanation for the design at the start of the results section.

COMMENT 2) Hirako et al. found that rhythmic oscillation in blood glucose levels was associated with *P. chabaudi* cell cycle synchronization. Yet, in light of the results of Prior et al., it appears that this would be an association and not causation. Nevertheless, the same study by Hirako et al. implicated TNFR in this mechanism, and sure enough *-/-* animals showed no rhythms in the parasite proliferation. These observations should be further discussed.

RESPONSE: The 3rd paragraph of the discussion explores how the results of Hirako et al. and indeed our previous work (Prior et al., 2018) can be reconciled with our findings about isoleucine. For example, glucose is essential for isoleucine uptake (Martin & Kirk, 2007), and high blood

glucose concentrations of isoleucine are associated with uptake of glucose by host cells, including by TNF stimulated immune cells (Elsed & Playfair, 1994; Hirako et al., 2018). We have altered this paragraph to conclude that whilst isoleucine appears to be the main driver of the IDC schedule, glucose likely mediates its impact via multiple, non-mutually exclusive, activities. Further, we have now included another experiment and new Figure providing more evidence that glucose is not a direct driver of the IDC schedule.

COMMENT 3) Fig 1, legend: “cycles of 12-hours with food followed by 12 h without access to food (TRF, time- restricted feeding”. To use TRF only here is somehow weird as it is equivalent to the other protocols (light or dark feeding). Thus, the other protocols are also equivalent to TRF. Indeed, in several of the papers cited in the MS (as well as in many other publications) what is presented here as light or dark feeding would be called TRF.

RESPONSE: Good point! We have clarified that these are all forms of TRF but to keep the acronyms of the treatment groups easy to follow we have retained our nomenclature.

COMMENT 4) Figur3 3B: Gray color is not explained

RESPONSE: This was just a background colour. To avoid confusion we have changed this to white.

COMMENT 5) “Mean blood-glucose concentration differed between the groups, being higher in DF and TRF mice (DF=8.55±0.14 mmol/L, TRF=8.59±0.13 mmol/L)” These numbers are hard to recognize in the data (Fig 4), where the lines and shading are mean ± SEM glucose concentrations. Indeed, in most cases, the SEM in the plots span at least 1 unit (quite different from the +/- 0.1 units referred to in the text). Are those glucose concentration average values, or values at a particular moment?

RESPONSE: We quoted the overall means ±SEM of the time series (i.e. intercepts) in the text and have clarified this. We note that visual inspection of Fig4 suggests the mean patterns for DF and TRF exceed 8mmol/L for most of the sampling window, unlike the LF and AL groups.

COMMENT 6) “Isoleucine cannot be perturbed in vivo without causing confounding off-target effects on the host or interference by host homeostasis”. References and a brief explanation are needed.

RESPONSE: We searched the literature for approaches to perturbing blood-isoleucine concentrations in mice but were unrewarded because there are several issues to overcome. Isoleucine is essential for mice as well as parasites, so hosts cannot be maintained on isoleucine-free food without negative health consequences, particularly via its roles in the immune system (Rivas- Santiago et al, 2011). For example, even short-term limitation of isoleucine to 50% of optimal for 3 weeks results in altered immune performance (Mao et al, 2018), including a reduction in complement making it difficult to separate the consequences isoleucine perturbations from depressed innate immune defences that are crucial in the acute phase of malaria infection. Isoleucine also modulates VEGF expression which is involved in recovery from anaemia (Murata et al, 2007), a hallmark of malaria infection. Thus, an in vitro system allows far more precise control over isoleucine levels and the timing of its availability to parasites for our proof of principle experiments. We have now summarised this in the discussion.

COMMENT 7) Fig 8. The orange bar for I (human) is missing and instead, the blue bar (rodent) has twice the width. Since Ile is absent from the human hemoglobin it should be replaced by a Y-axis 0 orange bar, instead of being replaced by a broader blue one.

RESPONSE: Thanks for spotting this “graphical typo”. Now corrected

COMMENT 8) Fig 9 contains important data, although it is plotted in a way that is not useful. Each of the columns in the x axis should have a number that allows identifying (from the legend) the corresponding metabolite. Likewise, changing the size of each sphere (dot) would allow better visual resolution between one set and the other

RESPONSE: Apologies for the confusion. We have added numbers to the x axis that correspond to a metabolite numbering system added to the legend. We have now plotted the median concentration for each treatment for every metabolite, instead of plotting the raw data. As this figure is not about the metabolite dynamics, but about which metabolites remain in our analysis, this results in 4 data points per metabolite and a much simpler figure.

COMMENT 9) Ideally one would like to see experiments that would compare the qualitative effect of diets on the parasite rhythms. In particular, it would be interesting to compare existing feeding paradigms (i.e high fat diet, TRF vs ad libitum) versus high protein diets (TRF vs ad libitum). Yet, conducting and analyzing these experiments would be another paper in itself. Nevertheless, there may be existing literature that allows establishing interesting correlations between protein (Ile) content and the characteristics of the disease. Likewise, one would expect that this effect would be clearer in humans, particularly in cases where dietary interventions/ food supplements modify isoleucine circulating levels. While such association would establish only correlation (and not causation), it would enrich the discussion.

RESPONSE: We agree these are all interesting avenues for future work – particularly, by comparing and contrasting parasite gene expression patterns across multiple feeding paradigms, insight into the parasite’s time keeping mechanism might be revealed. Given that malaria is almost exclusively a disease in LMICs and that controlled human infection trials do not consider dietary details, this kind of information is not available to discuss.

COMMENT 10) “Wild type C57BL/6 mice were housed” Specify sex of the animals.

RESPONSE: Sexes were already specified in the detailed descriptions of each experiment, in the methods section

REFERENCES

- Elased K, Playfair JH: Hypoglycemia and hyperinsulinemia in rodent models of severe malaria infection. *Infect Immun.* 1994;62(11):5157–5160. 7927799 10.1128/iai.62.11.5157-5160.1994 303239
- Hirako IC, Assis PA, Hojo-Souza NS, et al.: Daily Rhythms of TNF α Expression and Food Intake Regulate Synchrony of Plasmodium Stages with the Host Circadian Cycle. *Cell Host Microbe.* 2018;23(6):796–808.e6. 29805094
- Mao et al. 2018. I-Isoleucine Administration Alleviates Rotavirus Infection and Immune Response in the Weaned Piglet Model. *Front. Immunol.*, 16 July 2018. PMCID: PMC6054962
- Martin RE, Kirk K: Transport of the essential nutrient isoleucine in human erythrocytes infected with the malaria parasite Plasmodium falciparum. *Blood.* 2007;109(5):2217–2224. 17047158 10.1182/blood-2005-11-026963
- Murata et al. 2007. Isoleucine, an Essential Amino Acid, Prevents Liver Metastases of Colon Cancer by Antiangiogenesis. *Cancer Res* 2007 Apr 1;67(7):3263-8. PMID: 17409434
- O'Donnell AJ, Prior KF, Reece SE: Host circadian clocks do not set the schedule for the within-

host replication of malaria parasites. Proc Biol Sci. 2020;287(1932):20200347. 32781954 10.1098/rspb.2020.0347 7575513

Prior KF, van der Veen DR, O'Donnell AJ, et al.: Timing of host feeding drives rhythms in parasite replication. PLoS Pathog. 2018;14(2):e1006900. 29481559 10.1371/journal.ppat.1006900 5843352

Rijo-Ferreira F, Acosta-Rodriguez VA, Abel JH, et al.: The malaria parasite has an intrinsic clock. Science. 2020;368(6492):746–753. 32409471 10.1126/science.aba2658 7409452

Rivas-Santiago et al. 2011. Induction of β -defensins by l-isoleucine as novel immunotherapy in experimental murine tuberculosis. Clin Exp Immunol. 2011 Apr;164(1):80-9. PMCID: PMC3074220

Schwartz, W, Zimmerman, P. Circadian timekeeping in BALB/c and C57BL/6 inbred mouse strains. The Journal of Neuroscience. 1990 Nov;10(11):3685-3694.

Competing Interests: The authors have no competing interests