RESEARCH ARTICLE

Increased expression of Pol δ does not alter the canonical replication program *in vivo* [version 1; peer review: 2 approved]

Róbert Zach, Antony M. Carr

Genome Damage and Stability Centre, School of Life Sciences, Science Park Road, University of Sussex, Falmer, Brighton, BN1 9RQ, UK

v1 First published: 26 Feb 2021, 6:44
<https://doi.org/10.12688/wellcomeopenres.16600.1>
 Latest published: 04 May 2021, 6:44
<https://doi.org/10.12688/wellcomeopenres.16600.2>

Abstract

Background: *In vitro* experiments utilising the reconstituted *Saccharomyces cerevisiae* eukaryotic replisome indicated that the efficiency of the leading strand replication is impaired by a moderate increase in Pol δ concentration. It was hypothesised that the slower rate of the leading strand synthesis characteristic for reactions containing two-fold and four-fold increased concentration of Pol δ represented a consequence of a relatively rare event, during which Pol δ stochastically outcompeted Pol ϵ and, in an inefficient manner, temporarily facilitated extension of the leading strand. Inspired by this observation, we aimed to determine whether similarly increased Pol δ levels influence replication dynamics *in vivo* using the fission yeast *Schizosaccharomyces pombe* as a model system.

Methods: To generate *S. pombe* strains over-expressing Pol δ , we utilised Cre-Lox mediated cassette exchange and integrated one or three extra genomic copies of all four Pol δ genes. To estimate expression of respective Pol δ genes in Pol δ -overexpressing mutants, we measured relative transcript levels of *cdc1*⁺, *cdc6*⁺ (or *cdc6*^{L591G}), *cdc27*⁺ and *cdm1*⁺ by reverse transcription followed by quantitative PCR (RT-qPCR). To assess the impact of Pol δ over-expression on cell physiology and replication dynamics, we used standard cell biology techniques and polymerase usage sequencing.

Results: We provide an evidence that two-fold and four-fold over-production of Pol δ does not significantly alter growth rate, cellular morphology and S-phase duration. Polymerase usage sequencing analysis further indicates that increased Pol δ expression does not change activities of Pol δ , Pol ϵ and Pol α at replication initiation sites and across replication termination zones. Additionally, we show that mutants over-expressing Pol δ preserve WT-like distribution of replication origin efficiencies.

Conclusions: Our experiments do not disprove the existence of opportunistic polymerase switches; however, the data indicate that, if

Open Peer Review

Approval Status

	1	2
version 2 (revision) 04 May 2021	 view	
version 1 26 Feb 2021	 view	 view

1. **Jie Ren** , Chinese Academy of Sciences, Beijing, China

2. **Sarah Sabatinos**, Ryerson University, Toronto, Canada

Any reports and responses or comments on the article can be found at the end of the article.

stochastic replacement of Pol ϵ for Pol δ does occur *in vivo*, it represents a rare phenomenon that does not significantly influence canonical replication program.

Keywords

Polymerase δ , over-expression, DNA replication, polymerase switch, polymerase usage sequencing, *Schizosaccharomyces pombe*, fission yeast

Corresponding author: Antony M. Carr (a.m.carr@sussex.ac.uk)

Author roles: **Zach R:** Conceptualization, Data Curation, Formal Analysis, Methodology, Project Administration, Software, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Carr AM:** Conceptualization, Funding Acquisition, Methodology, Project Administration, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Wellcome Trust through a Wellcome Trust Investigator Award [110047].
The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Zach R and Carr AM. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Zach R and Carr AM. **Increased expression of Pol δ does not alter the canonical replication program *in vivo* [version 1; peer review: 2 approved]** Wellcome Open Research 2021, 6:44 <https://doi.org/10.12688/wellcomeopenres.16600.1>

First published: 26 Feb 2021, 6:44 <https://doi.org/10.12688/wellcomeopenres.16600.1>

Introduction

Unchallenged duplication of the eukaryotic genome requires the coordinated action of three replicative polymerase complexes: Pol α -primase (hereafter referred to as Pol α), Pol δ and Pole (Burgers & Kunkel, 2017). According to the canonical model of eukaryotic replication, Pol α and Pol δ cooperate to discontinuously synthesise the lagging strand via the iterative production of short Okazaki fragments (OF), ca. 150bp, whereas Pole carries out continuous leading strand replication (Clausen *et al.*, 2015; Daigaku *et al.*, 2015; Miyabe *et al.*, 2011). Interestingly, such strict division of labour does not always apply, and deviations have been documented (Guilliam & Yeeles, 2020a).

While polymerase activities of Pol α and Pol δ are indispensable for cell survival, the polymerase domain of Pole is not required for completion of replication in either *Saccharomyces cerevisiae* or *S. pombe* (Feng & D'Urso, 2001; Kesti *et al.*, 1999). In both yeast experimental models it has been demonstrated that Pol δ facilitates the leading strand synthesis when catalytically-inactive Pole is expressed (Garbacz *et al.*, 2018; Miyabe *et al.*, 2015). Such findings have found support in *in vitro* experiments utilising reconstituted replisome system (Yeeles *et al.*, 2017), confirming that, under certain circumstances, Pol δ is competent in the leading strand replication.

Indeed, it has been reported that Pol δ replicates both DNA strands during homologous recombination restarted replication in *S. pombe* (Miyabe *et al.*, 2015) and break induced replication in *S. cerevisiae* (Donnianni *et al.*, 2019). Additionally, genomic analysis by polymerase usage sequencing (Pu-Seq) or HydEn-seq revealed that Pol δ is involved in the initiation of the leading strand replication in unperturbed *S. cerevisiae* and *S. pombe* cells, respectively (Daigaku *et al.*, 2015; Garbacz *et al.*, 2018; Zhou *et al.*, 2019). In agreement with such findings, PCNA-associated Pol δ has been shown to play an important role in early stages of the leading strand replication *in vitro* (Aria & Yeeles, 2018; Yeeles *et al.*, 2017). Moreover, it has recently been proposed that Pol δ takes over the leading strand synthesis prior to replication fork termination (Zhou *et al.*, 2019). The exact role of Pol δ during the final stages of replisome progression is, however, yet to be clarified.

Apart from homologous recombination dependent DNA synthesis and replication initiation, Pol δ -mediated leading strand synthesis has been shown to occur in the context of polymerase uncoupling. It has been reported that cyclobutane pyrimidine dimer driven disengagement of CMG-associated Pole from the leading 3'OH generates a gap, the efficient filling of which requires the translesion synthesis machinery, as well as the action of Pol δ (Guilliam & Yeeles, 2020b). In further support of a more generic function of Pol δ in the leading strand synthesis, Pol δ has been shown to proofread errors introduced by Pole in hyper mutator *pol2-M644G* mutants (Bullock *et al.*, 2020). In line with all aforementioned observations, it has been shown that CMG-associated Pole exists in two mutually-exclusive conformations, of which only one facilitates DNA synthesis (Zhou *et al.*, 2017).

Intriguingly, according to *in vitro* studies of eukaryotic replication, two-fold and four-fold increase in Pol δ concentration reduces the rate of the leading strand synthesis (Yeeles *et al.*, 2017). It has been suggested that the observed retardation of leading strand replication represents a consequence of stochastic polymerase switching, during which Pol δ outcompetes Pole and temporarily facilitates inefficient extension of the leading 3' end. Since the effect of Pol δ concentration on replisome progression and the hypothetical phenomenon of leading strand polymerase switching has not been investigated *in vivo*, we aimed to test whether similar a phenomenon manifests in living cells, potentially shedding light on a yet uncomprehended promiscuity of replicative polymerases.

Methods

Yeast culture and transformation

S. pombe cells were grown in yeast extract (YE) (Formedium, PCM0155) with supplements (Formedium, PSU0110) medium according to standard procedures (Petersen & Russell, 2016). Briefly, cells (25% glycerol stocks stored at -80°C) were streaked onto an agar plate and incubated at 30°C for 2–3 days. Next, cells were inoculated into a liquid medium and cultivated at 30°C for ca. 36 h in the ISF-1-W shaker (Kuhner) with constant shaking (180 rpm). Cultures were diluted accordingly two times during the course of cultivation. Then appropriate amounts of cells were collected (depending on experiment) and processed further. Cells were transformed by the lithium-acetate based method (Bähler *et al.*, 1998). Optical density (OD) of liquid cell cultures was assessed by WPA CO8000 Cell Density Meter (Biochrom). Doubling times were calculated using the formula: $DT = 1/k$, where DT stands for doubling time and k represents the slope of linear regression computed from a time-series of log₂-transformed OD measurements. A list of strains used in this study is provided in Table 1.

Microscopy

1 mL of exponentially growing cells was centrifuged (1000 × g, 5 min, 25°C) and the cell pellet resuspended in 1 mL of 70% ethanol. 500 μ L of fixed cells were collected by centrifugation (1000 × g, 5 min, 25°C) and re-suspended in 50 μ L of H₂O containing 1 μ M 4',6-diamidino-2-phenylindole (DAPI). Cells were incubated at room temperature in the dark for at least 15 min, and then analysed by microscopy using a Nikon E400 system. Cell lengths were determined from DIC images by measuring the distance between the opposite poles of the cell using ImageJ software (version 1.51m9) (Schneider *et al.*, 2012). At least 200 cells per sample were scored.

Cell synchronisation and DNA content analysis

Exponentially growing *cdc2^{asM17}* cells ($OD_{600} = 0.1$ – 0.2 ; 1 – 2×10^6 cells/mL) were treated with 2 μ M 3-Br-PP1 (abcam, ab143756) for 3 h. A 50mL fraction 3-Br-PP1-treated culture was centrifuged (1000 × g, 5 min, 25°C) and the cell pellet washed with 50 mL of fresh YES medium pre-heated to 30°C. Washed cells were re-suspended in 50 mL of fresh pre-heated YES and incubated at 30°C. In 15-min intervals, 1mL aliquots of

Table 1. List of strains.

ID	Genotype	Origin
RZ42	<i>h- ade6-704 leu1-32 ura4-D18 I-5230932:[LoxP-cdc1-cdc27-ura4-cdc6-cdm1-LoxM3]</i>	This study
RZ47	<i>h- ade6-704 leu1-32</i>	Laboratory stock
RZ93	<i>h- ade6-704 leu1-32 ura4-D18 I-3325162:[LoxP-cdc1-cdc27-kanR-cdc6-cdm1-LoxM3] I-4734015:[LoxP-cdc1-cdc27-natR-cdc6-cdm1-LoxM3] I-5230932:[LoxP-cdc1-cdc27-ura4-cdc6-cdm1-LoxM3]</i>	This study
655	<i>h- ade6-? leu1-32 ura4-D18 rnh201::kanR cdc20M630F</i>	Laboratory stock
856	<i>h- ade6-704 leu1-32 ura4-D18 rnh201::kanR cdc6L591G</i>	Laboratory stock
1141	<i>h- ade6-704 leu1-32 ura4-D18 rnh201::kanR pol1L850F</i>	Laboratory stock
RZ57	<i>h- ade6-704 leu1-32 ura4-D18 I-5230932:[LoxP-cdc1-cdc27-ura4-cdc6L591G-cdm1-LoxM3] rnh201::hygR cdc6L591G</i>	This study
RZ62	<i>h- ade6-704 leu1-32 ura4-D18 I-5230932:[LoxP-cdc1-cdc27-ura4-cdc6-cdm1-LoxM3] rnh201::hygR cdc20M630F</i>	This study
RZ68	<i>h- ade6-704 leu1-32 ura4-D18 I-5230932:[LoxP-cdc1-cdc27-ura4-cdc6-cdm1-LoxM3] rnh201::hygR pol1L850F</i>	This study
RZ112	<i>h- ade6-704 leu1-32 ura4-D18 I-3325162:[LoxP-cdc1-cdc27-kanR-cdc6L591G-cdm1-LoxM3] I-4734015:[LoxP-cdc1-cdc27-natR-cdc6L591G-cdm1-LoxM3] I-5230932:[LoxP-cdc1-cdc27-ura4-cdc6L591G-cdm1-LoxM3] rnh201::hygR cdc6L591G</i>	This study
RZ116	<i>h- ade6-704 leu1-32 ura4-D18 I-3325162:[LoxP-cdc1-cdc27-kanR-cdc6-cdm1-LoxM3] I-4734015:[LoxP-cdc1-cdc27-natR-cdc6-cdm1-LoxM3] I-5230932:[LoxP-cdc1-cdc27-ura4-cdc6-cdm1-LoxM3] rnh201::hygR pol1L850F</i>	This study
RZ118	<i>h- ade6-704 leu1-32 ura4-D18 I-3325162:[LoxP-cdc1-cdc27-kanR-cdc6-cdm1-LoxM3] I-4734015:[LoxP-cdc1-cdc27-natR-cdc6-cdm1-LoxM3] I-5230932:[LoxP-cdc1-cdc27-ura4-cdc6-cdm1-LoxM3] rnh201::hygR cdc20M630F</i>	This study
RZ331	<i>h- ade6-704 leu1-32 cdc2asM17</i>	This study
RZ332	<i>h- ade6-704 leu1-32 cdc2asM17 I-5230932:[LoxP-cdc1-cdc27-ura4-cdc6-cdm1-LoxM3]</i>	This study
RZ333	<i>h- ade6-704 leu1-32 cdc2asM17 I-3325162:[LoxP-cdc1-cdc27-kanR-cdc6-cdm1-LoxM3] I-4734015:[LoxP-cdc1-cdc27-natR-cdc6-cdm1-LoxM3] I-5230932:[LoxP-cdc1-cdc27-ura4-cdc6-cdm1-LoxM3]</i>	This study

synchronous cell culture were centrifuged (1000 × g, 3 min, 25°C) and collected cells fixed in 1 mL of 70% ethanol. For each time point, 500 µL of fixed cells were centrifuged (1000 × g, 3 min, 25°C), the supernatant was discarded, and the cell pellet re-suspended in 500 µL of sodium citrate (50 mM, pH = 7) containing 1 mg/mL RNase A (NEB, T3018L). The resulting cell suspension was incubated for 3 h at 37°C, and then mixed with 500 µL of sodium citrate (50 mM, pH = 7) containing 2µM SYTOX Green (Invitrogen, S7020). Samples were analysed using an Accuri C6 flow cytometry system (Beckman Coulter). Data were analysed by BD CSampler software (version 1.0.264.21) and R (version 4.0.0) (<https://www.R-project.org/>; R Core Team, 2020).

Cre-recombination mediated cassette exchange (RMCE)
Leucine-auxotrophic cells carrying one of three LoxP-LoxM3 integration sites (I-3325162:[LoxP-rtS-ura4⁺], I-4734015:[LoxM3-kanR-LoxP], I-5230932:[LoxM3-kanR-LoxP]) were transformed with one of the Polδ Cre-Lox integration vectors (pRZ02, pRZ03, pRZ04, pRZ05, pRZ06, pRZ07) listed in Table 2. Leucine-prototrophic transformants (containing respective Polδ Cre-Lox integration vector) were selected on EMM plates lacking

leucine. Single clones were selected and grown over-night in liquid YES medium. Next, leucine-auxotrophic colonies (lacking the transformed Polδ Cre-Lox vector) carrying a selection marker associated with respective Polδ integration events were selected (Watson *et al.*, 2008).

Plasmids

Cre-Lox integration plasmids carrying 4 Polδ genes (*cdc1⁺*, *cdc27⁺*, *cdm1⁺*, *cdc6⁺* or *cdc6^{L591G}*) and one of the three selection markers (NatR, KanR, ura4⁺) were derived from the previously characterised pAW8 vector (Addgene, 110222) (Watson *et al.*, 2008) by standard restriction insertion cloning. Briefly, insert DNA fragments (*SphI-cdc1⁺-ApaI*, *ApaI-cdc27⁺-XhoI*, *XhoI-ura4⁺-SacI*, *SacI-cdc6⁺-SbfI*, *SacI-cdc6^{L591G}-SbfI*, *SbfI-cdm1⁺-SpeI*, *ApaI-cdc27⁺-NatMX6-SacI*, *ApaI-cdc27⁺-KanMX6-SacI*) were generated by high-fidelity PCR with KOD Hot Start DNA Polymerase (Merck Millipore, 71085-3) and purified with QIAquick PCR Purification Kit (QIAGEN, 28104) or QIAquick Gel Extraction Kit (QIAGEN, 28704). *SacI-cdc6^{L591G}-SbfI*, *ApaI-cdc27⁺-NatMX6-SacI* and *ApaI-cdc27⁺-KanMX6-SacI* were produced by overlap extension PCR. Generated Polδ gene fragments contained intact 5'UTR and 3'UTR sequences, as

well as upstream and downstream regions of 663–980 bp. pAW8 vector and insert DNA fragments were digested by respective restriction enzymes and ligated over-night at 18°C using T4 DNA ligase (NEB, M0202S). Each ligation reactions contained 50 ng of pAW8 vector and three-fold molar excess of respective DNA fragments. Ligation reactions were incubated in T3 Thermocycler (Biometra). Restriction digestion reactions were performed according to manufacturer's instructions with the following restriction enzymes: *SphI*-HF (NEB, R3182S); *ApaI* (NEB, R0114S); *XhoI* (NEB, R0146S); *SacI*-HF (NEB, R3156S); *SpeI*-HF (NEB, R3133S); *SbfI*-HF (NEB, R3642S); *SalI*-HF (NEB, R3138S); *BamHI*-HF (NEB, R3136S). Ligation products were transformed into DH5-Alpha *E. coli* competent cells. Plasmids were purified using a QIAprep Spin Miniprep Kit (QIAGEN, 27104). A list of generated plasmids is provided in [Table 2](#).

RT-qPCR

Total RNA was isolated from 1–2 mL of exponentially growing cells ($OD_{600} = 0.5$; 5×10^6 cells/mL) using MasterPure Yeast RNA purification kit (Cambio Ltd, MPY03100). RNA was converted to cDNA utilising a RevertAid First Strand cDNA

Synthesis Kit (ThermoFisher Scientific, K1621) and random hexamer primers. Relative transcript levels were determined by qPCR with Luna Universal qPCR Master Mix (NEB, M3003E) and an AriaMx Real-time PCR System (Agilent Technologies). qPCR reactions were prepared by mixing 10 μ L of 2 $\times$ Luna Universal qPCR Master Mix, 0.5 μ L of forward and reverse primers (10 μ M), 2 μ L of 1000-fold diluted cDNA and 7 μ L of nuclease-free H₂O. Thermal cycling conditions were: Hot Start: 95°C for 3 min; Cycling (45 $\times$): 95°C for 15 s, 60°C for 30 s. Relative transcript levels were calculated using the equation $RNA_{\text{target}} = 2^{-Cq(\text{target})} / 2^{-Cq(\text{reference})}$, where RNA_{target} represents the relative transcript level of a given target gene, and $Cq(\text{target})$ and $Cq(\text{reference})$ stand for PCR cycle quantification values of target and reference genes, respectively. *act1* was used as the reference gene. A list of qPCR primers (obtained from Integrated DNA Technologies) used in this study is provided in [Table 3](#).

Pu-Seq library preparation

For all strains presented, two sets of Pu-Seq libraries were prepared. One set was prepared as described previously ([Daigaku et al., 2015](#); [Keszthelyi et al., 2015](#)). Briefly,

Table 2. List of plasmids.

ID	Annotation	Origin
pRZ02	pAW8_cdc1-cdc27-ura4-cdc6-cdm1	This study, derived from pAW8 (Watson et al., 2008)
pRZ03	pAW8_cdc1-cdc27-ura4-cdc6L591G-cdm1	
pRZ04	pAW8_cdc1-cdc27-natR-cdc6-cdm1	
pRZ05	pAW8_cdc1-cdc27-natR-cdc6L591G-cdm1	
pRZ06	pAW8_cdc1-cdc27-kanR-cdc6-cdm1	
pRZ07	pAW8_cdc1-cdc27-kanR-cdc6L591G-cdm1	

Table 3. List of qPCR primers.

ID	Sequence (5' – 3')	Target	Origin
RZ67	CAACTATCCTTCTCAACAG	<i>cdc1</i> (134 bp)	This study
RZ68	GCTAGTAGCCAACACAAAATG		
RZ69	CGTTCACGATTCTGAAGATG	<i>cdc27</i> (102 bp)	This study
RZ70	ATAATTCTCTGAGGTTCTGTC		
RZ75	CCTGCAATAAATCTGAGAAG	<i>cdc6</i> (109 bp)	This study
RZ76	CATTGTCTAGTAACACCAAAC		
RZ81	TTCATTCTAGTACCGCAGTG	<i>cdm1</i> (142 bp)	This study
RZ82	TGTGGGATTGACTTGAATTAC		
RZ87	TCCTCATGCTATCATGCGTCTT	<i>act1</i> (78 bp)	(Převorovský et al., 2016)
RZ88	CCACGCTCCATGAGAATCTTC		

10–20 µg of genomic DNA containing increased quantities of misincorporated ribonucleotides (rNMPs) was treated with 0.3M NaOH for 2 h at 55°C. Digested DNA was run on a 2% agarose gel (in 0.5× TBE). 300–500bp ssDNA fragments were gel extracted and subjected to complementary second strand synthesis primed by random 8-mers (obtained from Integrated DNA technologies). Resulting double-stranded (dsDNA) fragments were converted to Illumina sequencing libraries using NEBNext Ultra DNA library prep kit for Illumina (NEB, E7645S) and NEBNext multiplex oligos for Illumina (NEB, E7335). The second set of Pu-Seq libraries was prepared according to a modified version of the established GLOE-Seq protocol (Sriramachandran *et al.*, 2020), which utilises two subsequent ligations of adapter/splinter oligonucleotides, first to rNMP-dependent phosphorylated 5' ends and, following sonication, to 3' ends of the ssDNA fragments. rNMP-dependent 5' ends were generated from genomic DNA containing increased quantities of misincorporated ribonucleotides treated by RNase H2 (NEB, M0288S) and subsequently denatured at 95°C. Sequencing was performed on an Illumina NextSeq 500 sequencer. Sequencing reads were mapped onto the reference genome using Bowtie2 (Langmead & Salzberg, 2012).

Pu-Seq data analysis

Polymerase tracks at any given 300bp bin were calculated using the equation $PT = (R_T + R_B) / (R_T - R_B)$, where PT represents polymerase track, and R_T and R_B stand for rNMPs mapped to the top and the bottom DNA strands, respectively. Polymerase tracks were determined for each biological repeat separately, then averages of the two repeats were used for subsequent analysis. Positions and efficiencies of origins of replication were determined from differential values of polymerase tracks, similarly to (Daigaku *et al.* (2015)). Briefly, for all three datasets (Polδ, Pole, Polα), the difference of each neighbouring datapoint of polymerase track values (smoothed by simple moving average of 3) was calculated as $Diff_i = PT_i - PT_{i-1}$, where $Diff_i$ represents differential value at position i , and $PT_i - PT_{i-1}$ stand for smoothed polymerase track values at positions i and $i-1$, respectively. Differential value of the first bin on a given chromosome was assigned 0. Pole differentials and the opposites of Polδ and Polα differentials were averaged and smoothed by simple moving average of 3. Then, positive peaks (indicating sharp inclinations in the data) were selected. Differential peaks containing two or more distinct maxima separated by at least four bins were treated as independent peaks. Peaks with maxima below 30th percentile were disregarded. Each independent differential peak represented an origin of replication, the efficiency of which was estimated as 50% of the sum of its values. 259 replication initiation regions and 147 termination zones were selected using wild-type (WT) origin efficiency data. For comparison purposes, origin efficiencies were normalised assuming that the efficiency of the most efficient origin was 100%. Data were analysed in R (<https://www.R-project.org>; R Core Team, 2020) using a custom script (see *Software availability*).

Results

Brief overview of polymerase usage sequencing

Pu-Seq methodology determines the genome-wide polymerase activities by detecting the traces of rNMPs misincorporated by mutated Polδ (*cdc6^{L591G}*), Pole (*cdc20^{M630F}*) or Polα (*pol1^{L850F}*) (Daigaku *et al.*, 2015; Keszthelyi *et al.*, 2015). In Pu-Seq, respective polymerase mutant strains also carry a deletion of *rnh201*, the catalytic subunit of RNase H2 complex, disruption of which abrogates ribonucleotide excision repair (RER) and thus stabilises misincorporated rNMPs (Daigaku *et al.*, 2015). To assess activities of individual replicative polymerases, we employed a strategy previously used to analyse Okazaki fragment sequencing data (Petryk *et al.*, 2016). Briefly, activities of Polδ, Pole and Polα at any given locus are expressed as polymerase tracks, which are proportional differences of rNMPs misincorporated in the top and the bottom DNA strands (Figure 1).

Construction and characterisation of Polδ-overexpressing strains

To achieve an approximate two-fold and four-fold upregulation of the whole Polδ complex, we aimed to increase the genomic copy number of all four Polδ genes. We constructed a set of Cre-Lox integration vectors, each of which carried a distinct selection marker (NatR, KanR, *ura4⁺*) and all four genes constituting either WT (*cdc6⁺*, *cdc1⁺*, *cdc27⁺*, *cdm1⁺*) or L591G-mutated (*cdc6^{L591G}*, *cdc1⁺*, *cdc27⁺*, *cdm1⁺*) Polδ (Figure 2A). Employing Cre-Lox recombination mediated cassette exchange (Watson *et al.*, 2008), we generated three distinct Polδ genomic integrations and created strains carrying either one (2×Polδ) or three (4×Polδ) extra copies of either WT or L591G-mutated Polδ holoenzyme (Figure 2B). Using WT Polδ integrations, we constructed 2×Polδ and 4×Polδ strains expressing Cdc2^{asM17} (Cdk1 variant inhibitable by the ATP analogue 3-Br-PP1), which allowed us to synchronise cells in G2 and assess their progression through the S-phase (Aoi *et al.*, 2014). Additionally, we constructed 2×Polδ *rnh201Δ* and 4×Polδ *rnh201Δ* mutants expressing either Cdc20^{M630F} or Pol1^{L850F}, which allowed us to determine whether the activities of Pole and Polα were altered in cells over-expressing Polδ. In a similar manner, utilising L591G-mutated Polδ integrations, we produced 2×Polδ *rnh201Δ* and 4×Polδ *rnh201Δ* mutants exclusively expressing Cdc6^{L591G}, which allowed us to assess activity of Polδ at different expression levels.

To validate that 2×Polδ and 4×Polδ mutants displayed increased expression of Polδ genes, we measured relative transcript levels of *cdc1⁺*, *cdc27⁺*, *cdc6⁺/cdc6^{L591G}*, and *cdm1⁺* by RT-qPCR. In all genetic backgrounds tested, 2×Polδ and 4×Polδ mutants displayed a significant increase in relative transcript levels of all four Polδ genes (Figure 2C). Unfortunately, due to the unavailability of commercial antibodies recognising Polδ subunits in *S. pombe*, we were unable to confirm that protein levels of the Polδ subunits were also

Figure 1. General Pu-Seq analysis. Top panel – cartoon representation of Pol α , Pol δ and Pol ϵ activities around an origin of replication (Ori). Respective polymerase mutations employed in Pu-Seq are indicated. Middle panel – Example of genomic ribonucleotides (rNMPs; presented as 300bp bins) detected by Pu-Seq in *rnh201 Δ* cells expressing Cdc6^{L591G}, Cdc20^{M630F} and Pol1^{L850F}. A representative locus surrounding an origin of replication is shown. Bottom panel – Polymerase tracks calculated for Pol α , Pol δ and Pol ϵ at the representative locus. For each polymerase, polymerase tracks are calculated from rNMPs mapped to the top and the bottom DNA strands as: $PT = (R_T + R_B) / (R_T - R_B)$, where PT represents polymerase track, and R_T and R_B stand for rNMPs mapped to the top and the bottom DNA strands, respectively. Positive and negative values indicate predominant polymerase activity on the top and the bottom DNA strands, respectively. Data from 1 experiment are shown.

increased. It has been previously reported, however, that plasmid-based over-expression of each of the four Pol δ subunits is achievable in *S. pombe* (Kang *et al.*, 2000; MacNeill *et al.*, 1996; Reynolds *et al.*, 1998). Consequently, we reasoned that elevation of Pol δ transcript levels represented sufficient proof of *bona-fide* upregulation.

To determine the fundamental cellular consequences of Pol δ -overexpression, we assessed growth rate and cellular morphology of WT, 2 $\times$ Pol δ and 4 $\times$ Pol δ cells. Pol δ -overexpressing mutants displayed WT-like growth parameters and did not develop any cellular or nuclear defects (Figure 2D and 2E). Accordingly, increased Pol δ expression did not alter the distribution of cell sizes (Figure 2F). To assess whether increased Pol δ expression influenced progression through S-phase specifically, we synchronised WT, 2 $\times$ Pol δ and 4 $\times$ Pol δ cells with a Cdc2^{asM17} background in G2 by the addition of 3-Br-PP1 and analysed changes in DNA content in 15-min intervals after release. Progression through S-phase in 2 $\times$ Pol δ and 4 $\times$ Pol δ mutants was undistinguishable from WT cells (Figure 2F), suggesting that the over-production of Pol δ did not change S-phase progression. Taken together, a moderate increase in Pol δ expression did not have a notable impact on cell cycle or replication progression.

Replication dynamics

To investigate the potential influence of Pol δ -overexpression on replication dynamics in greater detail, we performed two

independent Pu-Seq experiments, each of which addressed activities of Pol δ , Pol ϵ and Pol α , in WT, 2 $\times$ Pol δ and 4 $\times$ Pol δ cells. Overall, in all genetic backgrounds tested, Pol δ , Pol ϵ and Pol α tracks displayed very little variation (Figure 3), suggesting that increased Pol δ levels did not dramatically alter the properties of replication. To capture a genome-wide view of replication [characterised by estimated firing efficiency (Or_{eff}) of at least 40%] and regions constituting replication termination zones, which were defined by two efficient origins ($Or_{eff} > 40\%$) and did not contain any intermediary efficiency origins ($20\% < Or_{eff} < 40\%$). Analysis of Pol δ and Pol ϵ tracks associated with 259 efficient origins and 147 termination zones did not reveal any notable differences (Figure 4A and 4B). We observed that Pol α tracks in 2 $\times$ Pol δ cells displayed marginal deviation from the WT profile (Figure 4A and 4B); however, considering that the observed difference was not reflected in 4 $\times$ Pol δ cells, we concluded this observation represented a technical, rather than biological phenomenon. We reasoned that if increased Pol δ levels negatively affected replisome progression, 2 $\times$ Pol δ and 4 $\times$ Pol δ mutants would be expected to display increased activity of low and intermediary efficiency origins. Pol δ -overexpressing cells, however, retained a WT-like distribution of genome-wide origin efficiencies, which further indicated normal replication progression (Figure 4C and 4D). Taken together, we concluded that, in our experimental system, a moderate increase in Pol δ levels did not result in any observable changes in replication dynamics.

Figure 2. Construction and characterisation of mutants over-expressing Polδ. (A) Simplified map of Cre-Lox vector(s) that were used to integrate extra copies of Polδ genes. Each vector carries genes constituting wild-type (WT) or L591G-mutated Polδ holoenzyme and one of three selection markers: NatR or KanR or *ura4*⁺. Cre – Cre recombinase (B) Graphical representation of genomic Polδ integration site(s) in 2×Polδ and 4×Polδ cells. (C) Relative transcript levels of Polδ genes (*cdc1*⁺, *cdc27*⁺, *cdc6*⁺/*cdc6*^{L591G} and *cdm1*⁺) in the indicated mutants measured by RT-qPCR. Mutants designated as *pol1*^{L850F}, *cdc6*^{L591G} and *cdc20*^{M630F} also carried *rh201Δ*. Individual points represent data from independent experiments. Horizontal lines represent means. Statistical significance was determined by the unpaired two-sample t-test. ** p ≤ 0.01; *** p ≤ 0.001; **** p ≤ 0.0001 (D) Representative growth curves of WT, 2×Polδ and 4×Polδ cells. Optical density (OD) was measured in 1h intervals for total 10 h. Time-series of log₂-transformed OD measurements are presented. Red lines represent linear regression models. Slopes of linear regression models (k) and calculated doubling times are indicated. (E) Representative images of WT, 2×Polδ and 4×Polδ cells stained with DAPI. Composite images of DIC and DAPI channels are shown. Scale bar represents 5 μm. (F) Distributions of cell lengths of WT, 2×Polδ and 4×Polδ cells. Data from three independent experiments are shown. Squares represent medians of individual experiments. Horizontal line represents the median of merged data. Statistical significance was determined by the unpaired two-sample Wilcoxon test. n.s. = not significant. (G) DNA profiles of WT, 2×Polδ and 4×Polδ cells synchronised in G2. Results from two independent experiments are shown.

Conclusions

In this study, we tested whether a moderate (2–4-fold) increase in Polδ expression impairs, or in any way alters, replication dynamics under normal conditions in *S. pombe*. The presented

experiments were inspired by report that a two-fold and four-fold increase in Polδ concentration reduces the rate of the leading strand synthesis *in vitro*, hypothesised to be due to stochastic polymerase switching, during which Polδ outcompetes Pole

Figure 3. Representative polymerase tracks. Polδ, Polε and Polα tracks across the right arm of chromosome III in WT, 2×Polδ and 4×Polδ cells. Means of two independent experiments are shown.

Figure 4. Pu-Seq analysis of mutants over-expressing Polδ. (A, B) Polδ, Polε and Polα tracks around 259 efficient origins of replication (A) and across 147 termination zones (B). Individual regions and means are shown. Polδ expression levels are indicated: circles – wild-type (WT); squares – 2×Polδ; triangles – 4×Polδ. Chromosomal coordinates around efficient origins were centred relative to the position of an origin. Data constituting termination zones were equally binned, and bins were centred relative to the midpoint of a termination zone. (C, D) Distribution of normalised origin efficiencies in WT, 2×Polδ and 4×Polδ cells.

and temporarily facilitates the extension of the leading strand (Yeeles *et al.*, 2017).

We constructed a set of strains carrying either one or three extra copies of all Polδ genes and validated that these Polδ

integrations resulted in increased transcription of the respective Polδ components: *cdc1*⁺, *cdc27*⁺, *cdc6*⁺ and *cdm1*⁺. We were unable to explore if the Polδ subunits were upregulated at the protein level. However, considering that successful ectopic over-production of Polδ subunits has been reported in the

seminal literature (Kang *et al.*, 2000; MacNeill *et al.*, 1996; Reynolds *et al.*, 1998), we argue that our experimental design conveyed a genuine Pol δ over-production.

We determined that cells characterised by up to four-fold increased Pol δ expression do not exhibit defects in growth and cell cycle progression. Furthermore, utilising Pu-Seq methodology, we demonstrated that genome-wide replication dynamics in 2xPol δ and 4xPol δ mutants is virtually indistinguishable from WT, arguing against the notion of stochastic polymerase switching or any other impairment of DNA replication induced by over-production of Pol δ .

Naturally, it is still possible that we simply did not reach the threshold of Pol δ expression that is required for the polymerase-switch to occur at frequencies detectable by Pu-Seq. Higher cellular levels of Pol δ could be achieved by ectopic or strong promoter-driven expression of Pol δ genes; however, we argue that such an extensive Pol δ over-production would constitute a non-physiological system, which would no longer be biologically relevant in relation to the reported *in vitro* data (Yeeles *et al.*, 2017). Moreover, it has been shown that gross over-expression of *cdc6*⁺ is detrimental to overall cell physiology (MacNeill *et al.*, 1996), which would likely make Pu-Seq experiments difficult to interpret or impossible to carry out. We also argue that promoter manipulation or plasmid-based over-expression would disrupt the stoichiometry of Pol δ subunits, which could be detrimental to Pol δ folding and function.

While we established that moderate over-expression of Pol δ does not noticeably affect canonical replication, we acknowledge that presented data do not sufficiently disprove the natural occurrence of the stochastic switch from Pole to Pol δ . Nevertheless, our data do imply that, if such events occur *in vivo*, they manifest at low frequencies and likely represent only a marginal disturbance to an overwhelmingly robust replication program.

Data availability

Underlying data

Gene Expression Omnibus: Raw and processed Pu-Seq data, Accession number GSE165503; [https://identifiers.org/geo:GSE165503](https://identifiers.org/geo/GSE165503).

Zenodo: Increased expression of Pol δ does not alter the canonical replication program *in vivo*. <https://doi.org/10.5281/zenodo.4513956>.

This project contains the following underlying data:

- Data_cell-size.xlsx (cell size measurements)
- data_OD.xlsx (optical density measurements)
- data_RT-qPCR.xlsx (raw Cq values)
- .tif files (raw microscopy images; strain-repeat-channel-image)
- .fcs files (flow cytometry files; well-strain-repeat-time-point)

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Software availability

Source code available from: https://github.com/R-Zach/Pu-Seq_polymerase_delta_over-expression

Archived source code at time of publication: <https://doi.org/10.5281/zenodo.4516546> (Zach, 2021)

License: [Apache License 2.0](#)

Acknowledgments

We would like to express our gratitude to Karel Naiman and other members of the Carr lab for their valuable suggestions and comments regarding experimental design and interpretation of the data.

References

- Aoi Y, Kawashima SA, Simanis V, *et al.*: **Optimization of the analogue-sensitive Cdc2/Cdk1 mutant by *in vivo* selection eliminates physiological limitations to its use in cell cycle analysis.** *Open Biol.* 2014; 4(7): 140063.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Aria V, Yeeles JTP: **Mechanism of Bidirectional Leading-Strand Synthesis Establishment at Eukaryotic DNA Replication Origins.** *Mol Cell.* 2018; 73(2): 199–211.e10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bähler J, Wu JQ, Longtine MS, *et al.*: **Heterologous modules for efficient and versatile PCR-based gene targeting in *Schizosaccharomyces pombe*.** *Yeast.* 1998; 14(10): 943–951.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bullock CR, Xing X, Shcherbakova PV: **DNA polymerase δ proofreads errors made by DNA polymerase ϵ .** *Proc Natl Acad Sci U S A.* 2020; 117(11): 6035–6041.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Burgers PMJ, Kunkel TA: **Eukaryotic DNA Replication Fork.** *Annu Rev Biochem.* 2017; 86(1): 417–438.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Clausen AR, Lujan SA, Burkholder AB, *et al.*: **Tracking replication enzymology *in vivo* by genome-wide mapping of ribonucleotide incorporation.** *Nat Struct Mol Biol.* 2015; 22(3): 185–191.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Daigaku Y, Keszthelyi A, Müller CA, *et al.*: **A global profile of replicative polymerase usage.** *Nat Struct Mol Biol.* 2015; 22(3): 192–198.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Donnianni RA, Zhou ZX, Lujan SA, *et al.*: **DNA Polymerase Delta Synthesizes Both Strands during Break-Induced Replication.** *Mol Cell.* 2019; 76(3): 371–381.e4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Feng W, D'Urso G: ***Schizosaccharomyces pombe* cells lacking the amino-terminal catalytic domains of DNA polymerase epsilon are viable but require the DNA damage checkpoint control.** *Mol Cell Biol.* 2001; 21(14): 4495–4504.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Garbacz MA, Lujan SA, Burkholder AB, *et al.*: **Evidence that DNA polymerase δ contributes to initiating leading strand DNA replication in *Saccharomyces***

cerevisiae. *Nat Commun*. 2018; **9**(1): 858.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Guilliam TA, Yeeles JTP: **An updated perspective on the polymerase division of labor during eukaryotic DNA replication**. *Crit Rev Biochem Mol Biol*. 2020a; **55**(5): 469–481.

[PubMed Abstract](#) | [Publisher Full Text](#)

Guilliam TA, Yeeles JTP: **Reconstitution of translesion synthesis reveals a mechanism of eukaryotic DNA replication restart**. *Nat Struct Mol Biol*. 2020b; **27**(5): 450–460.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Kang HY, Choi E, Bae SH, *et al.*: **Genetic analyses of *Schizosaccharomyces pombe* dna2(+) reveal that dna2 plays an essential role in Okazaki fragment metabolism**. *Genetics*. 2000; **155**(3): 1055–1067.

[PubMed Abstract](#) | [Free Full Text](#)

Kesti T, Flick K, Keränen S, *et al.*: **DNA polymerase epsilon catalytic domains are dispensable for DNA replication, DNA repair, and cell viability**. *Mol Cell*. 1999; **3**(5): 679–685.

[PubMed Abstract](#) | [Publisher Full Text](#)

Keszthelyi A, Daigaku Y, Ptasińska K, *et al.*: **Mapping ribonucleotides in genomic DNA and exploring replication dynamics by polymerase usage sequencing (Pu-seq)**. *Nat Protoc*. 2015; **10**(11): 1786–1801.

[PubMed Abstract](#) | [Publisher Full Text](#)

Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2**. *Nat Methods*. 2012; **9**(4): 357–359.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

MacNeill SA, Moreno S, Reynolds N, *et al.*: **The fission yeast Cdc1 protein, a homologue of the small subunit of DNA polymerase delta, binds to Pol3 and Cdc27**. *EMBO J*. 1996; **15**(17): 4613–4628.

[PubMed Abstract](#) | [Free Full Text](#)

Miyabe I, Kunkel TA, Carr AM: **The major roles of DNA polymerases epsilon and delta at the eukaryotic replication fork are evolutionarily conserved**. *PLoS Genet*. 2011; **7**(12): e1002407.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Miyabe I, Mizuno K, Keszthelyi A, *et al.*: **Polymerase δ replicates both strands after homologous recombination-dependent fork restart**. *Nat Struct Mol Biol*. 2015; **22**(11): 932–938.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Petersen J, Russell P: **Growth and the Environment of *Schizosaccharomyces pombe***. *Cold Spring Harb Protoc*. 2016; **2016**(3): pdb.top079764.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Petryk N, Kahli M, d'Aubenton-Carafa Y, *et al.*: **Replication landscape of the human genome**. *Nat Commun*. 2016; **7**: 10208.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Převorovský M, Oravcová M, Zach R, *et al.*: **CSL protein regulates transcription of genes required to prevent catastrophic mitosis in fission yeast**. *Cell Cycle*. 2016; **15**(22): 3082–3093.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

R Core Team: **R: A language and environment for statistical computing**.

R Foundation for Statistical Computing, Vienna, Austria. 2020.

[Reference Source](#)

Reynolds N, Watt A, Fantes PA, *et al.*: **Cdm1, the smallest subunit of DNA polymerase d in the fission yeast *Schizosaccharomyces pombe*, is non-essential for growth and division**. *Curr Genet*. 1998; **34**(4): 250–258.

[PubMed Abstract](#) | [Publisher Full Text](#)

Schneider CA, Rasband WS, Eliceiri KW: **NIH Image to ImageJ: 25 years of image analysis**. *Nat Methods*. 2012; **9**(7): 671–675.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sriramachandran AM, Petrosino G, Méndez-Lago M, *et al.*: **Genome-wide Nucleotide-Resolution Mapping of DNA Replication Patterns, Single-Strand Breaks, and Lesions by GLOE-Seq**. *Mol Cell*. 2020; **78**(5): 975–985.e7.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Watson AT, Garcia V, Bone N, *et al.*: **Gene tagging and gene replacement using recombinase-mediated cassette exchange in *Schizosaccharomyces pombe***. *Gene*. 2008; **407**(1–2): 63–74.

[PubMed Abstract](#) | [Publisher Full Text](#)

Yeeles JTP, Janska A, Early A, *et al.*: **How the Eukaryotic Replisome Achieves Rapid and Efficient DNA Replication**. *Mol Cell*. 2017; **65**(1): 105–116.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Zach R: **R-Zach/Pu-Seq polymerase_delta_over-expression: Release 1.0**. (Version v1.0). *Zenodo*. 2021.

<http://www.doi.org/10.5281/zenodo.4516546>

Zhou JC, Janska A, Goswami P, *et al.*: **CMG-Pol epsilon dynamics suggests a mechanism for the establishment of leading-strand synthesis in the eukaryotic replisome**. *Proc Natl Acad Sci U S A*. 2017; **114**(16): 4141–4146.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Zhou ZX, Lujan SA, Burkholder AB, *et al.*: **Roles for DNA polymerase δ in initiating and terminating leading strand DNA replication**. *Nat Commun*. 2019; **10**(1): 3992.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 18 March 2021

<https://doi.org/10.21956/wellcomeopenres.18297.r42830>

© 2021 Sabatinos S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Sarah Sabatinos

Department of Chemistry and Biology, Ryerson University, Toronto, Canada

Zach and Carr examine the effect of polymerase delta over-expression on fission yeast DNA replication. This study tests whether the *in vitro* results of a 2017 study (Yeeles *et al.*) are found in fission yeast DNA replication, *in vivo*. Yeeles (2017) found that increasing polymerase delta concentration caused a decrease in synthesis, attributed to polymerase competition and leading strand polymerase switch.

Zach and Carr have generated fission yeast strains that express additional copies of the 4 polymerase delta genes so that there are either 2 total copies or 4 total copies present in the genome. This design parallels the Yeeles *in vitro* work that similarly compared a 2x and 4x increase in polymerase delta.

The authors do not have antibody recognizing their polymerase delta proteins, but instead show that transcript levels of all 4 genes scale by qPCR. They characterize the effect of 2x, and 4x polymerase delta expression for its effects on: cell morphology, growth/doubling, and DNA content/cell cycle progression. Their calculation of growth uses absorbance readings, and is accompanied by cell length measurements to show that absorbance increase is not attributable to altered cell size. All results suggest that 2x and 4x increases in polymerase delta genes have no apparent effect on fission yeast cell fitness. Further, this over expression model does not noticeably slow DNA synthesis or cause replication-dependent effects on cell health and growth.

The methods are clearly written, and provide thorough descriptions of materials and equipment used. Interestingly, the authors used 2 separate methods for Pu-Seq library preparation. The impact of using 2 different methods is not described in the text, and data from 1 experiment is shown in Figure 1. The "...Means of two independent experiments are shown." in figure 3, and the data suggests that differences between prepared libraries is minimal. Indication of replicates would be helpful if not explicitly stated (elaborated below).

Overall, the authors provide a clear test of a specific hypothesis: that increased polymerase delta expression will impede synthesis. The data is clean, the premise is interesting, and the subject is

worthy of study. The issue of not seeing protein level to characterize polymerase delta levels is minor in this initial *in vivo* characterization, given the characterization of transcript levels.

Minor comments:

- The methods are well-described and clear. Some elaboration of replicates would improve the text to be very clear throughout. Could experimental replicate number be indicated for 2C? Figure 3 suggests that the means of the 2 separate Pu-Seq methods are very close- is this the case? Are both Pu-Seq library replicates used in Figure 4 calculations?
- If polymerase delta switching to the leading strand impacts replisome speed, could this be seen in polymerase epsilon tracks? Figure 3 suggests that while the patterns are very similar for Pol e under over expression conditions, the overlap may not be perfect; in contrast, there is apparently tight overlay for normal/2x/4x in polymerase delta and alpha tracks. Is a non-perfect overlap in polymerase epsilon tracks the residual effect of a transient situation? Since Figure 3 shows the mean of 2 independent experiments, is the variation lost by combining individual experiments (prepared differently)?
- I have no complaint with the use of qPCR to quantify expression of the polymerases delta components, and transcript levels appear to scale by ~2x and ~4x relative to the baseline amount of each, independent transcript. Based on known promoter activities, how does the authors' 2x or 4x level of transcript increase (from amplified copies driven by the native promoters) compare to ectopic promoter-driven expression from MacNeill (1996), Kang (2000) or Reynolds (1998)?
- How do efficient origins of replication in this work (Figure 4A) compare with other published studies?

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Cell biology, molecular genetics.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 19 Apr 2021

Antony Carr, University of Sussex, Falmer, Brighton, UK

- **The methods are well-described and clear. Some elaboration of replicates would improve the text to be very clear throughout. Could experimental replicate number be indicated for 2C? Figure 3 suggests that the means of the 2 separate Pu-Seq methods are very close- is this the case? Are both Pu-Seq library replicates used in Figure 4 calculations?**

In Figure 2C description, we included the number of independent measurements for WT, 2xPolδ and 4xPolδ cells. In Figure 1, Pu-Seq data from both experiments are shown now so reader can judge the variance of two methods used. In Figure 4, means of two experiments are analysed. This is now clearly stated in the Figure 4 description.

- **If polymerase delta switching to the leading strand impacts replisome speed, could this be seen in polymerase epsilon tracks? Figure 3 suggests that while the patterns are very similar for Pol ε under over expression conditions, the overlap may not be perfect; in contrast, there is apparently tight overlay for normal/2x/4x in polymerase delta and alpha tracks. Is a non-perfect overlap in polymerase epsilon tracks the residual effect of a transient situation? Since Figure 3 shows the mean of 2 independent experiments, is the variation lost by combining individual experiments (prepared differently)?**

We argued that, if replisome speed was affected, we would detect changes in origin firing. Origins efficiency profiles in WT and Polδ-overexpressing cells; however, are comparable (Figure 4 – C, D). It is true that, according to Figure 3, Polε tracks in WT and Polδ-overexpressing cells do not display perfect overlap. We believe this is due to slightly higher inter-experimental variability in WT Polε datasets, as is now indicated in revised Figure 1.

- **I have no complaint with the use of qPCR to quantify expression of the polymerases delta components, and transcript levels appear to scale by ~2x and ~4x relative to the baseline amount of each, independent transcript. Based on known promoter activities, how does the authors' 2x or 4x level of transcript increase (from amplified copies driven by the native promoters) compare to ectopic promoter-driven expression from MacNeill (1996), Kang (2000) or Reynolds (1998)?**

In all listed studies, ectopic over-expression of non-catalytic Polδ subunits is inferred indirectly, based on a positive effect on cellular fitness of various replication mutants including *cdc1-P13*, *cdc6-121*, *cdc27-P11*, *cdc24-M38* and *dna2-C2*. Successful over-expression of Cdc6 was judged by the fact that cells carrying a *nmt1-cdc6* construct developed severe cellular defects. Expression levels of Polδ subunits in different over-expression systems have never been demonstrated, so we cannot make a meaningful comparison.

- **How do efficient origins of replication in this work (Figure 4A) compare with other published studies?**

The original study by Daigaku *et al.* concluded that > 90% of origins identified by conventional method are also identified by Pu-Seq. Pu-Seq; however, provides better resolution and accuracy.

Competing Interests: No competing interests were disclosed.

Reviewer Report 09 March 2021

<https://doi.org/10.21956/wellcomeopenres.18297.r42833>

© 2021 Ren J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Jie Ren

CAS Key Laboratory of Genomic and Precision Medicine, Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing, China

This manuscript by Róbert Zach and Antony M. Carr examines the *in vivo* impact of an increased expression of DNA polymerase Pol δ on cellular growth and replication dynamics.

Intriguingly, although the canonical model divides the workload of leading strand synthesis and lagging strand synthesis between Pol ϵ and Pol δ , *in vitro* and *in vivo* studies showed evidence of Pol δ activity on both strands under certain circumstances. Yet, *in vitro* reconstitution of replication indicated that although increased dosage of Pol δ can out compete Pol ϵ on the leading strand, this is much less efficient for its synthesis. The current work tests the hypothesis that whether the same effect exist *in vivo*, which will provide more understanding on replication dynamics in living cells.

Using a carefully designed Cre-Lox system, the authors constructed strains with two- or four-fold increased Pol δ holoenzyme with the ability to incorporate rNTPs for following assessment of polymerase usage. With various analysis on cell growth and genome-wide quantification of individual polymerase usage especially at origins and termination regions, they concluded that there is no detectable impact in replication dynamics or associated growth defect caused by increased amounts of Pol δ .

Major Comments:

1. It will be nice to also show an example for termination site identification through Pu-seq as the demonstration for origin in Fig 1.
2. In the method section, the authors mentioned that an independent set of modified GLOE-seq was also performed along side Pu-seq. Are the GLOE-seq results comparable with those of Pu-seq? Is this work still undergoing?
3. Increased amount of Pol δ transcripts were confirmed with the over-expression system, but

increased usage of Pol δ is barely evident as shown by this work. Although not required for the current work, it will be nice to be followed up by testing whether or not more Pol δ subunits can be incorporated into the replisome using iPOND or similar techniques.

Minor points:

1. In the analysis of Pu-seq data, PT is calculated as $(R_t + R_b)/(R_t - R_b)$. Is there certain mathematical conversion omitted here? Otherwise, how is the range of PT kept within $[-1, 1]$ as in Fig 1?
2. It is concluded that Pol α tracks in 2xPol δ is marginally different from the other two conditions. I assume it is demonstrated by the mean polymerase tracks in Fig 4A and 4B as a slight deviation of the curve of squares from the other two curves. It is better to make that clear in the figure legend, or with some statistical analysis.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Molecular genetics.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 19 Apr 2021

Antony Carr, University of Sussex, Falmer, Brighton, UK

Major Comments:

- **It will be nice to also show an example for termination site identification through Pu-seq as the demonstration for origin in Fig 1.**

Representative locus in Figure 1 was expanded. Now the origin of replication as well as the

adjacent termination zone are displayed. Additionally, data from both experiments are shown.

- **In the method section, the authors mentioned that an independent set of modified GLOE- seq was also performed alongside Pu-seq. Are the GLOE-seq results comparable with those of Pu-seq? Is this work still undergoing?**

Modified GLOE-Seq protocol was used to prepare the second set of Pu-Seq libraries (experiment/repeat 2). Figure 1 now shows Pu-Seq data generated by both procedures.

- **Increased amount of Polδ transcripts were confirmed with the over-expression system, but increased usage of Polδ is barely evident as shown by this work. Although not required for the current work, it will be nice to be followed up by testing whether or not more Polδ subunits can be incorporated into the replisome using iPOND or similar techniques.**

This is a valuable suggestion. It indeed would be interesting to investigate how replisome composition reacts to different levels of its particular components, such as Polδ.

Minor points:

- **In the analysis of Pu-seq data, PT is calculated as $(R_t + R_b) / (R_t - R_b)$. Is there certain mathematical conversion omitted here? Otherwise, how is the range of PT kept within [-1,1] as in Fig 1?**

In the manuscript we made a mistake and stated $PT = (R_T + R_B) / (R_T - R_B)$ instead of $PT = (R_T - R_B) / (R_T + R_B)$, which is the correct equation used in the analysis. Typo was corrected. The fact that denominator is always bigger than nominator ensures that PT falls within [-1,1].

- **It is concluded that Polα tracks in 2xPolδ is marginally different from the other two conditions. I assume it is demonstrated by the mean polymerase tracks in Fig 4A and 4B as a slight deviation of the curve of squares from the other two curves. It is better to make that clear in the figure legend, or with some statistical analysis.**

Minor deviations in Polα tracks are now signified by asterisks.

Additional changes to the manuscript:

- In the introduction section, we now reference a recent paper discussing the role of Polδ in bypassing oxidative DNA lesions:

"Additionally, it has been demonstrated that Polδ takes over the leading strand synthesis and performs an error-free bypass of oxidative DNA adducts thymine glycol and 8-oxoguanine (Guilliam & Yeeles, 2021)."

Competing Interests: No competing interests were disclosed.