OPEN LETTER

Defining the burden of febrile illness in rural South and Southeast Asia: an open letter to announce the launch of the Rural Febrile Illness project [version 1; peer review: 3 approved]

Arjun Chandna ^{1,2}, Nan Shwe Nwe Htun³, Thomas J. Peto ^{1,3}, Marco Liverani⁴⁻⁶, Tobias Brummaier ^{1,7-9}, Koukeo Phommason¹⁰, Sazid Ibna Zaman ^{1,3}, Aye Sandar Zaw¹¹, Elizabeth Batty ^{1,3}, Naomi Waithira ^{1,3}, Melissa Richard-Greenblatt ¹², Stuart D. Blacksell¹, Ladaporn Bodhidatta³, James J. Callery ³, Watcharintorn Fagnark³, Shayla Islam¹³, Sanchai Lertcharoenchoke³, Mavuto Mukaka^{1,3}, Tiengkham Pongvongsa¹⁴, William H.K. Schilling ^{1,3}, Janjira Thaipadungpanit ^{3,15}, Rupam Tripura ^{1,3}, Arjen M. Dondorp ^{1,3}, Mayfong Mayxay ^{1,10,16}, Nicholas J. White ^{1,3}, François Nosten ^{1,7}, Frank Smithuis ^{1,11}, Elizabeth A. Ashley ^{1,10}, Richard J. Maude ^{1,3,17,18}, Nicholas P.J. Day ^{1,3}, Yoel Lubell ^{1,3}

¹Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, UK

²Cambodia Oxford Medical Research Unit, Angkor Hospital for Children, Siem Reap, Cambodia

³Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand

⁴Department of Global Health and Development, London School of Hygiene & Tropical Medicine, London, UK

⁵Faculty of Public Health, Mahidol University, Bangkok, Thailand

⁶School of Tropical Medicine and Global Health, Nagasaki University, Nagasaki, Japan

⁷Shoklo Malaria Research Unit, Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Mae Sot, Thailand

⁸Swiss Tropical and Public Health Institute, Basel, Switzerland

⁹University of Basel, Basel, Switzerland

¹⁰Lao-Oxford-Mahosot Hospital Wellcome Trust Research Unit, Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao People's Democratic Republic

¹¹Myanmar Oxford Clinical Research Unit, Yangon, Myanmar

¹²University of Pennsylvania, Pennsylvania, USA

¹³Health Nutrition and Population Programme, BRAC, Dhaka, Bangladesh

¹⁴Savannakhet Provincial Health Department, Savannakhet, Lao People's Democratic Republic

¹⁵Department of Clinical Tropical Medicine, Mahidol University, Bangkok, Thailand

¹⁶Institute of Research and Education Development, University of Health Sciences, Vientiane, Lao People's Democratic Republic

¹⁷Harvard TH Chan School of Public Health, Harvard University, Boston, USA

¹⁸The Open University, Milton Keynes, UK

First published: 26 Mar 2021, 6:64

<https://doi.org/10.12688/wellcomeopenres.16393.1>

Latest published: 10 Mar 2022, 6:64

Open Peer Review

<https://doi.org/10.12688/wellcomeopenres.16393.2>

Abstract

In rural areas of South and Southeast Asia malaria is declining but febrile illnesses still account for substantial morbidity and mortality. Village health workers (VHWs) are often the first point of contact with the formal health system, and for patients with febrile illnesses they can provide early diagnosis and treatment of malaria. However, for the majority of febrile patients, VHWs lack the training, support and resources to provide further care. Consequently, treatable bacterial illnesses are missed, antibiotics are overused and poorly targeted, and patient attendance wanes along with declining malaria.

This *Open Letter* announces the start of a new initiative, the Rural Febrile Illness (RFI) project, the first in a series of projects to be implemented as part of the South and Southeast Asian Community-based Trials Network (SEACTN) research programme. This multi-country, multi-site project will begin in Bangladesh, Cambodia, Lao PDR, and Myanmar and will define the epidemiological baseline of febrile illness in five remote and underserved areas of Asia where malaria endemicity is declining and access to health services is limited. The RFI project aims to determine the incidence, causes and outcomes of febrile illness; understand the opportunities, barriers and appetite for adjustment of the role of VHWs to include management of non-malarial febrile illnesses; and establish a network of community healthcare providers and facilities capable of implementing interventions designed to triage, diagnose and treat patients presenting with febrile illnesses within these communities in the future.

Keywords

Community Health Workers, Etiology, Fever, Primary Health Care, Rural Health, Southeastern Asia, Telemedicine, Western Asia, Village Health Workers

This article is included in the [Mahidol Oxford Tropical Medicine Research Unit \(MORU\)](#) gateway.

Approval Status

	1	2	3
version 2
(revision)			view
10 Mar 2022
version 1
26 Mar 2021	view	view	view

1. **Samuel Akech** , KEMRI-Wellcome Trust Research Programme, Nairobi, Kenya
2. **Teresa B. Kortz** , University of California, San Francisco (UCSF), San Francisco, USA
3. **Emily Ciccone** , University of North Carolina, Chapel Hill, USA

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding authors: Arjun Chandna (arjun@tropmedres.ac), Yoel Lubell (yoel@tropmedres.ac)

Author roles: **Chandna A:** Conceptualization, Funding Acquisition, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Shwe Nwe Htun N:** Conceptualization, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Peto TJ:** Conceptualization, Methodology, Writing – Review & Editing; **Liverani M:** Conceptualization, Methodology, Writing – Review & Editing; **Brummaier T:** Methodology, Writing – Review & Editing; **Phommasone K:** Methodology, Writing – Review & Editing; **Ibna Zaman S:** Methodology, Visualization, Writing – Review & Editing; **Sandar Zaw A:** Methodology, Writing – Review & Editing; **Batty E:** Methodology, Writing – Review & Editing; **Waithira N:** Data Curation, Methodology, Software, Writing – Review & Editing; **Richard-Greenblatt M:** Methodology, Writing – Review & Editing; **Blacksell SD:** Methodology, Writing – Review & Editing; **Bodhidatta L:** Methodology, Writing – Review & Editing; **Callery JJ:** Methodology, Writing – Review & Editing; **Fagnark W:** Methodology, Writing – Review & Editing; **Islam S:** Methodology, Writing – Review & Editing; **Lertcharoenchoke S:** Methodology, Software, Writing – Review & Editing; **Mukaka M:** Methodology, Writing – Review & Editing; **Pongvongsa T:** Methodology, Writing – Review & Editing; **Schilling WHK:** Methodology, Writing – Review & Editing; **Thaipadungpanit J:** Methodology, Writing – Review & Editing; **Tripura R:** Methodology, Writing – Review & Editing; **Dondorp AM:** Funding Acquisition, Methodology, Writing – Review & Editing; **Mayxay M:** Funding Acquisition, Methodology, Writing – Review & Editing; **White NJ:** Funding Acquisition, Methodology, Writing – Review & Editing; **Nosten F:** Funding Acquisition, Methodology, Writing – Review & Editing; **Smithuis F:** Funding Acquisition, Methodology, Writing – Review & Editing; **Ashley EA:** Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing; **Maude RJ:** Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing; **Day NPJ:** Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing; **Lubell Y:** Conceptualization, Funding Acquisition, Methodology, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work is supported by Wellcome [219644]. The work in Cambodia is conducted as part of an RAI3E OR project, funded by the Global Fund [QSE-M-UNOPS-MORU-20864-007-42]. This diagnostics component of this partner project in Cambodia is supported by the Wellcome Trust.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Chandna A *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Chandna A, Shwe Nwe Htun N, Peto TJ *et al.* **Defining the burden of febrile illness in rural South and Southeast Asia: an open letter to announce the launch of the Rural Febrile Illness project [version 1; peer review: 3 approved]** Wellcome Open Research 2021, 6:64 <https://doi.org/10.12688/wellcomeopenres.16393.1>

First published: 26 Mar 2021, 6:64 <https://doi.org/10.12688/wellcomeopenres.16393.1>

Disclaimer

The views expressed in this article are those of the authors. Publication in Wellcome Open Research does not imply endorsement by Wellcome.

Introduction

The majority of individuals in South and Southeast Asia live in rural areas, often characterised by high levels of poverty and restricted access to healthcare^{1–3}. Data on causes of disease in these areas to prioritise interventions for scale up are limited. Despite this, there are indications that diseases of an infectious aetiology, ‘febrile illnesses’, account for substantial morbidity and mortality^{4,5}.

Malaria is the quintessential febrile illness. For decades, empiric antimalarials were recommended for patients presenting with fever⁶, reflective of the burden associated with this fatal but treatable illness. In many parts of Asia, village health workers (VHWs) were introduced to improve access to treatment. Rapid declines in malaria incidence have subsequently been observed⁷. As a result, today fever seldom means malaria⁸, and the exact cause of the illness often remains unknown, as does optimal management and what becomes of these individuals⁹.

Falling malaria incidence highlights the inadequacies and fragilities of this vertical approach to healthcare. VHWs receive limited training, support and remuneration, and once malaria has been ruled out, often cannot provide further testing or treatment for their patients. Consequently, treatable bacterial infections are missed^{10,11}, and antibiotics, if available, are overused and poorly targeted^{11–13}. Furthermore, the current inability to address non-malarial fever adversely affects uptake of malaria testing: patients have little to gain from having malaria ruled out in areas where it is already rare, but no other care offered for their illness^{14,15}. This hampers eradication efforts and risks resurgence of drug-resistant malaria^{16,17}. Encouragingly, it has been shown that the trend of decreasing patient attendance (and malaria testing) can be reversed when the remit of the VHW is extended to other basic healthcare provision¹⁴.

This *Open Letter* announces the start of the Rural Febrile Illness (RFI) project, the first project in the South and Southeast Asian Community-based Trials Network (SEACTN) research programme¹⁸. The RFI project will begin in Bangladesh, Cambodia, Lao PDR and Myanmar and aims to define the epidemiological baseline of febrile illness in remote and underserved areas of Asia where malaria endemicity has declined and access to health services is limited. The primary objective of the RFI project is to determine the incidence, causes and outcomes of febrile illness in the rural communities residing within the project areas. The longer-term objective of SEACTN is to establish a network of community healthcare providers and facilities, capable of implementing interventions designed to triage, diagnose and treat patients presenting with febrile illnesses and other causes of ill health within these communities in the future.

Project overview

The RFI project is a multi-country, multi-site initiative comprised of a number of prospective observational studies. The project is divided into three key Work Packages (WP) with parallel supporting activities (Table 1). Detailed protocols for individual components of the RFI project are available at www.seactn.org.

Work Package A (WP-A)

In collaboration with VHWs we will develop and deploy electronic data collection tools to capture the incidence, presenting syndromes and outcomes of febrile illness amongst individuals presenting to the most peripheral level of the health system. In Lao PDR, community healthcare providers working at primary health centres (PHCs) rather than VHWs will implement the project, as utilisation of VHWs is currently low in this area².

Data collected from VHWs and PHCs will be complemented by parallel health status and health seeking behaviour surveys, verbal autopsies, and village and health facility mapping projects, designed to gain a comprehensive understanding of the burden of febrile illness and access to care in the areas served by these health facilities and providers. We will also perform targeted aetiological investigations in a subset of patients presenting to WP-A providers and facilities.

Work Package B (WP-B)

WP-B activities will be concentrated around two higher-level health facilities (health clinics and/or hospitals) in each of the sites in Bangladesh, Laos PDR and Myanmar, located within (or nearby) the same geographical area as the VHWs and PHCs selected for WP-A. More extensive aetiological investigations, as well as assays of host biomarkers, will be performed in patients with febrile illnesses attending these facilities.

Work Package C (WP-C)

In WP-C we will draw on the data collected in WP-A and WP-B to create temporally- and spatially-explicit electronic decision-support tools (eDSTs), designed to assist community health workers (both VHWs and healthcare providers at PHCs) in their assessment, triage and treatment of patients presenting with febrile illnesses in rural and remote areas. The data from WP-A and WP-B will also be used to identify the most high-impact and cost-effective point-of-care tests (POCTs) that could be included within the eDSTs, as well as appropriate delivery mechanisms. Subsequent deployment and health system integration will be informed by stakeholder analyses in the same settings to better understand the opportunities, barriers and appetite for adjustment of the role of VHWs and other community healthcare providers to include management of non-malarial causes of febrile illness.

Study sites and implementing partners

Strong, long-standing partnerships have been leveraged to plan and implement this multi-disciplinary project, across at least

Table 1. Key objectives of the RFI project.

Overall	
1	Determine the incidence, causes and outcomes of febrile illness in five rural areas of Bangladesh, Cambodia, Lao PDR and Myanmar
2	Establish a network of community healthcare providers and facilities, capable of implementing interventions designed to triage, diagnose and treat patients presenting with febrile illnesses within these communities in the future
Work Package A	
1	Develop electronic data collection tools for patients presenting to village health workers and primary health centres with febrile illnesses in the study areas
2	Determine the incidence and outcomes of febrile illnesses amongst patients presenting to village health workers and primary health centres in the study areas
3	Describe and understand health status and health-seeking behaviour for febrile illnesses in the study areas
4	Describe common causes of mortality and events immediately preceding death in the study areas
5	Map the geographic location, accessibility, treatment availability and workforce capacity of health facilities within and nearby the study areas
Work Package B	
1	Describe the causes and outcomes of febrile illnesses amongst patients presenting to sentinel rural health facilities in the study areas
2	Determine the diagnostic performance of host biomarkers to distinguish bacterial from viral infections amongst patients presenting with febrile illnesses
3	Determine the prognostic performance of host biomarkers to identify patients with febrile illnesses at risk of severe outcomes
Work Package C	
1	Complete stakeholder analyses to identify the opportunities, barriers and appetite for adjustment of the role of community healthcare providers to include management of non-malarial causes of febrile illness
2	Model the cost-effectiveness of different combinations of interventions to improve the management of febrile illness in the study areas
3	Develop and pilot electronic decision-support tools and point-of-care tests that can assist community healthcare providers in their assessment, triage and treatment of patients with febrile illnesses

620 villages in five rural regions of Bangladesh, Cambodia, Lao PDR and Myanmar (Figure 1).

In Bangladesh the project will be implemented in partnership with BRAC, one of the largest non-governmental development organisations in the world. As of September 2020, 130 villages have been selected and local health facilities (Upazila Health Complexes and District Hospitals) identified in Cox's Bazar and Bandarban districts.

The project in Cambodia will be a collaboration with the National Center for Parasitology, Entomology and Malaria Control (CNM), the Provincial Health Departments of Battambang and Pailin provinces, and Action for Health Development (AHEAD), a civil society organisation working to improve health in rural communities. The study will take place in western Cambodia in 120 villages from Pailin and Battambang provinces.

Two study sites are planned in Myanmar. In Northern Kachin state, 89 villages and two health clinics (Himalaya Clinic I and II), have been selected across three townships (Putao, Machanbaw and Nawngmun), where Medical Action Myanmar (MAM), a non-governmental healthcare organisation, operates a network of VHWs and rural health clinics. The second site is in Kayah state, where the Shoklo Malaria Research Unit (SMRU) supervises a large community health worker programme focused on malaria elimination. Overall, 130 villages and two health clinics (Wang Pha and Mawker Thai) across the townships of Hlaingbwe, Myawaddy and Kawkaik are planned for selection.

In southern Lao PDR, 28 PHCs (serving 160 villages) within three districts (Atsaphangthong, Thaphalanxay and Phin), as well as Atsaphangthong district hospital and Savannakhet provincial hospital have been selected. The project will be conducted in partnership with Savannakhet Provincial Health Office (SPHO).

Figure 1. Study areas with selected villages and health facilities. AHEAD, Action for Health Development; MAM, Medial Action Myanmar; SMRU, Shoklo Malaria Research Unit; SPHO, Savannakhet Provincial Health Office; WP-B, Work Package B. Due to access difficulties related to the ongoing COVID-19 pandemic, participating villages in the townships of Hlaingbwe and Kawkaireik (Thai-Myanmar border region) and WP-B health facilities in Bangladesh are yet to be finalised. WP-B activities are currently not planned in Cambodia and village selection is underway.

WP-A

Electronic data collection tools

Electronic data collection forms are being developed using the CommCare platform (Cambridge, MA, USA) and loaded on to mobile Android tablets. The tablets, accompanied by solar chargers where necessary, will be distributed to all VHWs (PHCs in Lao PDR) in SEACTN villages. Their feedback will inform iteration of the data collection forms following a human-centred design approach¹⁹. Once finalised, the tablets will be used to capture data from patients with febrile illnesses who attend the VHWs and PHCs (see below).

Data will be securely uploaded in real-time. In areas without mobile internet uploads will occur periodically during visits by the local implementing partner (BRAC, AHEAD, MAM, SMRU or SPHO). The aim is to develop a data stream from source to web-based interface with near real-time geospatial mapping of the incidence and outcomes of febrile illness. A detailed Data Management Plan is available on request from the Mahidol-Oxford Tropical Medicine Research Unit (MORU) Data Access Committee.

Digital health education materials

Training materials are being developed in partnership with [DigitalMedic](#) and serve three primary purposes:

- To sensitise communities to the project aims and objectives (see below);
- To support in-person training of the VHWs and PHC workers, focussing on ensuring accurate syndromic classification of febrile illnesses, reliable collection of clinical data and biological samples, and confidence in operating the electronic data collection tool;
- To improve patient understanding of the health problems the project aims to address.

Community engagement activities

Prior to the initiation of the study the local implementing partner will meet with villagers, village leaders and relevant local authorities to explain the project rationale, aims and planned activities, using digital educational material as outlined

above. Community consent will be sought for the collection of basic demographic and syndromic data (using the same electronic data collection tool) from all patients attending VHWs and PHCs (both febrile and non-febrile patients) to better understand the health needs of the community and inform future directions of the SEACTN.

Determining the incidence and outcomes of febrile illnesses in patients presenting to village health workers and primary health centres

All patients who attend the VHW (or PHC in Lao PDR) will be screened, and those with a febrile illness who provide consent will be enrolled. Enrolment will be consecutive and is planned for 18 months. Based on current attendances we estimate that we will capture approximately 100,000 febrile illness episodes across the five study sites.

The VHW (or PHC worker) will use the electronic data collection tool to record a limited set of clinical data and the result of the malaria rapid diagnostic test (mRDT). No changes to patient management will occur as a result of the study, except that the health worker will be prompted to consider referral and/or inform the operations team of the implementing partner if any danger signs are elicited, subject to the specific context at each site²⁰⁻²². The participant will be followed-up by the VHW 28 days later to determine the outcome of their illness.

In participants outside the neonatal age range a capillary blood sample (finger- or heel-prick) will be collected at the same time as the mRDT and stored as a dried blood spot (DBS) on filter paper. In a subset of participants (at least 20,000 illness episodes) a convalescent DBS will be collected at the one-month follow-up. Targeted molecular and serological investigations (Table 2) will be performed on an initial set of 10,000 illness

Table 2. Aetiological investigations to be performed in the RFI project, including specimen type, target pathogen, diagnostic platform and laboratory location²⁴⁻²⁶. *Dengue NS1 testing will be performed on the acute (D0) sample only. **Blood cultures will be performed in-country at a quality-assured laboratory close to the WP-B health facility. RDT, rapid diagnostic test; JE, Japanese Encephalitis; NPS, nasopharyngeal swab.

Setting	Specimen	Target pathogen	Laboratory location and diagnostic platform		
			MORU PCR (Day 0)	MORU Paired serology (Day 0 and Day 28)	On site (Day 0)
WP-A: VHWs and PHCs	Capillary blood	<i>Plasmodium</i> spp.			RDT
		Dengue	✓	NS1* and IgM	
		Chikungunya	✓	IgM	
		Pan-Alphavirus	✓		
		Pan-Flavivirus	✓	Zika and JE IgM	
		<i>Orientia tsutsugamushi</i>		IgM	
		<i>Rickettsia typhi</i>		IgM	
		<i>Rickettsia</i> spp.		IgM	
		<i>Leptospira</i> spp.		IgM	
WP-B: Rural health facilities	Capillary blood	<i>Plasmodium</i> spp.			RDT or microscopy
	Venous blood	Dengue	✓	NS1* and IgM	
		Chikungunya	✓	IgM	
		Pan-Alphavirus	✓		
		Pan-Flavivirus	✓	Zika and JE IgM	
		<i>Orientia tsutsugamushi</i>	✓	IgM	
		<i>Rickettsia</i> spp.	✓	IgM	
		<i>Leptospira</i> spp.	✓	IgM	
		16S rRNA (eubacteria)	✓		
		Bacterial bloodstream infections			Blood culture**
	NPS	Respiratory pathogens	✓		

episodes to determine the aetiology of the febrile illness, with remaining samples stored for future targeted analyses.

Health-seeking behaviour surveys

We will conduct health status and health-seeking behaviour surveys to understand pathways to care for individuals with febrile illnesses. A better appreciation of how care for acute illness is currently sought in these settings will enable us to determine the overall burden of febrile illness in the study areas, and identify the best options for potential intervention in the future.

Community-level data, such as estimated vaccine coverage, availability of water, sanitation and hygiene (WASH) facilities, and assessments of indoor air quality, will be gathered to help contextualise the project's findings.

Verbal autopsies

Little information exists on the febrile (and non-febrile) causes of mortality in the study areas²³. In collaboration with researchers from the University of Toronto, we have adapted the World Health Organization's (WHO) Verbal Autopsy (VA) tool to conduct electronic VA questionnaires for all deaths that occur in SEACTN villages in Bangladesh, Cambodia, Lao PDR and the North Kachin site in Myanmar²⁷. Understanding the common causes of mortality and the circumstances that surround death will enable us to identify targets for interventions that can be implemented and evaluated within the SEACTN programme in the future.

Village and health facility mapping

A thorough understanding of local healthcare infrastructure is essential to direct referrals to higher-level care appropriately. Villages, transport networks and health facilities in the study areas will be mapped using field collection and satellite imagery, and detailed profiles created of the study villages including population statistics, communication and transport systems, health services and campaigns, physical environment and socioeconomic metrics. WHO's [AccessMod 5](#) software will be used to estimate travel time from the study villages to health facilities and potential gaps in local service provision identified. In addition to informing how future interventions can be deployed within the SEACTN, we will provide this information to health system planners and policy makers to help identify where new health facilities could be provided and existing services strengthened to achieve highest impact.

WP-B

Determining the aetiologies of febrile illnesses in patients attending rural health facilities

To gain a comprehensive understanding of the causes of febrile illness in the region, we will recruit a cohort of patients attending two sentinel health facilities within four of the project areas. The information from these cohorts will be combined with the aetiological data from WP-A, where we are limited to collecting low-volume DBS specimens from a subset of participants (the remote locations preclude collection of samples from all villages and prevent maintenance of a cold chain).

All patients aged > 28 days who attend the health facilities with a febrile illness will be screened, and those who provide consent will be enrolled. Six hundred participants (inpatients and outpatients) will be enrolled in each age stratum (> 28 days to < 5 years; ≥ 5 years to < 15 years; ≥ 15 years), across both health facilities within a single project area (i.e. 1,800 participants in each area). Recruitment is planned for a minimum 12 continuous months at each site to ensure seasonality is adequately captured.

Baseline data will be recorded including demographics, anthropometrics, presenting syndrome, vital signs, clinical signs, duration of illness and any care sought for the illness thus far. In Bangladesh, Lao PDR and the Kayin state site in Myanmar blood cultures will be collected and the results provided to the treating clinical teams.

Venous blood samples and nasopharyngeal swabs will be collected and transported to MORU's central laboratories in Bangkok, where molecular and serological aetiological investigations ([Table 2](#)) will be performed. Novel molecular techniques such as target enrichment sequencing will also be validated against existing molecular diagnostics and used to investigate a broader range of pathogens than can be detected by the pathogen-specific tests²⁸. Additional investigations for participants with specific syndromes (for example, urine, pus, throat swab and/or cerebrospinal fluid culture) will be considered, subject to feasibility at each of the facilities.

Admitted participants will be followed-up daily during their admission. All participants will be followed-up on day 2 (in person or via telephone) and asked to return to the health facility 28 days after enrolment to determine the outcome of their illness. A venous blood sample will be collected at this time for convalescent serological testing.

Identification of clinical features and host biomarkers that distinguish bacterial from viral infections in febrile patients

The venous blood samples collected from patients attending the rural health facilities will also be used to quantitatively measure host biomarkers that reflect immune activation and endothelial dysfunction. We will use this information, together with the baseline clinical data and results of the aetiological investigations, to construct diagnostic algorithms that can distinguish bacterial from viral infections^{29–32}. Host biomarkers feasible for measurement using POCTs, will be prioritised, so that the algorithms could be used to guide antimicrobial prescribing in resource-limited primary care settings.

Development of prognostic clinical prediction models for febrile patients at risk of severe outcomes

We will also use the host biomarker data, in conjunction with the baseline clinical and day 2 and 28 outcome data, to derive prognostic algorithms (clinical prediction models) to identify patients at risk of severe outcomes, which could be used to guide referral decisions from community healthcare settings to higher-level care^{33–38}. The baseline clinical data, host biomarker panels and outcome definitions have been harmonised with a

parallel study (NCT04285021) to facilitate data sharing and external validation of the prediction models.

WP-C

Stakeholder mapping and analyses

Throughout the RFI project, key stakeholders will be engaged and interviewed on the topic of expanding the remit of VHWs and/or other community healthcare providers to include management of non-malarial febrile illnesses. Stakeholders will include policy makers and managers within the Ministries of Health of the respective countries; representatives of international and national donor organisations; individuals and organisations responsible for the implementation and supervision of VHW and other community-based health programmes; and VHWs and community members.

During the interviews, information and views will be collected about operational challenges, opportunities and policy bottlenecks concerning the expansion of VHW programmes, with particular attention to the key issues of health system integration, capacity, and sustainability. After an initial set of interviews with seed informants, selected after a stakeholder mapping exercise, additional interviewees will be identified through snowball sampling. Collected material will be analysed thematically, while interim outputs such as matrix tables or position maps will be constructed to better understand the role and interest of different categories of stakeholders, their resources,

and their level of support to the expansion of VHW programmes. Findings will also inform participatory development of the eDSTs (see below).

Economic evaluation of interventions that could be deployed in the SEACTN to improve the management of febrile illness

We will draw on the data and findings from WP-A and WP-B to develop economic models to assess the cost-effectiveness of different (combinations of) interventions to improve the management of febrile illness, that could be trialled in subsequent projects within the SEACTN. Analyses will be conducted on standalone interventions such as the implementation of POCTs³⁹, as well as multi-layered approaches integrating regional data on causes of illness within eDSTs, alongside POCTs⁴⁰. The outputs of the cost-effectiveness analyses will also be provided to relevant governmental and non-governmental actors to support the planning and scale-up of national programmes.

Development of eDSTs for use by rural community healthcare providers

The results from these modelling assessments and the data from WP-A and WP-B, will be used to design bespoke, spatially-explicit eDSTs that can assist community healthcare providers in their assessment, triage and treatment of patients with febrile illnesses (Figure 2).

Figure 2. Overview of the long-term ambition for the management of febrile illnesses within SEACTN. (1) Patient with febrile illness presents to village health worker (VHW). (2) Simple clinical algorithm pre-loaded on mobile device helps VHW assess patient. (3) VHW performs point-of-care test (POCT) if recommended by algorithm. (4) VHW decides between community-based treatment or referral for higher-level care. (5) Regional health facilities (sentinel nodes) periodically provide data from patients attending with febrile illnesses (including patients referred by VHWs working within SEACTN). (6) Proportion of samples collected from febrile patients at the time POCTs are performed, stored on filter paper and transported to reference laboratories. (7) Data from (5) and (6) integrated with data from (2) to periodically update clinical algorithms to reflect seasonal and longitudinal changes in febrile illness landscape.

The eDSTs will be loaded on to the mobile tablets that the VHWs and PHC workers are already familiar with using for data collection. Interactive training modules will be developed (in partnership with DigitalMedic), and will include educational content on the common causes of febrile illness in the region, as well as instructional information on how to use the new eDSTs and any relevant POCTs. User feedback will be sought and the eDSTs iterated using a human-centred design process¹⁹. Once finalised the eDSTs will be put forward for evaluation in future projects to be conducted within the SEACTN.

Dissemination of findings

Interim findings from relevant aspects of the project (for example, aggregate results of the aetiological investigations for febrile illness and the VA study) will be periodically summarised and fed back to the local communities via village leaders and local authorities. This will allow important information to be actioned in a time-sensitive manner, whilst preserving the confidentiality of individual participants.

The final results generated from this study will be disseminated to key stakeholders (identified during the stakeholder mapping exercises) and the study communities (via the same community engagement forums used to launch the project) in both English and local languages. The results will also be shared with the scientific community via peer-reviewed publications and conference presentations.

Limitations

Studying febrile illness at the most peripheral level of a health system provides a unique opportunity to influence the course of a patient's illness at their first contact with formal health services. This is particularly important in settings where regulation of facilities, providers and treatments is often lacking or inadequately enforced. However, working at this level of the health system also poses certain challenges.

We are limited to collecting low-volume DBS specimens from patients attending VHWs and PHC workers. The aetiological yield of these specimens may be low. However, they are feasible for collection in large numbers, and without attempts to understand the causes of fever in rural areas of the region, meaningful improvements in the management of febrile illness will likely remain elusive. To mitigate this risk, we will recruit cohorts of patients attending sentinel health facilities within these areas (WP-B), where collection of a wider range of

specimens will permit more extensive aetiological investigations. Furthermore, the yield from the DBS specimens will be monitored and reviewed by the RFI Study Management Group. Depending on the results, DBS assaying may be expanded or replaced with an alternate strategy.

Expansion of the role of VHWs is both feasible and impactful¹⁴. However there is a limit to the number of roles that these skilled yet lay-people can be expected to fulfil, without adequate recognition, supervision and remuneration⁴¹. Stakeholder engagement, planned throughout the RFI project, will be crucial to understand the feasibility of long-term adjustments to the VHW role.

Conclusion

The RFI project aims to better understand and quantify the burden of febrile illness, the aetiological causes and the manner in which it affects people living in some of the most underserved areas of South and Southeast Asia, all on a scale which has not been attempted before. We will collect information to better understand and predict the outcomes of patients with febrile illnesses based on a multitude of factors, which will form the basis for interventions within the SEACTN in the future.

The foundational infrastructure established by the RFI project will include a network of upskilled and supported community healthcare providers, user-friendly electronic data collection tools, functioning biological specimen collection, transport and diagnostic pipelines, and robust data management systems for near real-time geospatial mapping of the incidence and outcomes of patients with febrile illness. SEACTN will be well positioned to support ongoing surveillance of febrile illnesses in the region, enabling earlier detection of disease outbreaks and the regular updating of treatment algorithms in response to seasonal and longitudinal changes in the regional febrile illness landscape.

Data availability

No data are associated with this article.

Acknowledgements

The authors are grateful for the support from the local authorities in participating countries and to our partners and collaborators. We thank the SEACTN Expert Review Group for helpful advice on the design of the project.

References

1. National Institute of Population Research and Training (NIPORT) and ICF: **Bangladesh Demographic and Health Survey 2017-18: Key Indicators**. Dhaka, Bangladesh and Rockville, Maryland, USA. 2019. [Reference Source](#)
2. Lao Statistics Bureau and UNICEF: **Lao Social Indicator Survey II 2017, Survey Findings Report**. Vientiane, Lao PDR. 2018. [Reference Source](#)
3. Ministry of Health and Sports (MoHS) and ICF: **Myanmar Demographic and Health Survey 2015-16**. Nay Pyi Taw Myanmar and Rockville, Maryland, USA. 2017. [Reference Source](#)
4. Crump JA, Kirk MD: **Estimating the Burden of Febrile Illnesses**. *PLoS Negl Trop Dis*. 2015; **9**(12): e0004040. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

5. Shrestha P, Dahal P, Ogbonnaa-Njoku C, *et al.*: **Non-malarial febrile illness: a systematic review of published aetiological studies and case reports from Southern Asia and South-eastern Asia, 1980-2015.** *BMC Med.* 2020; **18**(1): 299. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. World Health Organization: **Guidelines for the treatment of malaria.** Geneva, Switzerland, 2006. [Reference Source](#)
7. Landier J, Parker DM, Thu AM, *et al.*: **Effect of generalised access to early diagnosis and treatment and targeted mass drug administration on Plasmodium falciparum malaria in Eastern Myanmar: an observational study of a regional elimination programme.** *Lancet.* 2018; **391**(10133): 1916–1926. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. World Health Organization: **World Malaria Report.** Geneva, Switzerland, 2017. [Reference Source](#)
9. WHO: **WHO informal consultation on fever management in peripheral health care settings: a global review of evidence and practice.** 2013; 78. [Reference Source](#)
10. Mayxay M, Castonguay-Vanier J, Chansamouth V, *et al.*: **Causes of non-malarial fever in Laos: a prospective study.** *Lancet Glob Health.* 2013; **1**(1): e46–e54. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Lubell Y, Blacksell SD, Dunachie S, *et al.*: **Performance of C-reactive protein and procalcitonin to distinguish viral from bacterial and malarial causes of fever in Southeast Asia.** *BMC Infect Dis.* 2015; **15**: 511. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Wangrangsimakul T, Althaus T, Mukaka M, *et al.*: **Causes of acute undifferentiated fever and the utility of biomarkers in Chiangrai, northern Thailand.** *PLoS Negl Trop Dis.* 2018; **12**(5): e0006477. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Hopkins H, Bruxvoort KJ, Cairns ME, *et al.*: **Impact of introduction of rapid diagnostic tests for malaria on antibiotic prescribing: analysis of observational and randomised studies in public and private healthcare settings.** *BMJ.* 2017; **356**: j1054. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. McLean ARD, Wai HP, Thu AM, *et al.*: **Malaria elimination in remote communities requires integration of malaria control activities into general health care: an observational study and interrupted time series analysis in Myanmar.** *BMC Med.* 2018; **16**(1): 183. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Lubell Y, Chandna A, Smithuis F, *et al.*: **Economic considerations support C-reactive protein testing alongside malaria rapid diagnostic tests to guide antimicrobial therapy for patients with febrile illness in settings with low malaria endemicity.** *Malar J.* 2019; **18**(1): 442. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Ashley EA, Dhorda M, Fairhurst RM, *et al.*: **Spread of artemisinin resistance in Plasmodium falciparum malaria.** *N Engl J Med.* 2014; **371**(5): 411–23. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Hamilton WL, Amato R, van der Pluijm RW, *et al.*: **Evolution and expansion of multidrug-resistant malaria in southeast Asia: a genomic epidemiology study.** *Lancet Infect Dis.* 2019; **19**(9): 943–951. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. UK Wellcome Trust: **Innovations Flagships: funded projects.** (accessed 1 October 2020 2020). [Reference Source](#)
19. Adam M, McMahon SA, Prober C, *et al.*: **Human-Centered Design of Video-Based Health Education: An Iterative, Collaborative, Community-Based Approach.** *J Med Internet Res.* 2019; **21**(1): e12128. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. WHO: **Integrated Management of Childhood Illnesses.** 2014. [Reference Source](#)
21. World Health Organization: **Acute Care: Integrated Management of Adolescent and Adult Illness.** Geneva, Switzerland, 2004. [Reference Source](#)
22. WHO: **WHO/UNICEF Joint Statement - Integrated Community Case Management: an equity-focused strategy to improve access to essential treatment services for children.** 2012. [Reference Source](#)
23. Rao C: **Mortality estimates for South East Asia, and INDEPTH mortality surveillance: necessary but not sufficient?** *Int J Epidemiol.* 2013; **42**(4): 1196–9. [PubMed Abstract](#) | [Publisher Full Text](#)
24. Smit PW, Elliott I, Peeling RW, *et al.*: **An overview of the clinical use of filter paper in the diagnosis of tropical diseases.** *Am J Trop Med Hyg.* 2014; **90**(2): 195–210. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Woo PCY, Lau SKP, Teng JLL, *et al.*: **Then and now: use of 16S rDNA gene sequencing for bacterial identification and discovery of novel bacteria in clinical microbiology laboratories.** *Clin Microbiol Infect.* 2008; **14**(10): 908–34. [PubMed Abstract](#) | [Publisher Full Text](#)
26. Leber AL, Everhart K, Daly JA, *et al.*: **Multicenter Evaluation of BioFire FilmArray Respiratory Panel 2 for Detection of Viruses and Bacteria in Nasopharyngeal Swab Samples.** *J Clin Microbiol.* 2018; **56**(6): e01945–17. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. World Health Organization: **Verbal autopsy standards: ascertaining and attributing causes of death.** 2016. [Reference Source](#)
28. Goh C, Golubchik T, Ansari MA, *et al.*: **Targeted metagenomic sequencing enhances the identification of pathogens associated with acute infection.** *bioRxiv.* 2019. [Publisher Full Text](#)
29. Kapasi AJ, Ditttrich S, González IJ, *et al.*: **Biomarkers for Distinguishing Bacterial from Non-Bacterial Causes of Acute Febrile Illness: A Comprehensive Review.** *PLoS One.* 2016; **11**(8): e0160278. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Keitel K, Kagoro F, Samaka J, *et al.*: **A novel electronic algorithm using host biomarker point-of-care tests for the management of febrile illnesses in Tanzanian children (e-POCT): A randomized, controlled non-inferiority trial.** *PLoS Med.* 2017; **14**(10): e1002411. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Do NTT, Ta NTD, Tran NTH, *et al.*: **Point-of-care C-reactive protein testing to reduce inappropriate use of antibiotics for non-severe acute respiratory infections in Vietnamese primary health care: a randomised controlled trial.** *Lancet Glob Health.* 2016; **4**(9): e633–e41. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Althaus T, Greer RC, Swe MMM, *et al.*: **Effect of point-of-care C-reactive protein testing on antibiotic prescription in febrile patients attending primary care in Thailand and Myanmar: an open-label, randomised, controlled trial.** *Lancet Glob Health.* 2019; **7**(1): e119–e31. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Leligedowicz A, Conroy AL, Hawkes M, *et al.*: **Markers Of Endothelial Injury And Immune Activation Effectively Risk-Stratify Acute Febrile Syndromes In African Children.** *Am J Resp Crit Care Med.* Washington; 2020; **201**: A2818. [Reference Source](#)
34. Robinson ML, Workneh M, Ditttrich S, *et al.*: **Host biomarkers to predict the severity of acute febrile illness: A scoping review.** *medRxiv.* 2019. [Publisher Full Text](#)
35. Wright SW, Lovelace-Macon L, Hantrakun V, *et al.*: **sTREM-1 predicts mortality in hospitalized patients with infection in a tropical, middle-income country.** *BMC Med.* 2020; **18**(1): 159. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Richard-Greenblatt M, Boillat-Blanco N, Zhong K, *et al.*: **Prognostic Accuracy of Soluble Triggering Receptor Expressed on Myeloid Cells (sTREM-1)-based Algorithms in Febrile Adults Presenting to Tanzanian Outpatient Clinics.** *Clin Infect Dis.* 2020; **70**(7): 1304–1312. [PubMed Abstract](#) | [Publisher Full Text](#)
37. George EC, Walker AS, Kiguli S, *et al.*: **Predicting mortality in sick African children: the FEAST Paediatric Emergency Triage (PET) Score.** *BMC Med.* 2015; **13**: 174. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. McDonald CR, Weckman A, Richard-Greenblatt M, *et al.*: **Integrated fever management: disease severity markers to triage children with malaria and non-malarial febrile illness.** *Malar J.* 2018; **17**(1): 353. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Lubell Y, Althaus T, Blacksell SD, *et al.*: **Modelling the Impact and Cost-Effectiveness of Biomarker Tests as Compared with Pathogen-Specific Diagnostics in the Management of Undifferentiated Fever in Remote Tropical Settings.** *PLoS One.* 2015; **11**(3): e0152420. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Chandna A, White LJ, Pongvongsa T, *et al.*: **Accounting for aetiology: can regional surveillance data alongside host biomarker-guided antibiotic therapy improve treatment of febrile illness in remote settings? [version 2; peer review: 2 approved].** *Wellcome Open Res.* 2019; **4**: 1. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Pallas SW, Minhas D, Perez-Escamilla R, *et al.*: **Community Health Workers in Low- and Middle-Income Countries: What Do We Know About Scaling Up and Sustainability?** *Am J Pub Health.* 2013; **103**(7): e74–82. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 30 April 2021

<https://doi.org/10.21956/wellcomeopenres.18033.r43501>

© 2021 Ciccone E. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Emily Ciccone

Division of Infectious Diseases, School of Medicine, University of North Carolina, Chapel Hill, NC, USA

This open letter describes plans for a large, multi-country project focused on describing the etiology and improving the management of febrile illness in rural areas of South and Southeast Asia. This ambitious work is incredibly important as the widespread availability of malaria testing and treatment has led to a decline in malaria incidence, leaving most febrile illnesses undifferentiated. This diagnostic uncertainty leads to inappropriate and inadequate treatment, including the overuse of antibiotics, which drives antimicrobial resistance, a major threat to global public health.

In this article, the authors clearly outline their objectives and plans for implementation of the initiative, and the maps in Figure 1 distinctly show the varied locations of the included study sites. I also appreciated their acknowledgment of the challenges of working in rural areas with VHW providers, and think that there was sufficient explanation and justification of their plans. The continual reassessment of the feasibility of expanding the VHW role through stakeholder engagement itself will provide interesting information that will shape the evolution of such programs in response to the changing epidemiology of the conditions that VHW treat.

A few minor comments related to each Work Package are as follows:

WP-A

1. Please clarify the patient population that will be included. As the VHWs will be the primary providers that are included in the study, I assume the patient population will be children under 5 years of age, but would recommend clearly stating the target study population and if there are any differences in the age groups that will be enrolled at each site.
2. How will follow-up assessments be done? In other words, will they be in-person visits or phone calls? I assume in-person for at least a portion as convalescent DBS are being collected, but some clarification here would be useful.

3. How was 28 days chosen for the follow-up interval? This seems to be quite long for acute febrile illnesses. A brief justification of how this interval was chosen would be helpful.
4. Will the health-seeking behaviour surveys be conducted for all participants in WP-A or a subset? If a subset, how with the sampling be done?

WP-B

1. Again, a bit more clarification about age group would be helpful here - will any adults be enrolled or does the ≥ 15 age group only extend up to 18 years of age?
2. Would be interested in more detail regarding the host biomarker testing. Specifically, which tests are planned? Will the results of POCT be available to providers in real time?

WP-C

1. I found Figure 2 to be a bit confusing, particularly the group of icons in the center. Perhaps it would be helpful to include a few more arrows demonstrating the progression of the initial assessment (presentation --> mobile algorithm --> POCT --> location of care decision).

A few stylistic/grammatical things to consider are as follows:

- Could delete word "residing" in last paragraph of the introduction as communities themselves don't res
- Spell out BRAC when it is first mentioned in the Study sites and implementing partners section.
- Should be "local facilities have been identified" in same paragraph
- Remove the parentheses around "combinations of" in the paragraph about economic evaluation of interventions in WP-C section.

Is the rationale for the Open Letter provided in sufficient detail?

Yes

Does the article adequately reference differing views and opinions?

Yes

Are all factual statements correct, and are statements and arguments made adequately supported by citations?

Yes

Is the Open Letter written in accessible language?

Yes

Where applicable, are recommendations and next steps explained clearly for others to follow?

Not applicable

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: I am an early career physician scientist primarily focused on improving the management of pediatric febrile illness in sub-Saharan Africa (specifically Uganda and Malawi). My current projects study the use of rapid diagnostics, including biomarker and pathogen-specific testing, at peripheral health centers with limited laboratory capacity and by VHWs to improve antibiotic stewardship among children with acute respiratory illness.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 14 Dec 2021

Arjun Chandna, University of Oxford, Oxford, UK

WP-A: Please clarify the patient population that will be included. As the VHWs will be the primary providers that are included in the study, I assume the patient population will be children under 5 years of age, but would recommend clearly stating the target study population and if there are any differences in the age groups that will be enrolled at each site.

For WP-A, there are no age restrictions in the inclusion criteria, which will be uniform at all sites.

The relevant sentence has been amended to reflect this: "Enrolment is age-unrestricted, will be consecutive, and is planned for 18 months at all sites."

WP-A: How will follow-up assessments be done? In other words, will they be in-person visits or phone calls? I assume in-person for at least a portion as convalescent DBS are being collected, but some clarification here would be useful.

As far as possible, follow-up assessments will be conducted in person, and participants will have their travel costs reimbursed for attendance at follow-up visits. However, given the remoteness of some villages, telephone follow-up will be done if participants are not able to present in person.

The following has been added to clarify this point: "As far as possible, follow-up will be conducted face-to-face, with participants reimbursed for travel costs. However, given the remoteness of some of the study villages, follow-up may be conducted by telephone if the participant finds it too onerous to attend in person."

WP-A: How was 28 days chosen for the follow-up interval? This seems to be quite long for acute febrile illnesses. A brief justification of how this interval was chosen would

be helpful.

This interval was chosen because many of the diagnostic investigations involve paired serology, thus an adequate amount of time needs to have elapsed to allow for clinically significant changes in antibody titres. Furthermore, as many of the villages are relatively inaccessible, participants may find it too onerous to re-attend for follow-up if the interval is short. The preceding sentences have been added verbatim to the manuscript to explain our reasoning.

WP-A: Will the health-seeking behaviour surveys be conducted for all participants in WP-A or a subset? If a subset, how with the sampling be done?

The following has been added to the section on Household Health Surveys to address this query: "A cross-sectional household survey will be conducted targeting villages participating in WP-A, with households residing in the study area eligible for the survey. Two-stage cluster-randomized sampling of the households will be performed. In the first stage, villages of each study site will be selected; and in the second stage households located in the selected villages will be selected. All members of the selected households will be invited to participate in the survey. Participants will not be selected based on presence or absence of symptoms or diseases, and participation in WP-A does not affect the eligibility for survey participation."

WP-B: Again, a bit more clarification about age group would be helpful here - will any adults be enrolled or does the ≥ 15 age group only extend up to 18 years of age?

Yes, adults will be enrolled.

To avoid misunderstanding, the relevant sentence has been amended to "Six hundred participants (inpatients and outpatients) will be enrolled in each age stratum (>28 days to <5 years; ≥ 5 years to <15 years; and ≥ 15 years with no upper age limit), across all participating health facilities within a single project area (i.e. 1,800 participants in each area)."

WP-B: Would be interested in more detail regarding the host biomarker testing. Specifically, which tests are planned? Will the results of POCT be available to providers in real time?

We have added the following paragraph to the section on biomarkers, which addresses test selection and availability of POCT results: "Host biomarkers will be selected on the basis of biological plausibility, systematic review of the literature, and the results of ongoing prospective studies. The aim is to identify and test biomarkers that are implicated in the final common pathways to sepsis and severe febrile illness and shared across a range of microbial aetiologies. In community settings, it is not possible to obtain an aetiological diagnosis for the vast majority of patients with febrile illness during routine clinical care and, hence, our aim is to develop 'pathogen agnostic' diagnostic and prognostic algorithms

that can be applied at the syndromic level by limited-skill health workers and be used to inform common clinical management decisions e.g., whether to prescribe an antibiotic, which antibiotic to prescribe, and whether the patient requires referral to higher-level care. Biomarkers will be measured retrospectively using central laboratory-based assays (ELISAs and multi-analyte panels). The results will not be available to providers in real time during the initial observational phases of the project."

WP-C: I found Figure 2 to be a bit confusing, particularly the group of icons in the center. Perhaps it would be helpful to include a few more arrows demonstrating the progression of the initial assessment (presentation --> mobile algorithm --> POCT --> location of care decision).

We have added some additional arrows to the group of icons in the centre to clarify the expected sequence of events during a patient encounter. We hope that this, in combination with the legend of the figure, will provide a clear infographic summary of the overall conceptual framework for the medium- to long-term ambition of SEACTN.

General: Could delete word "residing" in last paragraph of the introduction as communities themselves don't res (sic)

'Residing' has been changed to 'located'.

General: Spell out BRAC when it is first mentioned in the Study sites and implementing partners section.

BRAC is now the official name of the organization and is not an acronym. We have reflected this in the manuscript.

General: Should be "local facilities have been identified" in same paragraph.

This sentence has now been modified for clarity.

General: Remove the parentheses around "combinations of" in the paragraph about economic evaluation of interventions in WP-C section.

The sentence has been amended to "We will draw on the data and findings from WP-A and WP-B to develop economic models to assess the cost-effectiveness of different interventions, or combinations of interventions, to improve the management of febrile illness suitable for assessment via clinical trials in subsequent projects within the SEACTN."

Competing Interests: I have no competing interests to declare

Reviewer Report 23 April 2021

<https://doi.org/10.21956/wellcomeopenres.18033.r43499>

© 2021 Kortz T. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Teresa B. Kortz

Department of Pediatrics, University of California, San Francisco (UCSF), San Francisco, CA, USA

Overall, a clear, well-written letter. This is an important topic and the proposed initiative has the potential to positively impact child health.

I may have missed it, but how is this being funded and what is the timeline?

Abstract

- Clear, no comments.

Introduction

- Suggest describing village health workers some more. For example, who are they, what role do they play, how do they improve access to treatment...
- Are the declines in malaria incidence related to VHWs? That is how I interpreted the second paragraph.

Project overview

- What is the proposed timeline for this project and the various work packages?

Study sites and implementing partners

- Was there a site selection process and if so, what was it? Was there an effort to recruit a certain type or number of community?
- How is "febrile illness" defined for the sake of study inclusion?
- Which "host biomarkers feasible for measurement using POCTs, will be prioritised"?

Dissemination/Limitations/Conclusion

- No specific comments.

Is the rationale for the Open Letter provided in sufficient detail?

Yes

Does the article adequately reference differing views and opinions?

Yes

Are all factual statements correct, and are statements and arguments made adequately supported by citations?

Yes

Is the Open Letter written in accessible language?

Yes

Where applicable, are recommendations and next steps explained clearly for others to follow?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: sepsis, severe febrile illness, pediatric critical care in resource-variable settings, host prognostic biomarkers, fever etiology, resource utilization

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 14 Dec 2021

Arjun Chandna, University of Oxford, Oxford, UK

Introduction: Suggest describing village health workers some more. For example, who are they, what role do they play, how do they improve access to treatment...

Please see the response above to the similar question asked by Reviewer 1, which addresses this comment in detail.

Introduction: Are the declines in malaria incidence related to VHWs? That is how I interpreted the second paragraph.

This is partly correct – VHWs were introduced as part of wider malaria control programmes, but were not the sole reason for the decline in malaria incidence.

We have clarified this point as follows: "In many parts of Asia, village health workers (VHWs) and/or village malaria workers (VMWs) were introduced to improve access to treatment as part of wider malaria eradication programmes. Rapid declines in malaria incidence have subsequently been observed."

Project overview: What is the proposed timeline for this project and the various work packages?

The project has an anticipated completion date of June 2023. Given site-specific circumstances, especially COVID-19 restrictions, sites will commence recruitment for the

work packages at different times, with the first site in Laos having started recruitment for WP-A in August 2021.

This has now been included as follows: "Sites will commence recruitment for WP-A and WP-B at different times given their unique circumstances, in particular the Covid-19 control measures in place at each site. WP-A will commence at the first site in August 2021, and we anticipate completing all work packages by June 2023. Detailed protocols for individual components of the RFI project are available at www.seactn.org."

Study sites and implementing partners: Was there a site selection process and if so, what was it? Was there an effort to recruit a certain type or number of community?

We have mentioned that the SEACTN project leverages on strong, long-standing partnerships, but have elaborated on this further to include details on site and village selection by adding the following sentence: "Site selection was based on these pre-existing collaborative relationships and village selection was delegated to the partner organizations, who are best placed to know the needs of, and challenges facing, the communities they serve."

How is "febrile illness" defined for the sake of study inclusion?

Acute febrile illness is defined as a documented fever ($\geq 37.5^{\circ}\text{C}$ axillary) or hypothermia ($< 35.5^{\circ}\text{C}$) at presentation and/or history of fever in the last 24 hours. This has now been added to the manuscript.

Which "host biomarkers feasible for measurement using POCTs, will be prioritised"?

Please see response to the comment on biomarkers from Reviewer 3 below, which answers this question comprehensively.

Competing Interests: I have no competing interests to declare.

Reviewer Report 20 April 2021

<https://doi.org/10.21956/wellcomeopenres.18033.r43297>

© 2021 Akech S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Samuel Akech

Health Services Unit, KEMRI-Wellcome Trust Research Programme, Nairobi, Kenya

The open letter communicates about a series of planned studies in several study sites in Asia to define aetiology of febrile illnesses, develop and implement point of care tests (POCTs) to differentiate bacterial versus viral infections, develop electronic decision support (eDST), and implement the POCTs and eDSTs using community health workers (village health workers and primary health care workers) for referral and treatment. In this letter, the authors have summarised linked pieces of research projects that are grouped into 3 work packages with several specific objectives. It is therefore expected that sufficient detail would be missing in a project that has several components. However, the ultimate success of this project as described would be a successful implementation of the eDST, POCTs, and referral/treatment by the Community Health Workers (CHWs).

Comments:

A brief description of training and capabilities of CHWs in the study settings will assist readers to gain insights into prospects of successful implementation. What other tasks do the CHWs already have? Will there be detailed implementation science conducted to understand implementation challenges for the project and is there a conceptual framework for the project implementation? It would be good to briefly mention how the implementation is expected to work at the level of CHWs who are quite central. Are the CHWs remunerated, what is their motivation, how much are researchers working with policy makers in the study countries-some insights will be good. Are the village health workers collecting heel/thumb prick samples? Is this something that they are permitted to do in this setting-this could be briefly explained.

Are there examples of treatments that CHWs would be allowed to give based on results of POCTs? If POCTs suggest a bacterial infection, it may be harmful to give an antibiotic and send the patient home since causes of a positive bacterial test could be varied e.g., UTI, sepsis, intra-abdominal infections (especially in adults), and it is difficult to imagine an antibiotic that can be given by CHWs that would cover all potential causes without need for a referral. In such cases, eDST would be configured to trigger a referral or no referral.

It would seem more logical if aetiological study and development of POCTs were done at earlier phases and then promising POCTs implemented plus eDST using CHWs. Can a diagram for sequence of activities be included?;

Is the rationale for the Open Letter provided in sufficient detail?

Yes

Does the article adequately reference differing views and opinions?

Yes

Are all factual statements correct, and are statements and arguments made adequately supported by citations?

Yes

Is the Open Letter written in accessible language?

Yes

Where applicable, are recommendations and next steps explained clearly for others to follow?

Not applicable

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Malaria, sepsis, clinical trials, paediatrics, epidemiology, severe childhood illness

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 14 Dec 2021

Arjun Chandna, University of Oxford, Oxford, UK

A brief description of training and capabilities of CHWs in the study settings will assist readers to gain insights into prospects of successful implementation. What other tasks do the CHWs already have? Will there be detailed implementation science conducted to understand implementation challenges for the project and is there a conceptual framework for the project implementation? It would be good to briefly mention how the implementation is expected to work at the level of CHWs who are quite central. Are the CHWs remunerated, what is their motivation, how much are researchers working with policy makers in the study countries-some insights will be good.

There is a high degree of variability in the roles, capabilities, and remuneration of village health workers, even within the same country. We have provided further details on these points:

“There is considerable variation in the training and capabilities of VHWs, and even within the same country there are different types of VHWs, funded by different actors and managed by different institutions. For example, in Lao PDR one such category of VHW is the Village Health Volunteer (VHV), who are managed by the Department of Hygiene and Health Promotion and receive no stipend. They undergo a 10-day training course to support health education, hygiene and sanitation programmes, vaccination, maternal health care, nutrition, the prevention and control of infectious diseases, and the management of revolving drug funds. In contrast, Lao VMWs are tasked with the diagnosis and treatment of uncomplicated malaria cases, referrals of severe case, distribution of bed nets, and health education/promotion but, unlike VHVs, receive a monthly stipend, usually financed through the Global Fund.”

Implementation research will, indeed, be conducted to identify and understand the relevant challenges and opportunities beginning with the stakeholder analysis component of the project, details of which are provided later in the open letter.

Are the village health workers collecting heel/thumb prick samples? Is this something that they are permitted to do in this setting-this could be briefly explained.

Yes, they are – this has been approved by the relevant ethics committees. All participating village health workers will be trained in study procedures prior to site activation.

The following sentence has been added to clarify this point: “VHWs/VMWs will be trained in dried blood spot collection prior to patient enrolment, for which the relevant site-specific ethical approvals have been obtained.”

Are there examples of treatments that CHWs would be allowed to give based on results of POCTs? If POCTs suggests a bacterial infection, it may be harmful to give an antibiotic and send the patient home since causes of a positive bacterial test could be varied e.g., UTI, sepsis, intra-abdominal infections (especially in adults), and it is difficult to imagine an antibiotic that can be given by CHWs that would cover all potential causes without need for a referral. In such cases, eDST would be configured to trigger a referral or no referral.

As mentioned, the ultimate aim of WP-C is to develop bespoke electronic algorithm-based decision support tools covering both diagnosis and management. To avoid management decisions based solely on POCT results, the algorithms will have the ability to aid primary health workers in formulating syndromic diagnoses and selecting the antibiotic regimen most likely to be of clinical benefit. POCTs will be but one component in these algorithms, and distinguishing bacterial from viral infection merely one of the several uses envisaged for them.

We have modified the relevant sentences to reflect this: “These eDSTs will be designed to assist community health workers (both VHWs and healthcare providers at PHCs) in their assessment, triage, and treatment of patients presenting with febrile illnesses in rural and remote areas. The data from WP-A and WP-B will also be used to identify the most high-impact and cost-effective point-of-care tests (POCTs) that could be included within the eDSTs, as well as appropriate delivery mechanisms.”

“POCTs will be but one component of the eDSTs, which will be dynamic, algorithm-based, and developed using artificial intelligence principles using the rich datasets resulting from WP-A and WP-B.”

It would seem more logical if aetiological study and development of POCTs were done at earlier phases and then promising POCTs implemented plus eDST using CHWs. Can a diagram for sequence of activities be included?

Yes, this is indeed the sequence of activities in this project – WP-A and WP-B will run concurrently. In these work packages, the causes of febrile illness will be defined, potential POCTs evaluated, and eDSTs developed. WP-C will focus on the implementation of these eDSTs, which will incorporate the use of POCTs found to be promising.

We have added the following to the Project Overview section, which sets out the timeline succinctly: "In brief, WP-A and WP-B, will define the causes of febrile illness in the region, evaluate potential point-of-care tests (POCTs), and develop electronic decision support tools (eDSTs). WP-C will focus on the implementation of these eDSTs, which will incorporate the use of POCTs found to be promising."

Competing Interests: No competing interests to declare.