RESEARCH ARTICLE

Adherence to iron and folic acid supplementation during pregnancy among postnatal mothers seeking maternal and child healthcare at Kakamega level 5 hospital in Kenya: a cross-sectional study [version 1; peer review: 2 approved with reservations]

Felix Bahati ^{1,2}, Salome Kairu-Wanyoike³, Japheth Mativo Nzioki ²

¹Health Services Research Unit, KEMRI Wellcome Trust, Nairobi, Nairobi, 43640-00100, Kenya

²Environmental Health and Disease Control, Jomo Kenyatta University of Agriculture and Technology, Nairobi, Nairobi, 62 000 – 00200, Kenya

³Directorate of Veterinary Services, Ministry of Agriculture, Livestock, Fisheries and Cooperatives, Nairobi, Nairobi, Kangemi 00605, Kenya

V1 First published: 12 Apr 2021, 6:80
<https://doi.org/10.12688/wellcomeopenres.16699.1>
 Latest published: 05 Jul 2021, 6:80
<https://doi.org/10.12688/wellcomeopenres.16699.2>

Abstract

Background: Maternal anaemia is a public health problem worldwide, and its aetiology is linked to iron deficiency. The high nutrient demand during pregnancy exacerbates the condition. To meet the increased nutritional demand, supplementation of iron and folic acid (IFA) is key. The supplements are provided freely to pregnant women during antenatal visits at public health facilities, however, their uptake and adherence in Kenya remain unacceptably low.

Methods: A hospital-based cross-sectional study involving 241 postnatal mothers seeking maternal and child healthcare (MCH) care at Kakamega level 5 hospital was conducted. Both quantitative and qualitative data were collected. Quantitative data were collected from 241 eligible postnatal mothers, while qualitative data were obtained through key informant interviews with community health volunteers and healthcare providers.

Results: There was a moderate adherence to IFA supplementation (60.6%) during pregnancy among postnatal mothers seeking MCH care at Kakamega level 5 hospital. Some of the reasons for non-adherence stated by the respondents included; IFA related side effects (41.3%), forgetfulness (37.3%) and bad smell of the IFA supplements (10.3%). Higher IFA adherence was noted among the primigravida participants (OR=2.704; 95% CI: 1.262, 5.793; p=0.010) compared to multigravida participants, and those with a higher knowledge level of anaemia (OR=3.215; 95% CI: 1.346, 7.68; p=0.009) compared to their

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
05 Jul 2021		
version 1		
12 Apr 2021		

1. **Katherine L. Dickin** , Cornell University, Ithaca, USA

2. **Martin N. Mwangi** , University of Malawi, Blantyre, Malawi

Any reports and responses or comments on the article can be found at the end of the article.

counterparts with low anaemia knowledge. Other factors that showed correlation with IFA adherence were: IFA education, pregnancy counselling before conception and the number of antenatal care visits attained.

Conclusion: There is a moderate adherence to IFA supplementation during pregnancy among mothers seeking MCH at Kakamega level 5 hospital. The greatest impediments of IFA compliance during pregnancy are IFA side effects, forgetfulness and the bad smell of the IFA tablets. Therefore, providing IFA education to pregnant mothers incorporated with probable ways of managing the IFA side effects would contribute to IFA supplementation adherence.

Keywords

Adherence, compliance, Iron and Folic Acid, postnatal, primigravida, multigravida, Kakamega, Kenya

This article is included in the [KEMRI | Wellcome Trust](#) gateway.

Corresponding author: Felix Bahati (FBahati@kemri-wellcome.org)

Author roles: **Bahati F:** Conceptualization, Data Curation, Formal Analysis, Investigation, Writing – Original Draft Preparation; **Kairu-Wanyoike S:** Conceptualization, Investigation, Methodology, Supervision, Writing – Review & Editing; **Nzioki JM:** Conceptualization, Methodology, Supervision, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome Trust [grant number: 207522/Z/17/Z] awarded to Prof. Michael English. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2021 Bahati F *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Bahati F, Kairu-Wanyoike S and Nzioki JM. **Adherence to iron and folic acid supplementation during pregnancy among postnatal mothers seeking maternal and child healthcare at Kakamega level 5 hospital in Kenya: a cross-sectional study [version 1; peer review: 2 approved with reservations]** Wellcome Open Research 2021, 6:80 <https://doi.org/10.12688/wellcomeopenres.16699.1>

First published: 12 Apr 2021, 6:80 <https://doi.org/10.12688/wellcomeopenres.16699.1>

Introduction

Maternal anaemia remains a public health problem worldwide¹. The World Health Organization (WHO) defines anaemia among pregnant women as having haemoglobin levels of less than 11.0g/dl². Anaemia in pregnant women has been attributed to iron deficiency³. The condition is exacerbated by the high nutrient demand during pregnancy⁴. Globally, 38% of pregnant women have been reported to be anaemic with the highest burden being in Africa¹. The WHO reports that 55.8% of pregnant women in Sub Saharan Africa (SSA) are anaemic¹. In Kenya, it is estimated that 55.1% of pregnant women have iron-deficiency anaemia⁵. A study conducted at Kakamega level 5 hospital noted a 38.9% anaemia prevalence among pregnant women seeking antenatal care (ANC) at the facility⁶.

To reduce incidences of maternal anaemia, the WHO recommends iron and folic acid (IFA) supplementation to all pregnant women⁷. According to WHO guidelines, daily 30mg–60mg of iron and 0.4 mg of folic acid supplements are essential to all pregnant women and their uptake should be commenced as early as possible once pregnancy has been confirmed⁸. However, strict adherence to IFA is required for better outcomes⁹. Evidence suggests that IFA uptake during pregnancy reduces the chances of iron-deficiency anaemia to a great extent¹⁰. Moreover, adequate IFA ensures the wellness of the developing foetus by reducing incidences of neural tube defects, puerperal sepsis, preterm birth, congenital heart defects, and low birth weight^{7,11}.

To ensure equitable access, the government provides IFA supplements to all pregnant women seeking ANC in public health facilities at no cost¹². However, despite all these efforts, adherence to IFA among pregnant women in Kenya remains poor^{12,13}. Although there has been a gradual national upward trend in self-reported IFA adherence from 2.5% in 2010¹⁴ to 8% in 2014, it is clear that the adherence is still low. In western Kenya, where Kakamega level 5 hospital is located, IFA adherence is even lower compared to the national prevalence (6.9%)¹⁵.

The causes of poor adherence to IFA uptake in Kenya are multidimensional. However, inaccessibility to ANC services could be the leading contributor to this poor trend¹⁶. Surprisingly, poor adherence has also been confirmed among pregnant women who regularly visit health facilities for ANC^{9,17,18}. Inadequate knowledge of anaemia, frequent stock-outs of IFA tablets in health facilities, and forgetfulness have been linked as key barriers to IFA supplementation adherence in various studies^{9,19,20}. The side effects resulting from IFA consumption also discourage pregnant mothers from IFA compliance¹³. Although these side effects are common with IFA use, many pregnant women seem to be unaware that practices such as taking the supplements alongside meals or just before going to bed help to alleviate the side effects. The persistently high prevalence of maternal anaemia coupled with poor adherence to IFA supplementation creates a need for more understanding of the possible causes of the poor adherence trend. Therefore, this study sought to investigate the possible determinants of IFA supplementation adherence during pregnancy among postnatal

mothers seeking maternal and child healthcare (MCH) care at Kakamega level 5 hospital.

Methods

Study site

The study was conducted at Kakamega level 5 hospital located in western Kenya. The hospital serves as the main referral public health facility in Kakamega county of Kenya. This facility is located within Kakamega town and has bed capacity of 449, according to the Kenya Master Health facility List (KMHFL) of 2021²¹.

Study design

A cross-sectional study was conducted between May and August 2020. The study targeted postnatal mothers of 0–6 months post-delivery seeking MCH care at Kakamega level 5 hospital. The sample size was calculated based on Cochran formula²² as shown below:

$$n = \frac{Z^2 P(1 - P)}{d^2}$$

To achieve our desired sample size (n), we set the standard normal deviation (Z) at 1.96 with 5% as the level of accuracy (d). A prevalence (P) of 32.7% was used basing on the findings of a similar study conducted in a similar county of Kenya. A total of 339 participants were therefore needed (n) but since our target population of 0–6 months postnatal mothers was less than 10000, the sample size was adjusted as shown²²:

$$nf = \frac{n}{1 + n/N}$$

The approximate total number of women that were anticipated to seek postnatal care (PNC) services during the 3 months of data collection in Kakamega level 5 hospital was attained by multiplying the average monthly PNC attendance by 3. According to the 2019 District Health Information system (DHIS2) data, the average number of women (with children aged 0–6 months) who sought MCH services at this facility every month was 277. We therefore anticipated that 831 mothers would be encountered during the 3 months of data collection (277 X 3=831). Therefore:

$$nf = \frac{339}{1 + (339/831)} = 241$$

We catered for nonresponse by proceeding to another respondent until the required sample size was achieved.

Data collection

Quantitative data – semi-structured questionnaire. Quantitative data were collected at the postnatal section of Kakamega level 5 hospital among postnatal mothers of 15–49 years. We targeted the postnatal mothers who had just delivered in the maternity wards and those who had brought their babies for a routine check-up within six months post-delivery. We were keen to ensure that all the postnatal mothers interviewed were

seeking healthcare services related to MCH care. However, the postnatal mothers who were critically ill such that they wouldn't speak to us comfortably were excluded from the study. An electronic version of a semi-structured questionnaire developed in the Research Electronic Data Capture tool (REDCap) was used to collect the data (see data dictionary codebook for questions in *Underlying data*²³). We scheduled appointments in two sessions every day; the morning session and the afternoon session. About 10 participants were interviewed each day with each of the interviews lasting for approximately 30 minutes. Study participants were recruited once they had obtained the MCH care that they needed.

The purpose of the study was explained to each potential participant after which a written consent was sought. Participants of less than 18 years were only interviewed if they were accompanied by a guardian who provided written consent on their behalf. Every potential study participant who consented to the study was directed to a separate room where a face-to-face interview was conducted. However, interviews with mothers who had just delivered were conducted at their respective bedsides in the ward.

The questionnaire used in the data collection was divided into five sections: socio-demographic characteristics of the participants, health system-related characteristics, participant's knowledge of anaemia, participant's attitude towards ANC healthcare provider and IFA adherence. Adherence to IFA supplementation was established by asking the participants about the average number of IFA tablets taken per week throughout the gestation period. Participant's knowledge of anaemia was ascertained using a set of 10 questions focusing on anaemia causes, haemoglobin level boosting foods, consequences of anaemia and its prevention mechanisms. All the responses provided by the respondents were entered in REDCap²², after which they were uploaded to KEMRI WELLCOME TRUST server every day.

Qualitative data – key informant interviews. The quantitative data were complemented by the key informant interviews (KIIs) with healthcare providers and the community health volunteers (CHVs) attached to the hospital. The first author administered all the KIIs. A total of five healthcare providers purposively selected within the hospital and all the 11 CHVs attached to the hospital participated in the KIIs. Neither of the targeted healthcare providers nor CHVs refused to take part in the study. The health care providers who provided feedback to the KIIs comprised of: PNC in-charge, ANC in-charge, PNC nurse, PNC pharmacist-in-charge and the MCH nutritionist in-charge. The KIIs with the healthcare providers took place at the hospital within their respective departments during individuals' shift. We were flexible in scheduling the KIIs with the healthcare providers and only spoke to each one of them at the time that seemed convenient for them. This was to ensure that the interviews did not compromise their service provision to the patients. The KIIs with CHVs either took place at their respective homes or within Kakamega level 5 hospital. Those who were interviewed at the facility had accompanied a patient to the facility to help them obtain care. We chose to

interview some CHVs at their respective homes to avoid them incurring transport and other costs related to travelling to the facility. The KIIs reported in this study were based on different, but strongly related KII guides prepared to suit either CHVs or healthcare providers (see KII guide in the *Extended data*²³). All the KIIs lasted between 30–45 minutes and were audio-recorded after the participants had given written informed consent to participate. All the interviews were conducted in secluded areas to ensure privacy of the respondents.

Data analysis

During analysis, the quantitative data were exported from the server to a Microsoft Excel CSV file after which a data cleaning process followed.

Adherence to IFA supplementation: This was the main outcome variable in this study. Respondents who took at least five IFA tablets per week throughout their gestation period were classified as adherent, while those who took less than five tablets per week throughout pregnancy were non-adherent.

Knowledge of anaemia: Anaemia knowledge was determined by scoring the responses to the ten questions assessing knowledge level of anaemia. The questions inquired on anaemia causes, consequences and its prevention. Each of the ten questions answered correctly was awarded a score of '1' while a wrong response or admitting not being aware was awarded a score of '0'. The total expected score of anaemia knowledge was 10 and this was used to calculate the percentage score of each participant. The respondents with a score that was equal to or greater than 50% were assumed to have a higher knowledge of anaemia while those with a score below 50% were treated as having low knowledge of anaemia²⁴.

Attitude towards ANC healthcare provider: We also established the attitude of the respondents towards ANC healthcare provider using a Likert scale. Five questions were asked on attitude, with responses ranging from strongly disagree, disagree, neutral, agree and strongly agree. The responses were scored in ascending order, with strongly disagree being scored a '1' while strongly agree was scored a '5'. The scores were summed up and later converted into percentages. Respondents were considered to have a positive attitude towards ANC healthcare provider if they scored at least 70%, while those who scored below 70% were assumed to have a negative attitude towards ANC healthcare provider²⁵.

The cleaned verified data were then imported to R statistical software version 3.5.2 [2018–12–20] for analyses. The respondents' socio-demographic characteristics and the prevalence of IFA compliance were summarized using descriptive statistics such as means, proportions and percentage frequencies. Some variables were also re-grouped into categories based on certain set thresholds. For example, mothers' age was re-categorized into groups, IFA adherence classified as adherent or non-adherent, gravidity as primigravida and multigravida, while the education level of the mother was categorized into primary, secondary and college levels. Equally, the number of ANC visits attained by the mothers were segregated

into two and below ANC visits and at least 3 ANC visits. Associations between adherence to IFA supplementation and its putative determinants were investigated through univariable and multivariable logistic regression. P values of less than 0.05 were interpreted as significant.

Qualitative data analysis began with the transcription of the recorded audio, but we did not return the transcripts to the interviewees for checking and verification. The first and second author conducted a deductive coding to organize the data into themes using the qualitative data analysis software NVivo 12 (QSR International, Australia).

Ethical clearance

Ethical clearance for this study was sought from the Ethical Review Board of Daystar University (approval number: DU-ERB- 000415). Daystar University Ethical Review Board was considered for ethical approval as we were keen to have the study protocol reviewed within the anticipated timelines. The National Commission for Science, Technology and Innovation (NACOSTI) granted the research license. Approvals were also obtained from the hospital in-charge of Kakamega level 5 hospital and the head of the PNC section.

Results

Socio-demographic characteristics of the respondents

A total of 241 mothers seeking MCH care at Kakamega level 5 hospital were interviewed between May and August 2020. The average age of the mothers was 24.9 ± 5.3 years with an age range of 15–43 years. As shown in [Table 1](#), the majority (68.1%) of the respondents belonged in the age category of 19–29 years. Only a few of the respondents had formal employment (19.9%) while 22.4% were students. Most of the mothers interviewed (41.1%) had attained a secondary school level of education. Also, all the respondents admitted to belonging to one religion or another with the majority being Christians (95.9%). More than three-quarters of the mothers (85.5%) resided at a distance greater than 30 minutes to their ANC facilities with motorcycles being the most preferred means of transport (62.2%). A large proportion of the mothers had a high knowledge of anaemia (85.5%). All the respondents admitted to having been given IFA at any one point during ANC and out of these, slightly over half (51.0%) reported having experienced side effects as a result of IFA consumption.

Obstetric related characteristics of the respondents

As shown in [Table 2](#), out of all the respondents interviewed, over two-thirds (73.9%) had completed at most 3 months post-delivery. Almost half of the mothers were multigravida (47.7%) and a few reported having had either a birth complication (12.1%) or miscarriage (4.6%). Most of the respondents (76.3%) had never had a pregnancy counselling session with a health-care provider before getting pregnant. The majority of the mothers (71.8%) sought their first ANC services with a gestation period of more than 8 weeks and many of them attended ANC at least 3 times (92.9%).

Although most of the mothers interviewed in this study had visited the facility to obtain delivery care (33.9%), other MCH

Table 1. Socio-demographic characteristics of the respondents, Kakamega level 5 hospital, Kenya, 2020.

Variable	n	%	95% CI
Mother's age (years)			
≤18	26	10.8	7.3, 15.6
19–29	164	68.1	61.7, 73.9
≥30	51	21.2	16.3, 27.0
Occupation			
Formal	48	19.9	15.2, 25.6
Non-formal	70	29.0	23.5, 35.3
Housewife	69	28.6	23.1, 34.9
Student	54	22.4	17.4, 28.3
Education level			
Primary	48	19.9	15.2, 25.6
Secondary	99	41.1	34.9, 47.6
College	94	39.0	32.9, 45.5
Marital status			
Married	172	71.4	65.1, 76.9
Single	69	28.6	23.1, 34.9
Religion			
Christian	231	95.9	92.3, 97.9
Muslim	10	4.1	2.1, 7.7
Distance to ANC facility			
>30 Minutes	206	85.5	80.2, 89.5
≤30 Minutes	35	14.5	10.5, 19.8
Transport mode to ANC facility			
Walking	73	30.3	24.6, 36.6
Motorcycle	150	62.2	55.8, 68.3
Matatu	18	7.5	4.6, 11.7
Anaemia knowledge			
Low	35	14.5	10.5, 19.8
High	206	85.5	80.2, 89.5
IFA side effects			
No	118	49.0	42.5, 55.4
Yes	123	51.0	44.6, 57.5

services sought included; child immunization (29.5%), child growth monitoring (28.2%), child treatment (6.4%) among others, as shown in [Figure 1](#).

Table 2. Obstetric related characteristics of the studied respondents, Kakamega level 5 hospital, Kenya, 2020.

Variables	n	%	95% CI
Postpartum duration			
0–3 Months	178	73.9	67.8, 79.2
4–6 Months	63	26.1	20.8, 32.3
Gravidity			
Primigravida	126	52.3	45.8, 58.7
Multigravida	115	47.7	41.3, 54.2
Birth complication			
No	212	88.0	83.0, 91.7
Yes	29	12.1	8.3, 17.0
Miscarriage			
No	230	95.4	91.8, 97.6
Yes	11	4.6	2.4, 8.2
Counselling before pregnancy			
No	184	76.3	70.4, 81.5
Yes	57	23.7	18.5, 29.6
First ANC gestation			
≤8 weeks	68	28.2	22.7, 34.4
>8 weeks	173	71.8	65.6, 77.3
ANC visits			
≥3 Times	224	92.9	88.7, 95.7
≤2 Times	17	7.1	4.3, 11.3

Prevalence of IFA adherence

The overall prevalence of IFA adherence among the studied respondents was 60.6% with 146 (146/241) mothers having reported taking at least five IFA tablets per week throughout pregnancy. Out of the IFA non-compliant cohort, slightly more than one third (35.3%) consumed three to four IFA tablets per week on average throughout pregnancy. The highest adherence was observed among the single mothers (79.7%), followed by younger mothers of 18 years and below (76.9%). There was a similar trend of IFA compliance among Muslims and Christians, 60% and 60.6%, respectively. About eight in every 10 of the mothers who never experienced any IFA related side effects (75.9%) were IFA compliant. Less than half (41.7%) of the respondents with a primary education level took IFA as recommended. Mothers who resided within 30 minutes distance from their ANC facilities had a slightly higher adherence (61.2%) as compared to their counterparts residing more than 30 minutes away from their facilities of ANC (57.1%), as shown in [Table 3](#).

A total of 95 (39.4%) respondents were not IFA compliant. Out of this cohort, the majority of the mothers (41.3%) attributed the non-compliance to IFA related side effects. Other common reasons for non-adherence stated by the respondents were: forgetfulness (37.3%), bad smell of the IFA tablets (10.3%) and pharmacophobia (6.3%), as shown in [Table 4](#).

A total of 123 (51%) mothers experienced IFA related side effects. Out of this, almost half (48.0%) were non-compliant. The most prevalent side effects were vomiting (41.3%), nausea (26.9%) and dizziness (18.6%). Slightly over half (52.2%) of the mothers who experienced vomiting while on IFA were non-compliant. About 48.4% of those who felt dizzy due to IFA consumption and the majority of the mothers who experienced IFA related diarrhoea could not comply with IFA supplementation ([Figure 2](#)).

Figure 1. Maternal and child healthcare services sought by respondents, Kakamega level 5 hospital, Kenya, 2020.

Table 3. Prevalence of IFA adherence among the respondents, Kakamega level 5 hospital, Kenya, 2020.

Variables		IFA adherence					
		Average number of IFA tablets taken by respondents per week					
	Total (N)	0-2	%	3-4	%	5-7	%
Mother's age							
≤18	26	0	0	6	23.1	20	76.9
19-29	164	6	3.7	56	34.1	102	62.2
≥30	51	4	7.8	23	45.1	24	47.1
Education level							
Primary	48	1	2.2	27	56.3	20	41.7
Secondary	99	4	4.0	29	29.3	66	66.7
College	94	5	5.3	29	30.9	60	63.8
Marital status							
Married	172	8	4.7	73	42.4	91	52.9
Single	69	2	2.9	12	17.4	55	79.7
Distance to ANC							
>30 mins	35	3	8.6	12	34.3	20	57.1
≤30 mins	206	7	3.4	73	35.4	126	61.2
Side effects							
No	118	4	3.7	32	29.6	82	75.9
Yes	123	6	5.0	53	43.8	64	52.9
Occupation							
Formal	48	3	6.3	14	29.2	31	64.6
Non-formal	70	2	2.9	30	42.9	38	54.3
Housewife	69	3	4.3	27	39.1	39	56.5
Student	54	2	3.7	14	25.9	38	70.4
Religion							
Christian	231	10	4.3	81	35.1	140	60.6
Muslim	10	0	0	4	40.0	6	60.0
Total		10	4.149	85	35.3	146	60.6

Overall Iron and Folic Acid (IFA) prevalence (**60.6%**).**Socio-demographic determinants of IFA adherence**

Investigation of IFA adherence trends through univariable logistic regression was conducted. Specifically, it was noted that postnatal mothers who were at least 30 years of age were less likely to be IFA compliant as compared to their younger counterparts of 18 years and below, (OR=0.267, 95% CI: 0.086, 0.741; p=0.015). Mothers with a primary level of education

had their odds of being IFA compliant reduced by 60% as compared to mothers who had acquired a college level of education, (OR=0.405, 95% CI: 0.196, 0.819; p=0.013). Similarly, the study established reduced odds of IFA adherence among the respondents who had attended ANC for at most two times as well as those who had reported experiencing IFA related side effects, (OR=0.327, 95%CI: 0.109, 0.893; p=0.034)

Table 4. Reasons for non-compliance as stated by the respondents, Kakamega level 5 hospital, Kenya, 2020.

Non-adherence reason	Responses	%	95% CI
Side effects	52	41.3	32.68, 50.40
Forgetfulness	47	37.3	29.00, 46.41
Bad smell	13	10.3	5.83, 17.32
Pharmacophobia	8	6.3	2.98, 12.53
Make the fetus grow bigger	4	3.2	1.02, 8.42
Unaware of IFA importance	1	0.8	0.04, 4.99
Ran out of supplements	1	0.8	0.04, 4.99
Total	126	100

Figure 2. Variations in iron and folic acid (IFA) compliance according to the side effects experienced among respondents, Kakamega level 5 hospital, Kenya, 2020.

and (OR=0.476, 95% CI: 0.279, 0.804; $p=0.006$), respectively. Unlike mothers in marriage, single mothers were 3.5 times more likely to be IFA compliant (OR=3.497, 95% CI: 1.852, 6.971; $p<0.001$). The primigravida mothers were almost four-fold more likely to be IFA compliant as compared to multigravida mothers (OR=3.559, 95% CI: 2.081, 6.187; $p<0.001$). Equally, the respondents who seemed to have higher anaemia knowledge were noted to have elevated odds of adhering to IFA supplementation, (OR=2.676, 95% CI: 1.296, 5.680; $p=0.009$) (Table 5).

In order to understand the independent association between IFA adherence and the socio-demographic factors, all the variables with crude P values of less than 0.2 were fitted to a multivariable model. These included: mother's age, mother's

education level, marital status, gravidity, mother's knowledge of anaemia, number of ANC visits, and IFA related side effects. The multivariable model confirmed the independent correlation between IFA adherence and gravidity (OR=2.704, 95% CI: 1.262, 5.793; $p=0.010$), mother's anaemia knowledge (OR=3.215, 95% CI: 1.346, 7.68; $p=0.009$), number of ANC visits (OR=0.273, 95% CI: 0.081, 0.916; $p=0.036$) and IFA side effects (OR=0.444, 95% CI: 0.246, 0.803; $p=0.007$), as shown in Table 6.

Health system-related determinants of IFA adherence
Univariable analysis was used to identify possible health system related factors influencing IFA supplementation adherence. Mothers who received education on IFA were almost three times more likely to be IFA compliant as compared to those

Table 5. Univariable logistic regression model demonstrating socio-demographic risk factors for IFA adherence, Kakamega level 5 hospital, Kenya, 2020.

Variables	Beta	Std. Error	Z value	OR (95% CI)	P value
Mother's age (Years)					
18	Ref				
19–29	-0.706	0.493	-1.434	0.494 (0.173, 1.23)	0.152
≥30	-1.322	0.544	-2.432	0.267 (0.086, 0.741)	0.015
Education level					
College	Ref				
Secondary	0.125	0.303	0.414	1.133 (0.626, 2.055)	0.679
Primary	-0.905	0.363	-2.491	0.405 (0.196, 0.819)	0.013
Marital status					
Married	Ref				
Single	1.252	0.4336	3.725	3.497 (1.852, 6.971)	<0.001
Gravidity					
Multigravida	Ref				
Primigravida	1.269	0.278	4.575	3.559 (2.081, 6.187)	<0.001
Birth complication					
No	Ref				
Yes	-0.412	0.398	-1.036	0.662 (0.303, 1.457)	0.300
Anaemia knowledge					
Low	Ref				
High	0.984	0.374	2.629	2.676 (1.296, 5.680)	0.009
ANC visits					
≥3	Ref				
≤2	-1.117	0.526	-2.124	0.327 (0.109, 0.893)	0.034
Distance to ANC					
>30 mins	Ref				
≤30 mins	0.167	0.370	0.450	1.181 (0.564, 2.432)	0.653
IFA side effects					
No	Ref				
Yes	-0.742	0.269	-2.754	0.476 (0.279, 0.804)	0.006

Standard Error (Std. Error), odds Ratio (OR), Significant (P<0.05).

who did not receive education on IFA (OR=2.728, 95% CI: 1.297, 5.921; p=0.009). Similarly, attending pregnancy counselling sessions prior to conception played a significant role on IFA adherence. The study revealed that mothers who sought pregnancy counselling sessions were more than twice

likely to be IFA compliant as compared to their counterparts who never attended pregnancy counselling sessions before conception (OR=2.415, 95% CI: 1.262, 4.859; p<0.01). Unlike mothers who had a negative attitude towards ANC healthcare provider, mothers who had a positive attitude towards ANC healthcare

Table 6. Multivariable logistic regression model demonstrating socio-demographic risk factors for IFA adherence, Kakamega level 5 hospital, Kenya, 2020.

Variables	Estimate	Std. Error	Z value	OR (95% CI)	P value
Mother's age (Years)					
≤18	Ref				
19-29	-0.356	0.615	-1.578	0.701 (0.21, 2.339)	0.563
≥30	-0.196	0.747	-0.263	0.822 (0.19, 3.553)	0.793
Education level					
College	Ref				
Secondary	0.295	0.359	0.822	1.344 (0.664, 2.718)	0.411
Primary	-0.481	0.428	-1.125	0.618 (0.267, 1.429)	0.260
Marital status					
Married	Ref				
Single	0.712	0.447	1.592	2.037 (0.848, 4.893)	0.111
Gravidity					
Multigravida	Ref				
Primigravida	0.995	0.389	2.559	2.704 (1.262, 5.793)	0.010
Anaemia knowledge					
Low	Ref				
High	1.168	0.444	2.629	3.215 (1.346, 7.68)	0.009
ANC visits					
≥3	Ref				
≤2	-1.300	0.618	-2.102	0.273 (0.081, 0.916)	0.036
IFA side effects					
No	Ref				
Yes	-0.811	0.302	-2.687	0.444 (0.246, 0.803)	0.007

Standard Error (Std. Error), odds Ratio (OR), Significant (P<0.05)

provider were up to 2.4 times more likely to be IFA compliant (OR=2.357, 95% CI: 1.124, 5.052; p=0.024), as shown in [Table 7](#).

According to the KIIs with CHVs, poor attitude towards ANC healthcare provider is partially contributed by the long turnaround time taken during ANC visits.

“There are still others who prefer enrolling for ANC a bit late say three to four months due in pregnancy. They just want to buy time, especially when they think of the long queue that awaits them at the ANC. They complain that they take a lot of time at ANC...” (KII: CHV).

All the health system related factors that showed a p≤0.2 in the univariable logistic regression were subjected to a multivariable model. Precisely, IFA education, having had a pregnancy counselling session and mother's attitude towards ANC healthcare provider were adjusted for in the multivariable model. Having had a pregnancy counselling session with a healthcare provider before conception and having received education on IFA showed independent association with IFA adherence, (OR=2.086, 95% CI: 1.071, 4.255; p=0.036) and (OR=2.372, 95% CI: 1.109, 5.218; p=0.028) respectively ([Table 8](#)).

The correlation between hospital IFA stockouts and IFA adherence was not statistically significant. However, KIIs with

Table 7. Univariable logistic regression model demonstrating health system-related risk factors for IFA adherence, Kakamega level 5 hospital, Kenya, 2020.

Variables	Beta	Std. Error	Z value	OR (95% CI)	P value
IFA education					
No	Ref				
Yes	1.004	0.384	2.611	2.728 (1.297, 5.921)	0.009
Pregnancy counselling					
No	Ref				
Yes	0.882	0.342	2.581	2.415 (1.262, 4.859)	<0.01
Turnaround Time					
Longer than expected	Ref				
As expected,	0.342	0.321	1.067	1.408 (0.747, 2.640)	0.286
IFA stock out					
No	Ref				
Yes	-0.253	0.400	-0.634	0.776 (0.355, 1.723)	0.526
Attitude towards ANC healthcare provider					
Negative	Ref				
Positive	0.856	0.381	2.253	2.357 (1.124, 5.052)	0.024

Table 8. Multivariable logistic regression model demonstrating health system-related risk factors for IFA adherence, Kakamega level 5 hospital, Kenya, 2020.

Variables	Beta	Std. Error	Z value	AOR (95% CI)	P value
IFA education					
No	Ref				
Yes	0.864	0.392	2.203	2.372 (1.109, 5.218)	0.028
Pregnancy counselling					
No	Ref				
Yes	0.735	0.350	2.102	2.086 (1.071, 4.255)	0.036
Attitude towards ANC healthcare provider					
Negative	Ref				
Positive	0.641	0.400	1.637	1.900 (0.884, 4.150)	0.102

healthcare workers revealed that the facility runs out of the supplements at times, although not very often. Despite the existence of clear channels on how to source for the IFA tablets, it would take a lot of time for the officer in charge to be notified

of the IFA stock-outs. The respondent felt that the situation was exacerbated by delayed communications among the staff and this, to some extent, resulted to missed opportunities for taking IFA.

“...As I said, we need to have a nutritionist in every room but that’s almost impossible for now. So sometimes when I’m compiling monthly reports, I notice that the mothers have not been issued with the supplements for some time. When you make follow-ups, you will be told that IFA had run out of stock. But the problem is that they did not communicate because I could have looked for the supplements somewhere else even if it means borrowing from another facility...” (KII: healthcare provider).

We also learned that the ANC healthcare providers experienced challenges in providing IFA education and general MCH talks to ANC seeking mothers. They attributed this to health system disruptions caused by the COVID-19 pandemic whereby the ANC services were shifted to another department to create more space for COVID 19 patients as indicated in transcripts below.

“Before we came here, we got some space at the eye clinic. But it was even smaller than this. We tried giving health talks there and you would find yourself talking to a group of eye patients, ANC mothers and under-five mothers. It was such a bad confusion and we had to move from there to this orthopaedic department. The space here still doesn’t allow for such talks though. That is as it is for now...” (KII: healthcare provider).

“Did you notice that even the sitting space is not enough? We have no space for more benches. Some mothers are standing as you can see. In fact, this has impacted negatively on our ANC visiting trends. It seems mothers are no longer motivated to come for ANC as they feel that the place is congested to an extent that they miss a place to sit...” (KII: healthcare provider).

As indicated in the following verbatim, healthcare workers at the ANC section experience a heavy workload very often. This is partly due to the limited number of staff as well as the rigorous writing involved in their line of duty.

“I’m the only one here and the two ladies you see over there are interns, the other lady is a casual on the hospital’s payroll. And there is a lot of writing involved by the way. We do not have enough computers and the few that we have keep on hanging...” (KII: healthcare provider).

Discussion

There is moderate adherence to IFA supplementation (60.6%) during pregnancy among mothers seeking MCH care at Kakamega level 5 hospital. The finding is consistent with similar studies conducted in a sub-city of Ethiopia (60.9%), as well as a metropolitan area of Ghana (58.8%)^{26,27}. The IFA supplementation prevalence was however higher as compared to 32.7% reported in Kiambu county of Kenya¹³ and 20.3% among rural communities of North Western Tanzania²⁸, but lower than

71% reported in India²⁹ and 68.6% in Niger³⁰. The moderately higher IFA supplementation prevalence recorded in the current study could be attributed to the fact that the study was conducted in a hospital located within a town. This implies that the hospital is very accessible and the mothers seeking MCH care within the hospital have more access to information as they are likely to be residing closer to town. Nevertheless, the moderate IFA adherence at Kakamega level 5 hospital suggests that pregnant mothers seeking MCH care in this facility and their unborn babies are still exposed to some risks of gestational maternal anaemia. This is because pregnant mothers are expected to comply with daily IFA uptake throughout the gestation period as in accordance with the WHO guidelines⁸.

Participants who experienced IFA related side effects were less likely to be IFA compliant. Equally, most of the respondents who admitted not having adhered to IFA supplementation during their gestation period mentioned side effects (41.3%), forgetfulness (37.3%) and the bad smell of the supplements (10.3%) as the main challenges. Similar reasons for non-adherence have also been reported among pregnant women in India³¹ and Ghana²⁷. In Ethiopia, Nasir *et al.* observed that more than half of the mothers who were IFA non-compliant either attributed the poor trend to IFA related side effects or forgetfulness³². Most mothers do not seem to know how to manage IFA related side-effects and this makes them give up on the supplements quite easily. Out of the 123 mothers who reported to have experienced some side effects due to IFA consumption, almost 90% of them either stopped taking the supplements or let the side effects subside on their own. There is a need for health care providers to include education on the management of IFA related side effects as part of ANC counselling. Practices such as taking IFA alongside meals or just before going to bed as well as eating plenty of vegetables and fruits have been linked to reduced IFA side effects¹³. Forgetfulness among IFA users could be reduced by encouraging mothers to embrace modern technology reminders or take the supplements at specific times, such as after meals, every morning, and just before going to bed. The women could also ask a reliable member of the family or friend to remind them to take the supplements.

Gravidity is a great determinant of IFA adherence. Primigravida mothers were almost three times more likely to be IFA compliant as compared to multigravida respondents. This is in line with other studies conducted in Kenya and India^{13,31}. However, this finding is not consistent with Alemayu *et al.* and Niguse *et al.* who observed higher odds of IFA compliance among multigravida mothers in Ethiopia^{17,33}. The low IFA compliance among the multigravida mothers could be attributed to their experience with delivery and IFA usage. Some studies have shown that women who experience unpleasant side effects with IFA or have a history of good birth outcomes may not appreciate the need to adhere to IFA in their consecutive pregnancies³⁴. Therefore, sensitizing multigravida mothers on the importance of adhering to IFA supplementation during subsequent pregnancies regardless of the previous past experiences is key.

Mothers who attained fewer ANC visits were exposed to IFA non-adherence. Precisely, the respondents who attained a maximum of two ANC visits had up to 70% reduced chances of being IFA compliant. A large population-based study conducted among pregnant women in SSA noted higher odds of IFA compliance among women with at least four ANC visits³⁵. In Ethiopia, Molla *et al.* observed that women with at least four ANC visits were almost seven times more likely to be IFA compliant while Tarekegn *et al.* reported higher IFA compliance among women who had attained at least 3 ANC visits.^{36,37} Generally, ANC visits correlate positively with IFA adherence. A higher number of ANC visits indicates more frequency of contact between the mother and the healthcare provider. This provides a good opportunity for the healthcare provider to encourage the mothers to use the supplements as required. The mothers also get their supplements replenished through such visits to ensure that they do not run out of IFA. Besides, they could share any challenges encountered while taking the supplements with the healthcare provider and be advised accordingly. The WHO recommends at least eight³ ANC visits in pregnancy and therefore encouraging pregnant women to complete the recommended number of visits is likely to have a positive influence on IFA compliance.

Women who have a higher knowledge of anaemia were more than three times likely to be IFA adherent as compared to their counterparts of low anaemia knowledge. Similar findings have also been reported in other studies elsewhere.^{27,36,38} It is possible that having higher knowledge of anaemia enables a mother to understand the aetiology of anaemia, its prevention measures as well as the deleterious effects that the condition could cause to the mother and her unborn baby. This makes the mothers appreciate the importance of taking IFA as recommended.

Pregnancy counselling before conception and ANC education on the importance of IFA adherence are good ways of ensuring compliance with the supplements. This study revealed that women who received education on the importance of IFA during ANC were 2.4 times more likely to be IFA compliant as compared to their counterparts who had no education with regards to IFA supplementation. Just like IFA education, mothers who attended pregnancy counselling sessions before conception had higher odds of IFA compliance. This is consistent with other reports where IFA education during ANC or pregnancy counselling improved the uptake of the supplements^{13,26}. Being educated on the importance of IFA helps mothers to appreciate the need to take the supplements consequently leading to compliance. Unfortunately, KIIs with ANC health care workers revealed that the staff were almost unable to provide this crucial education to ANC mothers as at the time of the study due to health system disruptions caused by the COVID-19 pandemic³⁹ that led to a shift in location of ANC services. Moreover, the understaffing at the ANC department and the rigorous documentation involved within this section barely leaves enough opportunity for individual health talk between a mother and the health care provider. The Ministry of Health should put measures in place to ensure uninterrupted

continued provision of essential health care services such as MCH even in times of pandemics like COVID-19.

Study limitation

The study was likely to have been affected by recall bias as the mothers had to recall their IFA usage trends during their gestation period. However, we minimized this bias by only selecting mothers of 0–6 months post-delivery.

Conclusion

There is moderate adherence to IFA supplementation during pregnancy among mothers seeking MCH at Kakamega level 5 hospital. This suggests that there is still a need for improvement to meet the WHO recommendations where pregnant mothers are encouraged to take IFA supplements every day throughout pregnancy. The greatest impediments of IFA compliance are IFA related side effects, forgetfulness and the bad smell of the IFA tablets. Therefore, education on how to cope up with such anticipated outcomes should be provided to all pregnant mothers during ANC visits.

Data availability

Underlying data

Harvard Dataverse: Replication Data for: Adherence to Iron and Folic Acid supplementation during pregnancy among postnatal mothers seeking Maternal and Child Healthcare at Kakamega level 5 hospital in Kenya, <https://doi.org/10.7910/DVN/ZEYYSZ>²³.

The interview transcripts derived from the KIIs conducted in this study have not been made available for open access because this would violate the privacy that we initially assured all our study participants. Intermediary data in the form of quotes are available throughout the Results section that reflect the data collected. Reasonable access to the interview transcripts may be granted after the application of access permission by downloading and filling [this](#) online form and then sending it to the Data Governing Committee at KEMRI-Wellcome Trust (dgc@kemri-wellcome.org). The applicant must however prove beyond reasonable doubt that the confidentiality of the study participants will not be compromised in any way before data access is granted.

Extended data

Harvard Dataverse: Replication Data for: Adherence to Iron and Folic Acid supplementation during pregnancy among postnatal mothers seeking Maternal and Child Healthcare at Kakamega level 5 hospital in Kenya, <https://doi.org/10.7910/DVN/ZEYYSZ>²³.

This project contains the following extended data:

- KII guide (CHV)
- KII guide (healthcare provider)

Reporting guidelines

Harvard Dataverse: COREQ checklist for 'Adherence to Iron and Folic Acid supplementation during pregnancy among

postnatal mothers seeking Maternal and Child Healthcare at Kakamega level 5 hospital in Kenya', <https://doi.org/10.7910/DVN/ZEYYSZ>²³.

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

References

- WHO: **WHA Global Nutrition Targets 2025: Anaemia Policy Brief**. 2014; (6). [Reference Source](#)
- WHO: **The global prevalence of anaemia in 2011**. 2015. [Reference Source](#)
- WHO: **WHO Recommendations on Antenatal Care for a Positive Pregnancy Experience**. 2018; 10: 1–10. [Reference Source](#)
- WHO: **WHO recommendation on antenatal care contact schedules**. 2018; 1–7.
- MoH: **National Iron and Folic Acid Supplementation Communication Strategy**. 2013.
- Siteti MC, Namasaka SD, Ariya OP, *et al.*: **Anaemia in pregnancy: Prevalence and possible risk factors in Kakamega County, Kenya**. *Science Journal of Public Health*. 2014; 2(3): 216–22. [Publisher Full Text](#)
- WHO: **WHO recommendations on antenatal care for a positive pregnancy experience**. 2016. [Reference Source](#)
- WHO: **Developing and validating an iron and folic acid supplementation indicator for tracking progress towards global nutrition monitoring framework targets**. 2018. [Reference Source](#)
- Kiwanuka TS, Ononge S, Kiondo P, *et al.*: **Adherence to iron supplements among women receiving antenatal care at Mulago National Referral Hospital, Uganda-cross-sectional study**. *BMC Res Notes*. 2017; 10(1): 510. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Daru J, Zamora J, Fernández-félix BM, *et al.*: **Risk of maternal mortality in women with severe anaemia during pregnancy and post partum: a multilevel analysis**. *Lancet Glob Health*. 2018; 6(5): e548–e54. [PubMed Abstract](#) | [Publisher Full Text](#)
- Moreno-fernandez J, Ochoa JJ, Latunde-dada GO, *et al.*: **Iron Deficiency and Iron Homeostasis in Low Birth Weight Preterm Infants: A Systematic Review**. *Nutrients*. 2019; 11(5): 1090. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- USAD: **Barriers to Maternal Iron-Folic Acid Supplementation and Compliance in Kisumu and Migori, Kenya**. 2017. [Reference Source](#)
- Kamau MW, Mirie W, Kimani S: **Compliance with Iron and folic acid supplementation (IFAS) and associated factors among pregnant women: Results from a cross-sectional study in Kiambu County, Kenya**. *BMC Public Health*. 2018; 18(1): 580. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- KDHS: **Kenya Health and Demographic Survey 2008-2009**. 2010; 157–8. [Reference Source](#)
- KDHS: **2014 Demographic and Health Survey Key Findings**. 2014. [Reference Source](#)
- UNICEF: **Approach to Nutrition Programming in the East Asia and Pacific Region 2014 - 2025**. 2014; 3.
- Niguse W, Murugan R: **Determinants of Adherence to Iron Folic Acid Supplementation among Pregnant Women Attending Antenatal Clinic in Asella Town, Ethiopia**. *Int J Ther Appl*. 2018; 35: 60–7. [Reference Source](#)
- Kassa GM, Muche AA, Berhe AK, *et al.*: **Prevalence and determinants of anemia among pregnant women in Ethiopia; a systematic review and meta-analysis**. *BMC Hematol*. 2017; 17: 17. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Agegnehu G, Atenafu A, Dagne H, *et al.*: **Adherence to Iron and Folic Acid Supplement and Its Associated Factors among Antenatal Care Attendant Mothers in Lay Armachiho Health Centers, Northwest, Ethiopia, 2017**. *Int J Reprod Med*. 2019; 2019: 5863737. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gebremariam AD, Tiruneh SA, Abate BA, *et al.*: **Adherence to iron with folic acid supplementation and its associated factors among pregnant women attending antenatal care follow up at Debre Tabor General Hospital, Ethiopia, 2017**. *PLoS One*. 2019; 14(1): e0210086. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- We acknowledge the MCH nurses and CHVs of Kakamega level 5 hospital for creating time to speak to us during KIIs. Special thanks to all the KEMRI Wellcome Trust staff who provided guidance and ideas to make this work successful.
- MoH: **Kenya Master Health Facility List**. [Reference Source](#)
- Cochran WG: **Sampling Techniques third edition**. New York John Wiley Sons. 1977. [Reference Source](#)
- Bahati F, Kairu-Wanyoike S, Nzioki Mativo J: **Replication Data for: Adherence to Iron and Folic Acid supplementation during pregnancy among postnatal mothers seeking Maternal and Child Healthcare at Kakamega level 5 hospital in Kenya**. Harvard Dataverse. 2021; 2: 7910. <http://www.doi.org/10.7910/DVN/ZEYYSZ>
- Ahamed NH, Kotb SAM, Hassanen RH: **Knowledge and Attitude of Pregnant Women about Iron Deficiency Anemia in Assist University Women Health Hospital, Egypt**. *J Nurs Health Sci*. 2018; 7(3): 49–58. [Reference Source](#)
- Kamau M, Mirie W, Kimani S, *et al.*: **Effect of community based health education on knowledge and attitude towards iron and folic acid supplementation among pregnant women in Kiambu County, Kenya: A quasi experimental study**. *PLoS One*. 2019; 14(11): e0224361. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gebreamlak B, Dadi AF, Atnafu A: **High adherence to iron/folic acid supplementation during pregnancy time among antenatal and postnatal care attendant mothers in Governmental Health Centers in Akaki Kality Sub City, Addis Ababa, Ethiopia: Hierarchical negative binomial poisson regression**. *PLoS One*. 2017; 12(1): e0169415. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wemakor A, Garti H, Akai MM, *et al.*: **Prevalence and factors associated with compliance to iron and folic acid supplementation in pregnancy in Tamale Metropolis, Ghana**. *Nutrire*. 2020; 45(2): 16. [Publisher Full Text](#)
- Lyoba WB, Mwakatoga JD, Festo C, *et al.*: **Adherence to Iron-Folic Acid Supplementation and Associated Factors among Pregnant Women in Kasulu Communities in North-Western Tanzania**. *Int J Reprod Med*. 2020; 2020: 3127245. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Manasa K, Chandrakumar SG, Prashantha B: **Assessment of compliance with iron-folic acid therapy during pregnancy among postnatal mothers in a tertiary care centre, Mysuru**. *Int J Community Med Public Heal*. 2019; 6(4): 1665. [Publisher Full Text](#)
- Begum K, Ouédraogo CT, Wessells KR, *et al.*: **Prevalence of and factors associated with antenatal care seeking and adherence to recommended iron-folic acid supplementation among pregnant women in Zinder, Niger**. *Matern Child Nutr*. 2018; 14 Suppl 1(Suppl 1): e12466. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lavanya P, Jayalakshmy R, Rajaa S, *et al.*: **Adherence to iron and folic acid supplementation among antenatal mothers attending a tertiary care center, Puducherry: A mixed-methods study**. *J Family Med Prim Care*. 2020; 9(10): 5205–5211. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nasir BB, Fentie AM, Adisu MK: **Adherence to iron and folic acid supplementation and prevalence of anemia among pregnant women attending antenatal care clinic at Tikur Anbessa Specialized Hospital, Ethiopia**. *PLoS One*. 2020; 15(5): e0232625. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gebremariam AD, Tiruneh SA, Abate BA, *et al.*: **Adherence to iron with folic acid supplementation and its associated factors among pregnant women attending antenatal care follow up at Debre Tabor General Hospital, Ethiopia, 2017**. *PLoS One*. 2019; 14(1): e0210086. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Getachew M, Abay M, Zelalem H, *et al.*: **Magnitude and factors associated with adherence to Iron-folic acid supplementation among pregnant women in Eritrean refugee camps, northern Ethiopia**. *BMC Pregnancy*

Childbirth. 2018; **18**(1): 83.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

35. Ba DM, Ssentongo P, Kjerulff KH, *et al.*: **Adherence to Iron Supplementation in 22 Sub-Saharan African Countries and Associated Factors among Pregnant Women: A Large Population-Based Study**. *Curr Dev Nutr*. 2019; **3**(12): nzz120.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

36. Molla T, Guadu T, Muhammad EA, *et al.*: **Factors associated with adherence to iron folate supplementation among pregnant women in West Dembia district, northwest Ethiopia: A cross sectional study**. *BMC Res Notes*. 2019; **12**(1): 6.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

37. Tarekegn M, Wubshet M, Atenafu A, *et al.*: **Antenatal care and mothers' education improved iron-folic acid adherence at Denbiya district health centers, Northwest Ethiopia: Using pills count method**. *Arch Public Health*. 2019; **77**(1): 30.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

38. Sendeku FW, Azeze GG, Fenta SL: **Adherence to iron-folic acid supplementation among pregnant women in Ethiopia: A systematic review and meta-analysis**. *BMC Pregnancy Childbirth*. 2020; **20**(1): 138.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

39. Adeigbe S: **Where do women birth during a pandemic? Changing perspectives on Safe Motherhood during the COVID-19 pandemic**. *J Glob Heal Sci*. 2020; **2**(1): 4–5.

Open Peer Review

Current Peer Review Status: ? ?

Version 1

Reviewer Report 15 June 2021

<https://doi.org/10.21956/wellcomeopenres.18414.r43854>

© 2021 Mwangi M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Martin N. Mwangi

Training and Research Unit of Excellence (TRUE), College of Medicine, University of Malawi, Blantyre, Malawi

This is a well written paper about a study that sought to investigate the possible determinants of Iron and Folic Acid (IFA) supplementation adherence during pregnancy among postnatal mothers seeking maternal and child healthcare at a referral hospital in Western Kenya. Using a mixed methods approach, the authors explore barriers to IFA uptake among pregnant women.

Introduction

The background information section is well written. Important and recent references on the topic are cited. In paragraph 2, as part of our research in the same Western region of Kenya, the effect of iron supplementation on low birth weight was clearly demonstrated and thus this work deserves mention in relation to the effect of iron supplements on birth weight (see cited article).

Methods

The methods section is well structured and contains most of the relevant information.

1. Adherence to IFA supplementation was the primary outcome and was defined as taking at least five IFA tablets per week throughout the gestation period. The authors do not explain why other less strict definitions such as the WHO definition (iron folate supplements ≥ 90 days or 4 days per week during the pregnancy period) were not sufficient for this population. One wonders to what magnitude the conclusions would change if the criteria stated here (≥ 90 days or 4 days per week) is used.
2. Qualitative data – key informant interviews (KIIs): the KIIs with healthcare providers and the community health volunteers (CHVs) attached to the hospital were well received and accepted by the respondents. What is missing is an explanation of how the authors determined whether all vital information was already collected from a respondent. In addition, how the authors determined the stop point for the KIIs. Did they rely on a saturation point for example?

Results

The results section is clearly structured with adequate tables and figures. Because correlations are not causal the authors should be careful not to ascribe causality to otherwise correlated outcomes.

Discussion

Paragraph one of the discussion reports the main finding of the study. More comparative data on the same outcome from other Kenyan counties could have been reported to aid generalisability.

Paragraph two of the discussion ends with suggestions on how to reduce forgetfulness among IFA users. The authors should include recent developments or strategies such as the use of intravenous iron for example ferric carboxymaltose to combat maternal anaemia in pregnancy.

Bad smell has been noted to be among the top three reasons for non-compliance as reported by the participants of the study. The discussion could explore ways to overcome bad smells, for example in the manufacture of iron gel capsules so that pregnant mothers do not smell or taste the capsules before ingesting them. Other strategies could be outlined as well.

Recall bias is not adequately discussed in the study yet the primary outcome (adherence to IFA supplementation during pregnancy) relied on the ability of participants to recall past events. The recall period is long especially if a participant was recruited in the 3-6 months postpartum period. The paper could benefit from a discussion on strategies applied to reduce the impact of recall bias on the primary outcome.

General edits:

- Introduction, paragraph 3 first sentence should read: "To ensure equitable access, the government **of Kenya** provides..."
- Introduction, last sentence, the word "care" seems to be repeated.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Anaemia and Iron Deficiency Anaemia in pregnancy and early childhood, Nutrition and Infectious Diseases, Nutritional epidemiology, Public Health Nutrition.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 29 Jun 2021

Felix Bahati, KEMRI Wellcome Trust, Nairobi, Kenya

This is a well written paper about a study that sought to investigate the possible determinants of Iron and Folic Acid (IFA) supplementation adherence during pregnancy among postnatal mothers seeking maternal and child healthcare at a referral hospital in Western Kenya. Using a mixed methods approach, the authors explore barriers to IFA uptake among pregnant women.

Introduction

The background information section is well written. Important and recent references on the topic are cited. In paragraph 2, as part of our research in the same Western region of Kenya, the effect of iron supplementation on low birth weight was clearly demonstrated and thus this work deserves mention in relation to the effect of iron supplements on birth weight (see cited article).

We appreciate the reviewer for making us aware about this interesting study. The findings of the interventional study cited by the reviewer in Western Kenya proved that the supplements reduced the risk of low birth weight by 58% as well as prematurity. We have cited this work in paragraph 2 of our introduction to clearly show the significance of IFA supplementation.

Methods

The methods section is well structured and contains most of the relevant information.

1. Adherence to IFA supplementation was the primary outcome and was defined as taking at least five IFA tablets per week throughout the gestation period. The authors do not explain why other less strict definitions such as the WHO definition (iron folate supplements ≥ 90 days or 4 days per week during the pregnancy period) were not sufficient for this population. One wonders to what magnitude the conclusions would change if the criteria stated here (≥ 90 days or 4 days per week) is used.

As the reviewer notes, this study had slightly higher threshold of defining IFA adherence compared to other studies that used a lower threshold. Our adherence definition was motivated by the current WHO guidelines that require pregnant mothers to take IFA tablets daily throughout pregnancy. Unlike the previous guidelines, the current WHO guidelines are stressing on the need for daily intake rather than achieving the previous cut-off of 90 tablets. We reasoned that the daily IFA intake advocated by the WHO throughout gestation coupled with the current ANC guideline of 8 ANC visits from the previous 4, signifies the need for mothers to consume the supplements daily. We therefore designed our study with an aim of defining adherence that

would be in line with the current guidelines which gives it a greater chance of having significant policy implication. We, however, acknowledge the reviewers thought that maybe there would be some significant changes were it that another less strict adherence was used. We have brought out this issue by explaining in our first paragraph of the discussion that probably this could be the reason as to why our adherence prevalence could not match other studies done elsewhere. Besides, we have added a second last paragraph in the discussion section which partly highlights this issue.

1. Qualitative data – key informant interviews (KIIs): the KIIs with healthcare providers and the community health volunteers (CHVs) attached to the hospital were well received and accepted by the respondents. What is missing is an explanation of how the authors determined whether all vital information was already collected from a respondent. In addition, how the authors determined the stop point for the KIIs. Did they rely on a saturation point for example?

Actually, the KII guides used by the interviewer also served as the checklists to ensure that all the relevant information required was discussed during the interview. We conducted KIIs with two groups of respondents in this study; the CHVs and healthcare providers. We conducted KIIs with all the CHVs (11) attached to the hospital. This was because, we knew that most of them came from different sub-ethnicities within the area and therefore this would give us an opportunity to understand deeply the socio-cultural factors affecting IFA adherence (data not presented in this article). As for the healthcare providers, we were keen to cause very minimal inconveniences to care provision and therefore stopped immediately the themes reached the saturation point. We realize that this is a crucial information that we should have provided in the methodology section as the reviewer suggests. We added this information in the methodology section within the 'Qualitative data – key informant interviews' section.

Results

The results section is clearly structured with adequate tables and figures. Because correlations are not causal the authors should be careful not to ascribe causality to otherwise correlated outcomes.

This is a very crucial observation that was also raised by reviewer #1. We have therefore rephrased all the sentences in the discussion section that we thought could be easily misinterpreted to mean causality when in real sense the scope of our analysis was correlation.

Discussion

Paragraph one of the discussion reports the main finding of the study. More comparative data on the same outcome from other Kenyan counties could have been reported to aid generalisability.

Although there is a paucity of data in this area, the authors have identified a few relevant studies conducted in other Kenyan counties and made the necessary comparison. Please see references ...and...

Paragraph two of the discussion ends with suggestions on how to reduce forgetfulness among IFA users. The authors should include recent developments or strategies such as the

use of intravenous iron for example ferric carboxymaltose to combat maternal anaemia in pregnancy.

We commend the reviewer for this insight. We learnt that more than 1000mg doses of Ferric carboxymaltose could be administered to mothers in smaller divided doses administered at an interval on 7 days. This could save the mothers from the daily intake of the IFA tablets. We have included this in paragraph two of our discussion as suggested by the reviewer.

Bad smell has been noted to be among the top three reasons for non-compliance as reported by the participants of the study. The discussion could explore ways to overcome bad smells, for example in the manufacture of iron gel capsules so that pregnant mothers do not smell or taste the capsules before ingesting them. Other strategies could be outlined as well.

Although many participants in our study complained about the bad smell of these tablets, we had not thought of a way of eliminating the bad smell without interfering with the efficacy of the supplements. We think that the adoption of an iron gel capsule as suggested by the reviewer would be helpful in eliminating the smell. We have suggested this in our discussion as recommended by the reviewer. Unfortunately, we couldn't think of other appropriate strategies that could help alleviate this problem but stand ready to cite them if the reviewer suggests such other strategies.

Recall bias is not adequately discussed in the study yet the primary outcome (adherence to IFA supplementation during pregnancy) relied on the ability of participants to recall past events. The recall period is long especially if a participant was recruited in the 3-6 months postpartum period. The paper could benefit from a discussion on strategies applied to reduce the impact of recall bias on the primary outcome.

We have added a whole paragraph in the discussion section to address the recall bias issue. We have cited evidence supporting our recall period of 0-6 months. Generally, pregnancy is a very important event in the lifetime of most women. According to other researchers, when an event is salient, the respondents tend to remember it even 20 months afterwards. This coupled with overall assessment of an event i.e asking the respondent to gauge their overall use of IFA in gestation rather than specific weeks/stages of gestation makes it easier for participants to remember. We are therefore confident that the findings reported in this article are relevant and were not influenced by recall bias to a great extent. We have also edited table 2 and provided the total number of respondents that we interviewed at every monthly interval post-delivery as suggested by the reviewer. The limitation section has also been updated to clearly show the recall period was a weakness in this study.

General edits:

- Introduction, paragraph 3 first sentence should read: "To ensure equitable access, the government **of Kenya** provides..."

This has been rectified

- Introduction, last sentence, the word “care” seems to be repeated.

This has been edited.

- **Is the work clearly and accurately presented and does it cite the current literature?**

Yes

- **Is the study design appropriate and is the work technically sound?**

Yes

- **Are sufficient details of methods and analysis provided to allow replication by others?**

Partly

- **If applicable, is the statistical analysis and its interpretation appropriate?**

Partly

- **Are all the source data underlying the results available to ensure full reproducibility?**

No source data required

- **Are the conclusions drawn adequately supported by the results?**

Yes

Competing Interests

No competing interests were disclosed.

Reviewer Expertise

Anaemia and Iron Deficiency Anaemia in pregnancy and early childhood, Nutrition and Infectious Diseases, Nutritional epidemiology, Public Health Nutrition.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

We very much appreciate Reviewer 2's comments and we believe the paper is better as a result of the changes that we have made. We have made significant changes to the text in the discussion section to reduce the assumption that we are ascribing causality which is far beyond the scope of this study. Further, the limitations section is expanded, and we have edited the text to ensure the reader understands the limitations of the data. We, however, remain confident, that the key points of this article are still entirely valid. Most importantly, the reviewer has helped in spotting areas requiring edits, made some suggestions as possible solution to the problems encountered by poor participants and also reminded us of the study limitations.

Competing Interests: I'm the lead author of this article

Reviewer Report 28 May 2021

<https://doi.org/10.21956/wellcomeopenres.18414.r43773>

© 2021 Dickin K. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Katherine L. Dickin

Master of Public Health Program, Population Medicine and Diagnostic Sciences, Cornell University, Ithaca, NY, USA

The purpose of this research was to identify possible determinants of IFA supplementation adherence during pregnancy at a hospital in Kenya. The study used mixed methods, surveying mothers postnatally and interviewing health care providers on factors reported to influence adherence. This study adds to a body of literature documenting barriers to IFA supplementation across contexts and identifies implications for programs to improve nutritional care during pregnancy. The paper is well-written and clear. Concerns include the lack of clarity on how the primary outcome was defined and collected and the analytic choice to create binary variables, as well as the tendency to ascribe causal relationships to associations.

Specific comments

Introduction

Importantly, the introduction acknowledged multiple factors that contribute to poor “adherence” since this term can imply a lack of cooperation. The authors note other barriers such as stock-outs at health facilities as such their use of “adherence” reflects more than just behavioral lapses on part of women.

Methods

1. The recall period of up to 6 months for postnatal mothers reporting IFA intake seems long. What evidence is there that asking so long after pregnancy did not affect the data? The paper states that many were interviewed within 3 months of delivery, but it would be helpful to clarify how many responded at various intervals and whether any differences were found in data across these time points periods. Appropriately, the possibility of recall bias acknowledged under “study limitations”.
2. A strength of this paper is that the survey data was complemented with key informant interviews (KIIs) with health care providers. In the future, it would be useful to include KIIs with mothers to gather their in-depth perspectives on barriers and facilitators of supplementation.
3. I am not clear on the specifics of how the primary outcome was determined and the questionnaire does not appear to be available. The paper states “adherence to IFA supplementation was established by asking the participants about the average number of IFA tablets taken per week throughout the gestation period.” Was the question open-ended or multiple choice? Were mothers asked about the number of weeks or was it assumed to be “throughout gestation”? Since the majority of women attended ANC after 8 weeks, and likely quite late in pregnancy for some, how was this handled when asking about adherence “throughout gestation”? How did mothers estimate average intake per week? What was

done to guard against social desirability, i.e. mothers answering as they believed was expected of them?

4. The definition of adherence as “at least five IFA tablets per week throughout gestation period” is quite a high bar, given evidence of the effectiveness of even intermittent supplementation and reports that women may adhere better when first experiencing resolution of fatigue and other anemia symptoms or in early stages of supplementation. Consider analyzing adherence as a continuous variable, or at least with multiple levels of adherence.
5. The same concern applies to other variables such as knowledge, attitudes about providers, and even the number of ANC visits attained by the mothers. Converting the data to binary variables results in a loss of information (See Altman *et al.* BMJ. 2006 *The cost of dichotomising continuous variables* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1458573/>)¹ and reduces the power of the analysis. It also makes results hard to interpret, since no evidence is presented on the basis for the cut points, making them appear rather arbitrary.

Results

1. In Table 3, it seems adherence was considered in increments (e.g. 0-2 and 3-4 and 5-7 tabs/day) but without statistical testing of differences. There is also a lack of statistical testing reported on side effects and other compliant/non-compliant comparisons. For example: “A total of 123 (51%) mothers experienced IFA related side effects. Out of this, almost half (48.0%) were non-compliant. The most prevalent side effects were vomiting (41.3%), nausea (26.9%) and dizziness (18.6%). Slightly over half (52.2%) of the mothers who experienced vomiting while on IFA were non-compliant.” There is inadequate analysis provided to support the statement in the discussion that “Participants who experienced IFA related side effects were less likely to be IFA compliant.”
2. The study includes helpful findings on factors related to potential barriers such as distance from the facility. It is notable that most mothers had a high level of knowledge of anemia. Given the concern about stockouts, it would have been good to know how often women were given IFA.
3. There are important program implications from the finding that “mothers who received education on IFA were almost three times more likely to be IFA compliant as compared to those who did not receive education on IFA (OR=2.728, 95% CI: 1.297, 5.921; p=0.009)”. Women’s desire for clear information and counseling on antenatal supplementation has been reported elsewhere (e.g. *Ethiopian women's perspectives on antenatal care and iron-folic acid supplementation: Insights for translating global antenatal calcium guidelines into practice*. Birhanu, *et al.* Maternal & child nutrition 2018 <https://onlinelibrary.wiley.com/doi/pdf/10.1111/mcn.12424>)²
4. Caution is needed in interpreting associations. The paper suggests causality in the statement that “attending pregnancy counselling sessions prior to conception played a significant role on IFA adherence” but this could be due to confounding if, for example, women who are very health-oriented or who have concerns due to reproductive history are more likely to seek counseling prior to pregnancy and also to be more adherent to supplementation.

Discussion

1. While it is useful to compare results to what has been reported in other papers, not all cited papers define compliance in the same way, limiting interpretation. Adherence definitions could be responsible for differences attributed by the authors to accessibility of the hospital, etc. For example, a common definition that was used in the cited Ethiopian and Tanzanian studies is 90 days of consumption, whereas the compliance definition in this analysis would have been over 125.
2. Some interpretations assumed that correlations were causal. There is likely to be collinearity of multiple variables with the number of ANC visits (which provide more info, more supplements, potentially more support, and indicate time and access to services), for example. Associations explored in this paper are a step toward understanding supplementation but there is a need to recognize underlying factors related to women's mental health, autonomy, workloads, mobility, and social support, as well as quality and accessibility of health services.
3. **Study limitations:** consider also noting the risk of social desirability of responses, given quite high levels of reported adherence, and the difficulty in estimating "on average" intake. Also the limits on drawing conclusions about causality from the associations found in this analysis.

Conclusions are reasonable, despite some over-interpretation of causality in the discussion.

References

1. Altman DG, Royston P: The cost of dichotomising continuous variables. *BMJ*. 2006; **332** (7549): 1080 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Birhanu Z, Chapleau GM, Ortolano SE, Mamo G, et al.: Ethiopian women's perspectives on antenatal care and iron-folic acid supplementation: Insights for translating global antenatal calcium guidelines into practice. *Matern Child Nutr*. **14 Suppl 1**. [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Maternal and child nutrition, social and behavior change, qualitative and mixed methods research, nutrition program implementation and effectiveness

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 29 Jun 2021

Felix Bahati, KEMRI Wellcome Trust, Nairobi, Kenya

The purpose of this research was to identify possible determinants of IFA supplementation adherence during pregnancy at a hospital in Kenya. The study used mixed methods, surveying mothers postnatally and interviewing health care providers on factors reported to influence adherence. This study adds to a body of literature documenting barriers to IFA supplementation across contexts and identifies implications for programs to improve nutritional care during pregnancy. The paper is well-written and clear. Concerns include the lack of clarity on how the primary outcome was defined and collected and the analytic choice to create binary variables, as well as the tendency to ascribe causal relationships to associations.

Specific comments

Introduction

Importantly, the introduction acknowledged multiple factors that contribute to poor “adherence” since this term can imply a lack of cooperation. The authors note other barriers such as stock-outs at health facilities as such their use of “adherence” reflects more than just behavioral lapses on part of women.

We appreciate that we had used the term ‘adherence’ to imply that mothers are not taking the supplements as required without acknowledging the fact that some of the reasons behind such non-adherence trends are way beyond their control. A good example is the IFA stock-out within facilities as pointed out by the reviewer, something that these mothers may not have control about. We have therefore edited the fourth paragraph of our introduction to include a statement clearly stating that some of these causes of poor adherence are beyond the control of these women.

Methods

1. The recall period of up to 6 months for postnatal mothers reporting IFA intake seems long. What evidence is there that asking so long after pregnancy did not affect the data? The paper states that many were interviewed within 3 months of delivery, but it would be helpful to clarify how many responded at various intervals and whether any differences were found in data across these time points periods. Appropriately, the

possibility of recall bias acknowledged under “study limitations”.

We have added a whole paragraph in the discussion section to address the recall bias issue. We have cited evidence supporting our recall period of 0-6 months. Generally, pregnancy is a very important event in the lifetime of most women. According to other researchers, when an event is salient, the respondents tend to remember it even 20 months afterwards. This coupled with overall assessment of an event i.e asking the respondent to gauge their overall use of IFA in gestation rather than specific weeks/stages of gestation makes it easier for participants to remember. We are therefore confident that the findings reported in this article are relevant and were not influenced by recall bias to a great extent. We have also edited table 2 and provided the total number of respondents that we interviewed at every monthly interval post-delivery as suggested by the reviewer. The limitation section has also been updated to clearly show the recall period was a weakness in this study.

1. A strength of this paper is that the survey data was complemented with key informant interviews (KIIs) with health care providers. In the future, it would be useful to include KIIs with mothers to gather their in-depth perspectives on barriers and facilitators of supplementation.

We thank the reviewer for this useful direction. This is a good suggestion that could help us unmask more barriers and facilitators of IFA supplementation that we may not have achieved with the semi-structured questionnaire that was used in the current study. We shall consider this in our future studies.

1. I am not clear on the specifics of how the primary outcome was determined and the questionnaire does not appear to be available. The paper states “adherence to IFA supplementation was established by asking the participants about the average number of IFA tablets taken per week throughout the gestation period.” Was the question open-ended or multiple choice? Were mothers asked about the number of weeks or was it assumed to be “throughout gestation”? Since the majority of women attended ANC after 8 weeks, and likely quite late in pregnancy for some, how was this handled when asking about adherence “throughout gestation”? How did mothers estimate average intake per week? What was done to guard against social desirability, i.e. mothers answering as they believed was expected of them?

We measured adherence to IFA supplementation during pregnancy using a set of two questions. Firstly, we asked the respondent whether they took the IFA tablets faithfully throughout pregnancy as recommended by the healthcare provider, “Did you take the IFA supplements faithfully as recommended by the health care provider? (Yes/No)”. This was confirmed further by asking about the average number of IFA tablets that the respondents consumed per week throughout the gestation, “On average, how many IFA tablets did you used to take every week throughout pregnancy? (open ended).” Although we had assumed taking the IFA tablets faithfully as recommended to mean taking IFA daily as WHO recommends, we noticed that some respondents who reported taking the tablets faithfully also reported taking as low as three IFA tablets per week on average. For the sake of standardization of our primary outcome, we defined adherence as having taken at least five IFA tablets per week throughout pregnancy (Similar to having used the supplements for an average of at least 5 days per week throughout the gestation). Although the respondents were classified as either adherent or non-adherent based on the number of tablets consumed, we did not classify respondents who started ANC less than 3 months due to delivery as compliant. This was because we reasoned that they had consumed the

supplements for a shorter duration compared to others. Nonetheless, the authors acknowledge that by measuring adherence throughout gestation without specifying weeks, we failed to establish a specific interval over which adherence was measured. We have further acknowledged this in the limitation section. All in all, it would have been difficult for the respondents to remember their IFA use patterns had we specified some gestation weeks. We avoided instances where the participants could provide inaccurate responses for social desirability by: 1) Explaining the justification of the study that we only intended to use the data to influence policies meant to improve IFA uptake in pregnancy and avoid pregnancy complications and 2). Assuring the participants of their confidentiality such that they were not going to be victimised in any way based on their responses. Further, we sought permission from the hospital management to allow our interviewer to interact with the mothers within the clinics without wearing a medical regalia such as a white coat to avoid creating any impression that he was a healthcare provider consequently risking responses of social desirability.

1. The definition of adherence as “at least five IFA tablets per week throughout gestation period” is quite a high bar, given evidence of the effectiveness of even intermittent supplementation and reports that women may adhere better when first experiencing resolution of fatigue and other anemia symptoms or in early stages of supplementation. Consider analyzing adherence as a continuous variable, or at least with multiple levels of adherence.

We understand the point that the reviewer is trying to put across here. Taking an average of five tablets of IFA per week may seem quite higher. We have added a small section in the discussion section to highlight this and make our readers aware. We realized that we based much of this study on the current WHO guidelines of IFA supplementation and tried as much as possible to stick to their recommendation of adherence as daily intake of IFA right from conception to birth. This might have made us set the bar quite high. Our reasoning was also informed by other studies conducted in other parts of Kenya where more than 4 IFA tablets per week were used as a cut off for adherence. Despite such studies having focused a specific week among ANC seeking mothers, the fact remains that the assumption is that the mothers will stick to the same trend throughout pregnancy. We are confident that our findings suit well with the current WHO IFA supplementation guidelines and can easily inform policies within this context. The reviewer's preference on the use of adherence as a continuous variable is a keen observation and justified. We overlooked this during our analysis, and fully concur with the reviewer that it would have been better to use adherence as a continuous variable or establish various levels of adherence. This is a weakness in our data analysis approaches and we have stated it in the limitations. We believe that our table 3 highlights some various levels of adherence to some extent.

1. The same concern applies to other variables such as knowledge, attitudes about providers, and even the number of ANC visits attained by the mothers. Converting the data to binary variables results in a loss of information (See Altman et al. BMJ. 2006 *The cost of dichotomising continuous variables* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1458573/>)¹ and reduces the power of the analysis. It also makes results hard to interpret, since no evidence is presented on the basis for the cut points, making them appear rather arbitrary.

Actually, the reviewer made us realize that by converting the continuous variables into binary we sacrificed clarity and the statistical rigor in favour of brevity. As already stated above, we have discussed this as a limitation in the limitation section.

Results

1. In Table 3, it seems adherence was considered in increments (e.g. 0-2 and 3-4 and 5-7 tabs/day) but without statistical testing of differences. There is also a lack of statistical testing reported on side effects and other compliant/non-compliant comparisons. For example: "A total of 123 (51%) mothers experienced IFA related side effects. Out of this, almost half (48.0%) were non-compliant. The most prevalent side effects were vomiting (41.3%), nausea (26.9%) and dizziness (18.6%). Slightly over half (52.2%) of the mothers who experienced vomiting while on IFA were non-compliant." There is inadequate analysis provided to support the statement in the discussion that "Participants who experienced IFA related side effects were less likely to be IFA compliant."

"Participants who experienced IFA related side effects were less likely to be IFA compliant" was based on the odds ratios obtained from the multivariable logistic regression presented in table 6 and not the descriptive statistics as the reviewer depicts.

1. The study includes helpful findings on factors related to potential barriers such as distance from the facility. It is notable that most mothers had a high level of knowledge of anemia. Given the concern about stockouts, it would have been good to know how often women were given IFA.
 - *We did not collect this information directly as stated by the reviewer. This information was collected indirectly by asking the health care providers about IFA stock-outs. As one of the health care providers noted during the KII, "... As I said, we need to have a nutritionist in every room but that's almost impossible for now. So sometimes when I'm compiling monthly reports, I notice that the mothers have not been issued with the supplements for some time. When you make follow-ups, you will be told that IFA had run out of stock. But the problem is that they did not communicate because I could have looked for the supplements somewhere else even if it means borrowing from another facility..." (KII: healthcare provider). We also asked the mothers whether there was a time that they visited ANC and were not issued with supplements due to stock outs. Only 12 % (n=29) of the mothers said that they failed to get IFA tablets during ANC due to stock outs of the commodity at the facility. In almost all the 12% of the cases, this only happened once. We have included the sentences below in the results section to show how often the participants were given the supplements. "Only 12% of the participants interviewed reported not having been given IFA supplements during ANC visits because the facility had run out of stock. However, this only happened once in almost all the 12% of the cases. Separately, KIIs with healthcare workers also revealed that the facility runs out of the supplements at times, although not very often."*
1. There are important program implications from the finding that "mothers who received education on IFA were almost three times more likely to be IFA compliant as compared to those who did not receive education on IFA (OR=2.728, 95% CI: 1.297, 5.921; p=0.009)". Women's desire for clear information and counseling on antenatal supplementation has been reported elsewhere (e.g. *Ethiopian women's perspectives on antenatal care and iron-folic acid supplementation: Insights for translating global antenatal calcium guidelines into practice*. Birhanu, et al. Maternal & child nutrition 2018 <https://onlinelibrary.wiley.com/doi/pdf/10.1111/mcn.12424>)²

We have read the suggested work and even cited it in the discussion (ref no..) to make our discussion even better.

1. Caution is needed in interpreting associations. The paper suggests causality in the

statement that “attending pregnancy counselling sessions prior to conception played a significant role on IFA adherence” but this could be due to confounding if, for example, women who are very health-oriented or who have concerns due to reproductive history are more likely to seek counseling prior to pregnancy and also to be more adherent to supplementation.

We acknowledge that the statement highlighted by the reviewer could easily be misinterpreted to mean that we are ascribing ANC attendance to causality which is wrong. We have rephrased the sentence to imply an association between ANC counselling and adherence.

Discussion

1. While it is useful to compare results to what has been reported in other papers, not all cited papers define compliance in the same way, limiting interpretation. Adherence definitions could be responsible for differences attributed by the authors to accessibility of the hospital, etc. For example, a common definition that was used in the cited Ethiopian and Tanzanian studies is 90 days of consumption, whereas the compliance definition in this analysis would have been over 125.

The reviewer's thought on the possible reason for the disparities in IFA adherence prevalence between our study and other similar studies conducted elsewhere is indeed more apt than what we thought could be due to hospital accessibility. We have therefore re-adjusted the first paragraph to include the reviewer's thought.

1. Some interpretations assumed that correlations were causal. There is likely to be collinearity of multiple variables with the number of ANC visits (which provide more info, more supplements, potentially more support, and indicate time and access to services), for example. Associations explored in this paper are a step toward understanding supplementation but there is a need to recognize underlying factors related to women's mental health, autonomy, workloads, mobility, and social support, as well as quality and accessibility of health services.

The scope of this study was to establish the associations and not to ascribe causality. We have therefore rephrased all the sentences that seemed to portray causality rather than association. Besides, we have provided a section in the discussion clearly stating that the association revealed in this study are subject to collinearity as elaborated by the reviewer.

1. **Study limitations:** consider also noting the risk of social desirability of responses, given quite high levels of reported adherence, and the difficulty in estimating “on average” intake. Also the limits on drawing conclusions about causality from the associations found in this analysis.

All the observation raised here by the reviewer have been considered partly in the discussion and fully in the limitation section.

Conclusions are reasonable, despite some over-interpretation of causality in the discussion.

- **Is the work clearly and accurately presented and does it cite the current literature?**

Yes

- **Is the study design appropriate and is the work technically sound?**

Yes

- **Are sufficient details of methods and analysis provided to allow replication by others?**

Partly

- **If applicable, is the statistical analysis and its interpretation appropriate?**

Partly

- **Are all the source data underlying the results available to ensure full reproducibility?**

No source data required

- **Are the conclusions drawn adequately supported by the results?**

Partly

Generally, we very much appreciate Reviewer 1's comments and we believe the paper is stronger and more precise as a result of the changes we have made in response. We have made significant changes to the text to reduce assumptions/causality misinterpretations and instead speak to the data we have. Further, the limitations section is expanded, and we have edited the text to ensure the reader understands the limitations of the study. We believe, however, that the key points of the text are still entirely valid.

Competing Interests: I'm the lead author of this article